arXiv:1001.0030v1 [math.CO] 30 Dec 2009

CYCLIC SIEVING FOR GENERALISED NON-CROSSING PARTITIONS ASSOCIATED TO COMPLEX REFLECTION GROUPS OF EXCEPTIONAL TYPE — THE DETAILS

Christian Krattenthaler[†] and Thomas W. Müller[‡]

[†] Fakultät für Mathematik, Universität Wien,
Nordbergstraße 15, A-1090 Vienna, Austria.
WWW: `http://www.mat.univie.ac.at/~kratt`

[‡] School of Mathematical Sciences,
Queen Mary & Westfield College, University of London,
Mile End Road, London E1 4NS, United Kingdom.
WWW: `http://www.maths.qmw.ac.uk/~twm/`

Abstract. We prove that the generalised non-crossing partitions associated to well-generated complex reflection groups of exceptional type obey two different cyclic sieving phenomena, as conjectured by Armstrong, respectively by Bessis and Reiner. This manuscript accompanies the paper *"Cyclic sieving for generalised non-crossing partitions associated to complex reflection groups of exceptional type"* [arχiv:1001.0028], for which it provides the computational details.

1. Introduction

In his memoir [1], Armstrong introduced *generalised non-crossing partitions* associated to finite (real) reflection groups, thereby embedding Kreweras' non-crossing partitions [19], Edelman's m-divisible non-crossing partitions [10], the non-crossing partitions associated to reflection groups due to Bessis [4] and Brady and Watt [8] into one uniform framework. Bessis and Reiner [7] observed that Armstrong's definition can be straightforwardly extended to *well-generated complex reflection groups* (see Section 2 for the precise definition). These generalised non-crossing partitions possess a wealth of beautiful properties, and they display deep and surprising relations to other combinatorial objects defined for reflection groups (such as the generalised cluster complex

---

2000 *Mathematics Subject Classification.* Primary 05E15; Secondary 05A10 05A15 05A18 06A07 20F55.

Key words and phrases. complex reflection groups, unitary reflection groups, m-divisible non-crossing partitions, generalised non-crossing partitions, Fuß–Catalan numbers, cyclic sieving.

[†]Research partially supported by the Austrian Science Foundation FWF, grants Z130-N13 and S9607-N13, the latter in the framework of the National Research Network "Analytic Combinatorics and Probabilistic Number Theory."

[‡]Research supported by the Austrian Science Foundation FWF, Lise Meitner grant M1201-N13.

of Fomin and Reading [11], or the extended Shi arrangement respectively the geometric multichains of filters of Athanasiadis [2, 3]); see Armstrong's memoir [1] and the references given therein.

On the other hand, *cyclic sieving* is a phenomenon brought to light by Reiner, Stanton and White [23]. It extends the so-called "(-1)-phenomenon" of Stembridge [27, 28]. Cyclic sieving can be defined in three equivalent ways (cf. [23, Prop. 2.1]). The one which gives the name can be described as follows: given a set S of combinatorial objects, an action on S of a cyclic group $G = \langle g \rangle$ with generator g of order n, and a polynomial $P(q)$ in q, we say that the triple (S, P, G) *exhibits the cyclic sieving phenomenon*, if the number of elements of S fixed by g^k equals $P(e^{2\pi ik/n})$. In [23] it is shown that this phenomenon occurs in surprisingly many contexts, and several further instances have been discovered since then.

In [1, Conj. 5.4.7] ([7, Conj. 6.4]) and [7, Conj. 6.5], Armstrong, respectively Bessis and Reiner conjecture that generalised non-crossing partitions for irreducible well-generated complex reflection groups exhibit two different cyclic sieving phenomena (see Sections 3 and 6 for the precise statements).

According to the classification of these groups due to Shephard and Todd [25], there are two infinite families of irreducible well-generated complex reflection groups, namely the groups $G(d, 1, n)$ and $G(e, e, n)$, where n, d, e are positive integers, and there are 26 exceptional groups. For the infinite families of types $G(d, 1, n)$ and $G(e, e, n)$, the two cyclic sieving conjectures follow from the results in [16].

The purpose of the present article is to prove the cyclic sieving conjectures of Armstrong, respectively of Bessis and Reiner for the 26 exceptional types, thus completing the proof of these conjectures. Since the generalised non-crossing partitions feature a parameter m, from the outset this is *not* a finite problem. Consequently, we first need several auxiliary results to reduce the conjectures for each of the 26 exceptional types to a *finite* problem. Subsequently, we use Stembridge's *Maple* package `coxeter` [29] and the *GAP* package `CHEVIE` [12] to carry out the remaining *finite* computations. It is interesting to observe that, for the verification of the type E_8 case, it is essential to use the decomposition numbers in the sense of [14, 15, 17] because, otherwise, the necessary computations would not be feasible in reasonable time with the currently available computer facilities. We point out that, for the special case where the aforementioned parameter m is equal to 1, the first cyclic sieving conjecture has been proved in a uniform fashion by Bessis and Reiner in [7]. The crucial result on which their proof is based is (4.5) below, and it plays an important role in our reduction of the conjectures for the 26 exceptional groups to a finite problem.

Our paper is organised as follows. In the next section, we recall the definition of generalised non-crossing partitions for well-generated complex reflection groups and of decomposition numbers in the sense of [14, 15, 17], and we review some basic facts. The first cyclic sieving conjecture is subsequently stated in Section 3. Section 4 contains the announced auxiliary lemmas which, for the 26 exceptional types, allow a reduction of the conjecture to a finite problem. The remaining case-by-case verification of the conjecture is then carried out in Section 5. The second cyclic sieving conjecture is stated in Section 6. Section 7 contains the auxiliary lemmas which, for the 26 exceptional types, allow a reduction of the conjecture to a finite problem, while Section 8 contains the remaining case-by-case verification of the conjecture.

2. Preliminaries

A *complex reflection group* is a group generated by (complex) reflections in $\mathbb{C}^n$. (Here, a reflection is a non-trivial element of $GL(\mathbb{C}^n)$ which fixes a hyperplane pointwise and which has finite order.) We refer to [21] for an in-depth exposition of the theory complex reflection groups.

Shephard and Todd provided a complete classification of all *finite* complex reflection groups in [25] (see also [21, Ch. 8]). According to this classification, an arbitrary complex reflection group W decomposes into a direct product of *irreducible* complex reflection groups, acting on mutually orthogonal subspaces of the complex vector space on which W is acting. Moreover, the list of irreducible complex reflection groups consists of the infinite family of groups $G(m,p,n)$, where m,p,n are positive integers, and 34 exceptional groups, denoted $G_4, G_5, \dots, G_{37}$ by Shephard and Todd.

In this paper, we are only interested in finite complex reflection groups which are *well-generated*. A complex reflection group of rank n is called *well-generated* if it is generated by n reflections.[1] Well-generation can be equivalently characterised by a duality property due to Orlik and Solomon [22]. Namely, a complex reflection group of rank n has two sets of distinguished integers $d_1 \le d_2 \le \cdots \le d_n$ and $d_1^* \ge d_2^* \ge \cdots \ge d_n^*$, called its *degrees* and *codegrees*, respectively (see [21, p. 51 and Def. 10.27]). Orlik and Solomon observed, using case-by-case checking, that an irreducible complex reflection group W of rank n is well-generated if and only if its degrees and codegrees satisfy

$$d_i + d_i^* = d_n$$

for all $i = 1, 2, \dots, n$. The reader is referred to [21, App. D.2] for a table of the degrees and codegrees of all irreducible complex reflection groups. Together with the classification of Shephard and Todd [25], this constitutes a classification of well-generated complex reflection groups: the irreducible well-generated complex reflection groups are

- — the two infinite families $G(d,1,n)$ and $G(e,e,n)$, where d,e,n are positive integers,
- — the exceptional groups $G_4, G_5, G_6, G_8, G_9, G_{10}, G_{14}, G_{16}, G_{17}, G_{18}, G_{20}, G_{21}$ of rank 2,
- — the exceptional groups $G_{23} = H_3, G_{24}, G_{25}, G_{26}, G_{27}$ of rank 3,
- — the exceptional groups $G_{28} = F_4, G_{29}, G_{30} = H_4, G_{32}$ of rank 4,
- — the exceptional group G_{33} of rank 5,
- — the exceptional groups $G_{34}, G_{35} = E_6$ of rank 6,
- — the exceptional group $G_{36} = E_7$ of rank 7,
- — and the exceptional group $G_{37} = E_8$ of rank 8.

In this list, we have made visible the groups $H_3, F_4, H_4, E_6, E_7, E_8$ which appear as exceptional groups in the classification of all irreducible *real* reflection groups (cf. [13]).

Let W be a well-generated complex reflection group of rank n, and let $T \subseteq W$ denote the set of *all* (complex) reflections in the group. Let $\ell_T : W \to \mathbb{Z}$ denote the word length in terms of the generators T. This word length is called *absolute length* or *reflection length*. Furthermore, we define a partial order $\le_T$ on W by

$$u \le_T w \quad \text{if and only if} \quad \ell_T(w) = \ell_T(u) + \ell_T(u^{-1}w). \tag{2.1}$$

---

[1] We refer to [21, Def. 1.29] for the precise definition of "rank." Roughly speaking, the rank of a complex reflection group W is the minimal n such that W can be realized as reflection group on $\mathbb{C}^n$.

This partial order is called *absolute order* or *reflection order.* As is well-known and easy to see, the relation (2.1) is equivalent to the statement that every shortest representation of u by reflections occurs as an initial segment in some shortest product representation of w by reflections.

Now fix a (generalised) Coxeter element[2] $c \in W$ and a positive integer m. The *m-divisible non-crossing partitions* $NC^m(W)$ are defined as the set

$$NC^m(W) = \big\{(w_0; w_1, \dots, w_m) : w_0 w_1 \cdots w_m = c \text{ and } \\ \ell_T(w_0) + \ell_T(w_1) + \cdots + \ell_T(w_m) = \ell_T(c)\big\}.$$

We have suppressed the dependence on c, since we understand this definition up to conjugation. To be more precise, it can be shown that any two Coxeter elements are conjugate to each other (see [26, Theorem 4.2] or [21, Cor.11.25]), and hence the resulting sets $NC^m(W)$ are conjugate to each other. If $m = 1$, then $NC^1(W)$ can be identified with the set $NC(W)$ of non-crossing partitions for the (complex) reflection group W as defined by Bessis and Corran (cf. [6] and [5, Sec. 13]; their definition extends the earlier definition by Bessis [4] and Brady and Watt [8] for real reflection groups).

The following result has been proved by a collaborative effort of several authors (see [5, Prop. 13.1]).

Theorem 1. *Let W be an irreducible well-generated complex reflection group, and let $d_1 \le d_2 \le \cdots \le d_n$ be its degrees and $h := d_n$ its Coxeter number. Then*

$$|NC^m(W)| = \prod_{i=1}^{n} \frac{mh + d_i}{d_i}. \tag{2.2}$$

Remark 1. (1) The number in (2.2) is called the *Fuß–Catalan number* for the reflection group W.

(2) If c is a Coxeter element of a well-generated complex reflection group W of rank n, then $\ell_T(c) = n$. (This follows from [5, Sec. 7].)

We conclude this section by recalling the definition of decomposition numbers from [14, 15, 17]. Although we need them here only for (very small) real reflection groups, and although, strictly speaking, they have been only defined for real reflection groups in [14, 15, 17], this definition can be extended to well-generated complex reflection groups without any extra effort, which we do now.

Given a well-generated complex reflection group W of rank n, types $T_1, T_2, \dots, T_d$ (in the sense of the classification of well-generated complex reflection groups) such that the sum of the ranks of the T_i's equals n, and a Coxeter element c, the *decomposition number*

---

[2] An element of an irreducible well-generated complex reflection group W of rank n is called a *Coxeter element* if it is *regular* in the sense of Springer [26] (see also [21, Def. 11.21]) and of order d_n. An element of W is called regular if it has an eigenvector which lies in no reflecting hyperplane of a reflection of W. It follows from an observation of Lehrer and Springer, proved uniformly by Lehrer and Michel [20] (see [21, Theorem 11.28]), that there is always a regular element of order d_n in an irreducible well-generated complex reflection group W of rank n. More generally, if a well-generated complex reflection group W decomposes as $W \cong W_1 \times W_2 \times \cdots \times W_k$, where the W_i's are irreducible, then a Coxeter element of W is an element of the form $c = c_1 c_2 \cdots c_k$, where c_i is a Coxeter element of W_i, $i = 1, 2, \dots, k$. If W is a *real* reflection group, that is, if all generators in T have order 2, then the notion of generalised Coxeter element given above reduces to that of a Coxeter element in the classical sense (cf. [13, Sec. 3.16]).

$N_W(T_1, T_2, \dots, T_d)$ is defined as the number of "minimal" factorisations $c = c_1 c_2 \cdots c_d$, "minimal" meaning that $\ell_T(c_1) + \ell_T(c_2) + \cdots + \ell_T(c_d) = \ell_T(c) = n$, such that, for $i = 1, 2, \dots, d$, the type of c_i as a parabolic Coxeter element is T_i. (Here, the term "parabolic Coxeter element" means a Coxeter element in some parabolic subgroup. It follows from [24, Prop.6.3] that any element c_i is indeed a Coxeter element in a parabolic subgroup of W. By definition, the type of c_i is the type of this parabolic subgroup.) Since any two Coxeter elements are conjugate to each other, the decomposition numbers are independent of the choice of the Coxeter element c.

The decomposition numbers for real reflection groups have been computed in [14, 15, 17]. To compute the decomposition numbers for well-generated complex reflection groups is a task that remains to be done.

3. Cyclic sieving I

In this section we present the first cyclic sieving conjecture due to Armstrong [1, Conj. 5.4.7], and to Bessis and Reiner [7, Conj. 6.4].

Let $\phi : NC^m(W) \to NC^m(W)$ be the map defined by

$$(w_0; w_1, \dots, w_m) \mapsto \big((cw_mc^{-1})w_0(cw_mc^{-1})^{-1}; cw_mc^{-1}, w_1, w_2, \dots, w_{m-1}\big). \tag{3.1}$$

For $m = 1$, this action reduces to conjugation by the Coxeter element c (applied to w_1). Cyclic sieving arising from conjugation by c has been the subject of [7].

It is easy to see that ϕ^{mh} acts as the identity, where h is the Coxeter number of W (see (4.1) below). By slight abuse of notation, let C_1 be the cyclic group of order mh generated by ϕ. (The slight abuse consists in the fact that we insist on C_1 to be a cyclic group of order mh, while it may happen that the order of the action of ϕ given in (3.1) is actually a proper divisor of mh.)

Given these definitions, we are now in the position to state the first cyclic sieving conjecture of Armstrong, respectively of Bessis and Reiner. By the results of [16] and of this paper, it becomes the following theorem.

Theorem 2. *For an irreducible well-generated complex reflection group W and any $m \ge 1$, the triple $(NC^m(W), \operatorname{Cat}^m(W;q), C_1)$, where $\operatorname{Cat}^m(W;q)$ is the q-analogue of the Fuß–Catalan number defined by*

$$\operatorname{Cat}^m(W;q) := \prod_{i=1}^{n} \frac{[mh+d_i]_q}{[d_i]_q}, \tag{3.2}$$

exhibits the cyclic sieving phenomenon in the sense of Reiner, Stanton and White [23]*. Here, n is the rank of W, $d_1, d_2, \dots, d_n$ are the degrees of W, h is the Coxeter number of W, and $[\alpha]_q := (1-q^\alpha)/(1-q)$.*

Remark 2. We write $\operatorname{Cat}^m(W)$ for $\operatorname{Cat}^m(W;1)$.

By definition of the cyclic sieving phenomenon, we have to prove that

$$|\operatorname{Fix}_{NC^m(W)}(\phi^p)| = \operatorname{Cat}^m(W;q)\big|_{q=e^{2\pi i p/mh}}, \tag{3.3}$$

for all p in the range $0 \le p < mh$.

4. Auxiliary results I

This section collects several auxiliary lemmas which allow us to reduce the problem of proving Theorem 2, respectively the equivalent statement (3.3), for the 26 exceptional groups listed in Section 2 to a finite problem. While Lemmas 4 and 5 cover special choices of the parameters, Lemmas 3 and 7 afford an inductive procedure. More precisely, if we assume that we have already verified Theorem 2 for all groups of smaller rank, then Lemmas 3 and 7, together with Lemmas 4 and 8, reduce the verification of Theorem 2 for the group that we are currently considering to a finite problem; see Remark 3. The final lemma of this section, Lemma 9, disposes of complex reflection groups with a special property satisfied by their degrees.

Let $p = am + b$, $0 \le b < m$. We have

$$\begin{multline}\phi^p\big((w_0; w_1, \dots, w_m)\big)\\ = (*; c^{a+1} w_{m-b+1} c^{-a-1}, c^{a+1} w_{m-b+2} c^{-a-1}, \dots, c^{a+1} w_m c^{-a-1},\\ c^a w_1 c^{-a}, \dots, c^a w_{m-b} c^{-a}\big), \quad (4.1)\end{multline}$$

where $*$ stands for the element of W which is needed to complete the product of the components to c.

Lemma 3. *It suffices to check* (3.3) *for p a divisor of mh. More precisely, let p be a divisor of mh, and let k be another positive integer with* $\gcd(k, mh/p) = 1$, *then we have*

$$\operatorname{Cat}^m(W; q)\big|_{q=e^{2\pi i p/mh}} = \operatorname{Cat}^m(W; q)\big|_{q=e^{2\pi i kp/mh}} \tag{4.2}$$

and

$$|\operatorname{Fix}_{NC^m(W)}(\phi^p)| = |\operatorname{Fix}_{NC^m(W)}(\phi^{kp})|. \tag{4.3}$$

Proof. For (4.2), this follows immediately from

$$\lim_{q \to \zeta} \frac{[\alpha]_q}{[\beta]_q} = \begin{cases} \frac{\alpha}{\beta} & \text{if } \alpha \equiv \beta \equiv 0 \pmod d, \\ 1 & \text{otherwise,} \end{cases} \tag{4.4}$$

where ζ is a d-th root of unity and α, β are non-negative integers such that $\alpha \equiv \beta \pmod d$.

In order to establish (4.3), suppose that $x \in \operatorname{Fix}_{NC^m(W)}(\phi^p)$, that is, $x \in NC^m(W)$ and $\phi^p(x) = x$. It obviously follows that $\phi^{kp}(x) = x$, so that $x \in \operatorname{Fix}_{NC^m(W)}(\phi^{kp})$. Conversely, if $\gcd(k, mh/p) = 1$, then there exists k' with $k'k \equiv 1 \pmod{\frac{mh}{p}}$. It follows that, if $x \in \operatorname{Fix}_{NC^m(W)}(\phi^{kp})$, that is, if $x \in NC^m(W)$ and $\phi^{kp}(x) = x$, then $x = \phi^{k'kp}(x) = \phi^p(x)$, whence $x \in \operatorname{Fix}_{NC^m(W)}(\phi^p)$. □

Lemma 4. *Let p be a divisor of mh. If p is divisible by m, then* (3.3) *is true.*

Proof. According to (4.1), the action of ϕ^p on $NC^m(W)$ is described by

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) = (*; c^{p/m} w_1 c^{-p/m}, \dots, c^{p/m} w_m c^{-p/m}\big).$$

Hence, if $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, then each individual w_i must be fixed under conjugation by $c^{p/m}$.

Using the notation $W' = \operatorname{Cent}_W(c^{p/m})$, the previous observation means that $w_i \in W'$, $i = 1, 2, \dots, m$. Springer [26, Theorem 4.2] (see also [21, Theorem 11.24(iii)]) proved that W' is a well-generated complex reflection group whose degrees coincide with those degrees of W that are divisible by mh/p. It was furthermore shown in [7, Lemma 3.3] that

$$NC(W) \cap W' = NC(W'). \tag{4.5}$$

Hence, the tuples $(w_0; w_1, \dots, w_m)$ fixed by ϕ^p are in fact identical with the elements of $NC^m(W')$, which implies that

$$|\operatorname{Fix}_{NC^m(W)}(\phi^p)| = |NC^m(W')|. \tag{4.6}$$

Application of Theorem 1 with W replaced by W' and of the "limit rule" (4.4) then yields that

$$|NC^m(W')| = \prod_{\substack{1 \le i \le n \\ \frac{mh}{p} | d_i}} \frac{mh + d_i}{d_i} = \operatorname{Cat}^m(W; q)\big|_{q = e^{2\pi i p/mh}}. \tag{4.7}$$

Combining (4.6) and (4.7), we obtain (3.3). This finishes the proof of the lemma. □

Lemma 5. *Equation* (3.3) *holds for all divisors p of m.*

Proof. Using (4.4), we see that

$$\operatorname{Cat}^m(W; q)\big|_{q = e^{2\pi i p/mh}} = \begin{cases} m + 1 & \text{if } m = p, \\ 1 & \text{if } m \neq p. \end{cases}$$

On the other hand, if $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, then, because of the action (4.1), we must have $w_1 = w_{p+1} = \cdots = w_{m-p+1}$ and $w_1 = c w_{m-p+1} c^{-1}$. In particular, $w_1 \in \operatorname{Cent}_W(c)$. By the theorem of Springer cited in the proof of Lemma 4, the subgroup $\operatorname{Cent}_W(c)$ is itself a complex reflection group whose degrees are those degrees of W that are divisible by h. The only such degree is h itself, hence $\operatorname{Cent}_W(c)$ is the cyclic group generated by c. Moreover, by (4.5), we obtain that $w_1 = \varepsilon$ or $w_1 = c$. Therefore, for $m = p$ the set $\operatorname{Fix}_{NC^m(W)}(\phi^p)$ consists of the $m + 1$ elements $(w_0; w_1, \dots, w_m)$ obtained by choosing $w_i = c$ for a particular i between 0 and m, all other w_j's being equal to ε, while, for $m \neq p$, we have

$$\operatorname{Fix}_{NC^m(W)}(\phi) = \big\{(c; \varepsilon, \dots, \varepsilon)\big\},$$

whence the result. □

Lemma 6. *Let W be an irreducible well-generated complex reflection group all of whose degrees are divisible by d. Then each element of W is fixed under conjugation by $c^{h/d}$.*

Proof. By the theorem of Springer cited in the proof of Lemma 4, the subgroup $W' = \operatorname{Cent}_W(c^{h/d})$ is itself a complex reflection group whose degrees are those degrees of W that are divisible by d. Thus, by our assumption, the degrees of W' coincide with the degrees of W, and hence W' must be equal to W. Phrased differently, each element of W is fixed under conjugation by $c^{h/d}$, as claimed. □

Lemma 7. *Let W be an irreducible well-generated complex reflection group of rank n, and let $p = m_1 h_1$ be a divisor of mh, where $m = m_1 m_2$ and $h = h_1 h_2$. We assume that* $\gcd(h_1, m_2) = 1$. *Suppose that Theorem 2 has already been verified for all irreducible well-generated complex reflection groups with rank $< n$. If h_2 does not divide all degrees d_i, then Equation* (3.3) *is satisfied.*

Proof. Let us write $h_1 = am_2 + b$, with $0 \le b < m_2$. The condition $\gcd(h_1, m_2) = 1$ translates into $\gcd(b, m_2) = 1$. From (4.1), we infer that

$$\begin{multline}\phi^p\big((w_0; w_1, \dots, w_m)\big)\\ = (*; c^{a+1}w_{m-m_1b+1}c^{-a-1}, c^{a+1}w_{m-m_1b+2}c^{-a-1}, \dots, c^{a+1}w_mc^{-a-1},\\ c^aw_1c^{-a}, \dots, c^aw_{m-m_1b}c^{-a}\big). \quad (4.8)\end{multline}$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$\begin{aligned} w_i &= c^{a+1}w_{i+m-m_1b}c^{-a-1}, \quad i = 1, 2, \dots, m_1b,\\ w_i &= c^aw_{i-m_1b}c^{-a}, \quad i = m_1b + 1, m_1b + 2, \dots, m, \end{aligned}$$

which, after iteration, implies in particular that

$$w_i = c^{b(a+1)+(m_2-b)a}w_ic^{-b(a+1)-(m_2-b)a} = c^{h_1}w_ic^{-h_1}, \quad i = 1, 2, \dots, m.$$

It is at this point where we need $\gcd(b, m_2) = 1$. The last equation shows that each w_i, $i = 1, 2, \dots, m$, and thus also w_0, lies in $\mathrm{Cent}_W(c^{h_1})$. By the theorem of Springer cited in the proof of Lemma 4, this centraliser subgroup is itself a complex reflection group, W' say, whose degrees are those degrees of W that are divisible by $h/h_1 = h_2$. Since, by assumption, h_2 does not divide *all* degrees, W' has rank strictly less than n. Again by assumption, we know that Theorem 2 is true for W', so that in particular,

$$|\operatorname{Fix}_{NC^m(W')}(\phi^p)| = \mathrm{Cat}^m(W'; q)\big|_{q=e^{2\pi ip/mh}}.$$

The arguments above together with (4.5) show that $\operatorname{Fix}_{NC^m(W)}(\phi^p) = \operatorname{Fix}_{NC^m(W')}(\phi^p)$. On the other hand, using (4.4) it is straightforward to see that

$$\mathrm{Cat}^m(W; q)\big|_{q=e^{2\pi ip/mh}} = \mathrm{Cat}^m(W'; q)\big|_{q=e^{2\pi ip/mh}}.$$

This proves (3.3) for our particular p, as required. □

Lemma 8. *Let W be an irreducible well-generated complex reflection group of rank n, and let $p = m_1h_1$ be a divisor of mh, where $m = m_1m_2$ and $h = h_1h_2$. We assume that* $\gcd(h_1, m_2) = 1$. *If $m_2 > n$ then*

$$\operatorname{Fix}_{NC^m(W)}(\phi^p) = \big\{(c; \varepsilon, \dots, \varepsilon)\big\}.$$

Proof. Let us suppose that $(w_0; w_1, \dots, w_m) \in \operatorname{Fix}_{NC^m(W)}(\phi^p)$ and that there exists a $j \ge 1$ such that $w_j \ne \varepsilon$. By (4.8), it then follows for such a j that also $w_k \ne \varepsilon$ for all $k \equiv j - lm_1b \pmod m$, where, as before, b is defined as the unique integer with $h_1 = am_2 + b$ and $0 \le b < m_2$. Since, by assumption, $\gcd(b, m_2) = 1$, there are exactly m_2 such k's which are distinct mod m. However, this implies that the sum of the absolute lengths of the w_i's, $0 \le i \le m$, is at least $m_2 > n$, a contradiction. □

Remark 3. (1) If we put ourselves in the situation of the assumptions of Lemma 7, then we may conclude that Equation (3.3) only needs to be checked for pairs (m_2, h_2) subject to the following restrictions:

$$m_2 \ge 2, \quad \gcd(h_1, m_2) = 1, \quad \text{and } h_2 \text{ divides all degrees of } W. \tag{4.9}$$

Indeed, Lemmas 4 and 7 together imply that Equation (3.3) is always satisfied except if $m_2 \ge 2$, h_2 divides all degrees of W, and $\gcd(h_1, m_2) = 1$.

(2) Still putting ourselves in the situation of Lemma 7, if $m_2 > n$ and m_2h_2 does not divide any of the degrees of W, then Equation (3.3) is satisfied. Indeed, Lemma 8 says

that in this case the left-hand side of (3.3) equals 1, while a straightforward computation using (4.4) shows that in this case the right-hand side of (3.3) equals 1 as well.

(3) It should be observed that this leaves a finite number of choices for m_2 to consider, whence a finite number of choices for (m_1, m_2, h_1, h_2). Altogether, there remains a finite number of choices for $p = h_1 m_1$ to be checked.

Lemma 9. *Let W be an irreducible well-generated complex reflection group of rank n with the property that $d_i \mid h$ for $i = 1, 2, \dots, n$. Then Theorem 2 is true for this group W.*

Proof. By Lemma 3, we may restrict ourselves to divisors p of mh.

Suppose that $e^{2\pi i p/mh}$ is a d_i-th root of unity for some i. In other words, mh/p divides d_i. Since d_i is a divisor of h by assumption, the integer mh/p also divides h. But this is equivalent to saying that m divides p, and Equation (3.3) holds by Lemma 4.

Now assume that mh/p does not divide any of the d_i's. Then, by (4.4), the right-hand side of (3.3) equals 1. On the other hand, $(\varepsilon; \varepsilon, \dots, \varepsilon)$ is always an element of $\operatorname{Fix}_{NC^m(W)}(\phi^p)$. To see that there are no others, we make appeal to the classification of all irreducible well-generated complex reflection groups, which we recalled in Section 2. Inspection reveals that all groups satisfying the hypotheses of the lemma have rank $n \le 2$. Except for the groups contained in the infinite series $G(d, 1, n)$ and $G(e, e, n)$ for which Theorem 2 has been established in [16], these are the groups $G_5, G_6, G_9, G_{10}, G_{14}, G_{17}, G_{18}, G_{21}$. We now discuss these groups case by case, keeping the notation of Lemma 7. In order to simplify the argument, we note that Lemma 8 implies that Equation (3.3) holds if $m_2 > 2$, so that in the following arguments we always may assume that $m_2 = 2$.

CASE G_5. The degrees are $6, 12$, and therefore Remark 3.(1) implies that Equation (3.3) is always satisfied.

CASE G_6. The degrees are $4, 12$, and therefore, according to Remark 3.(1), we need only consider the case where $h_2 = 4$ and $m_2 = 2$, that is, $p = 3m/2$. Then (4.8) becomes

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) = (*; c^2 w_{\frac{m}{2}+1} c^{-2}, c^2 w_{\frac{m}{2}+2} c^{-2}, \dots, c^2 w_m c^{-2}, c w_1 c^{-1}, \dots, c w_{\frac{m}{2}} c^{-1}). \tag{4.10}$$

If $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p and not equal to $(c; \varepsilon, \dots, \varepsilon)$, there must exist an i with $1 \le i \le \frac{m}{2}$ such that $\ell_T(w_i) = 1$, $w_i c w_i c^{-1} = c$, and all w_j, $j \ne i, \frac{m}{2} + i$, equal ε. However, with the help of the *GAP* package `CHEVIE` [12], one verifies that there is no such solution to this equation. Hence, the left-hand side of (3.3) is equal to 1, as required.

CASE G_9. The degrees are $8, 24$, and therefore, according to Remark 3.(1), we need only consider the case where $h_2 = 8$ and $m_2 = 2$, that is, $p = 3m/2$. This is the same p as for G_6. Again, the *GAP* package `CHEVIE` [12] finds no solution. Hence, the left-hand side of (3.3) is equal to 1, as required.

CASE G_{10}. The degrees are $12, 24$, and therefore, according to Remark 3.(1), we need only consider the case where $h_2 = 12$ and $m_2 = 2$, that is, $p = 3m/2$. This is the same p as for G_6. Again, the *GAP* package `CHEVIE` [12] finds no solution. Hence, the left-hand side of (3.3) is equal to 1, as required.

CASE G_{14}. The degrees are $6, 24$, and therefore Remark 3.(1) implies that Equation (3.3) is always satisfied.

Case G_{17}. The degrees are $20, 60$, and therefore, according to Remark 3.(1), we need only consider the cases where $h_2 = 20$ and $m_2 = 2$, respectively that $h_2 = 4$ and $m_2 = 2$. In the first case, $p = 3m/2$, which is the same p as for G_6. Again, the *GAP* package `CHEVIE` [12] finds no solution. In the second case, $p = 15m/2$. Then (4.8) becomes

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^8 w_{\frac{m}{2}+1} c^{-8}, c^8 w_{\frac{m}{2}+2} c^{-8}, \dots, c^8 w_m c^{-8}, c^7 w_1 c^{-7}, \dots, c^7 w_{\frac{m}{2}} c^{-7}\big). \quad (4.11)$$

By Lemma 6, every element of $NC(W)$ is fixed under conjugation by c^3, and, thus, on elements fixed by ϕ^p, the above action of ϕ^p reduces to the one in (4.10). This action was already discussed in the first case. Hence, in both cases, the left-hand side of (3.3) is equal to 1, as required.

Case G_{18}. The degrees are $30, 60$, and therefore Remark 3.(1) implies that Equation (3.3) is always satisfied.

Case G_{21}. The degrees are $12, 60$, and therefore, according to Remark 3.(1), we need only consider the cases where $h_2 = 5$ and $m_2 = 2$, respectively that $h_2 = 15$ and $m_2 = 2$. In the first case, $p = 5m/2$, so that (4.8) becomes

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^3 w_{\frac{m}{2}+1} c^{-3}, c^3 w_{\frac{m}{2}+2} c^{-3}, \dots, c^3 w_m c^{-3}, c^2 w_1 c^{-2}, \dots, c^2 w_{\frac{m}{2}} c^{-2}\big). \quad (4.12)$$

If $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p and not equal to $(c; \varepsilon, \dots, \varepsilon)$, there must exist an i with $1 \le i \le \frac{m}{2}$ such that $\ell_T(w_i) = 1$ and $w_i c^2 w_i c^{-2} = c$. However, with the help of the *GAP* package `CHEVIE` [12], one verifies that there is no such solution to this equation. In the second case, $p = 15m/2$. Then (4.8) becomes the action in (4.11). By Lemma 6, every element of $NC(W)$ is fixed under conjugation by c^5, and, thus, on elements fixed by ϕ^p, the action of ϕ^p in (4.11) reduces to the one in the first case. Hence, in both cases, the left-hand side of (3.3) is equal to 1, as required.

This completes the proof of the lemma. □

5. Case-by-case verification of Theorem 2

In the sequel we write ζ_d for a primitive d-th root of unity.

Case G_4. The degrees are $4, 6$, and hence we have

$$\operatorname{Cat}^m(G_4; q) = \frac{[6m+6]_q\,[6m+4]_q}{[6]_q\,[4]_q}.$$

Let ζ be a $6m$-th root of unity. In what follows, we abbreviate the assertion that "ζ is a primitive d-th root of unity" as "$\zeta = \zeta_d$." The following cases on the right-hand side of (3.3) do occur:

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_4; q) = m+1, \quad \text{if } \zeta = \zeta_6, \zeta_3, \tag{5.1a}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_4; q) = \tfrac{3m+2}{2}, \quad \text{if } \zeta = \zeta_4,\ 2 \mid m, \tag{5.1b}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_4; q) = \operatorname{Cat}^m(G_4), \quad \text{if } \zeta = -1 \text{ or } \zeta = 1, \tag{5.1c}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_4; q) = 1, \quad \text{otherwise.} \tag{5.1d}$$

We must now prove that the left-hand side of (3.3) in each case agrees with the values exhibited in (5.1). The only cases not covered by Lemmas 4 and 5 are the ones in (5.1b) and (5.1d). On the other hand, the only case left to consider according to Remark 3 is the case where $h_2 = m_2 = 2$, that is the case (5.1b) where $p = 3m/2$. In particular, m must be divisible by 2. The action of ϕ^p is the same as the one in (4.10). With the help of the *GAP* package `CHEVIE` [12], one finds that each of the 3 (complex) reflections in G_4 which are less than the (chosen) Coxeter element is a valid choice for w_i, and each of these choices gives rise to $m/2$ elements in $NC^m(G_4)$ since the index i ranges from 1 to $m/2$.

Hence, in total, we obtain $1 + 3\frac{m}{2} = \frac{3m+2}{2}$ elements in $\mathrm{Fix}_{NC^m(G_4)}(\phi^p)$, which agrees with the limit in (5.1b).

Case G_8. The degrees are $8, 12$, and hence we have

$$\mathrm{Cat}^m(G_8; q) = \frac{[12m+12]_q\,[12m+8]_q}{[12]_q\,[8]_q}.$$

Let ζ be a $12m$-th root of unity. The following cases on the right-hand side of (3.3) do occur:

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_8; q) = m+1, \quad \text{if } \zeta = \zeta_{12}, \zeta_6, \zeta_3, \tag{5.2a}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_8; q) = \tfrac{3m+2}{2}, \quad \text{if } \zeta = \zeta_8,\ 2 \mid m, \tag{5.2b}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_8; q) = \mathrm{Cat}^m(G_8), \quad \text{if } \zeta = \zeta_4, -1, 1, \tag{5.2c}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_8; q) = 1, \quad \text{otherwise.} \tag{5.2d}$$

We must now prove that the left-hand side of (3.3) in each case agrees with the values exhibited in (5.2). The only cases not covered by Lemmas 4 and 5 are the ones in (5.2b) and (5.2d). On the other hand, the only case left to consider according to Remark 3 is the case where $h_2 = 4$ and $m_2 = 2$, that is the case (5.2b) where $p = 3m/2$. In particular, m must be divisible by 2. The action of ϕ^p is the same as the one in (4.10). With the help of the *GAP* package `CHEVIE` [12], one finds that each of the 3 (complex) reflections in G_8 which are less than the (chosen) Coxeter element is a valid choice for w_i, and each of these choices gives rise to $m/2$ elements in $NC^m(G_8)$ since the index i ranges from 1 to $m/2$.

Hence, in total, we obtain $1 + 3\frac{m}{2} = \frac{3m+2}{2}$ elements in $\mathrm{Fix}_{NC^m(G_8)}(\phi^p)$, which agrees with the limit in (5.2b).

Case G_{16}. The degrees are $20, 30$, and hence we have

$$\mathrm{Cat}^m(G_{16}; q) = \frac{[30m+30]_q\,[30m+20]_q}{[30]_q\,[20]_q}.$$

Let ζ be a $30m$-th root of unity. The following cases on the right-hand side of (3.3) do occur:

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{16};q) = m+1, \quad \text{if } \zeta=\zeta_{30},\zeta_{15},\zeta_6,\zeta_3, \tag{5.3a}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{16};q) = \tfrac{3m+2}{2}, \quad \text{if } \zeta=\zeta_{20},\zeta_4,\ 2\mid m, \tag{5.3b}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{16};q) = \operatorname{Cat}^m(G_{16}), \quad \text{if } \zeta=\zeta_{10},\zeta_5,-1,1, \tag{5.3c}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{16};q) = 1, \quad \text{otherwise.} \tag{5.3d}$$

We must now prove that the left-hand side of (3.3) in each case agrees with the values exhibited in (5.3). The only cases not covered by Lemmas 4 and 5 are the ones in (5.3b) and (5.3d). On the other hand, the only cases left to consider according to Remark 3 are the cases where $h_2 = 10$ and $m_2 = 2$, respectively $h_2 = m_2 = 2$. Both cases belong to (5.3b). In the first case, we have $p = 3m/2$, while in the second case we have $p = 15m/2$. In particular, m must be divisible by 2. In the first case, the action of ϕ^p is the same as the one in (4.10). With the help of the *GAP* package `CHEVIE` [12], one finds that each of the 3 (complex) reflections in G_{16} which are less than the (chosen) Coxeter element is a valid choice for w_i, and each of these choices gives rise to $m/2$ elements in $NC^m(G_{16})$ since the index i ranges from 1 to $m/2$. On the other hand, if $p = 15m/2$, then the action of ϕ^p is the same as the one in (4.11). By Lemma 6, every element of $NC(W)$ is fixed under conjugation by c^3, and, thus, on elements fixed by ϕ^p, the action of ϕ^p reduces to the one in the first case.

Hence, in total, we obtain $1+3\frac{m}{2} = \frac{3m+2}{2}$ elements in $\operatorname{Fix}_{NC^m(G_{16})}(\phi^p)$, which agrees with the limit in (5.3b).

Case G_{20}. The degrees are $12, 30$, and hence we have

$$\operatorname{Cat}^m(G_{20};q) = \frac{[30m+30]_q\,[30m+12]_q}{[30]_q\,[12]_q}.$$

Let ζ be a $30m$-th root of unity. The following cases on the right-hand side of (3.3) do occur:

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{20};q) = m+1, \quad \text{if } \zeta=\zeta_{30},\zeta_{15},\zeta_{10},\zeta_5, \tag{5.4a}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{20};q) = \tfrac{5m+2}{2}, \quad \text{if } \zeta=\zeta_{12},\zeta_4,\ 2\mid m, \tag{5.4b}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{20};q) = \operatorname{Cat}^m(G_{20}), \quad \text{if } \zeta=\zeta_6,\zeta_3,-1,1, \tag{5.4c}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{20};q) = 1, \quad \text{otherwise.} \tag{5.4d}$$

We must now prove that the left-hand side of (3.3) in each case agrees with the values exhibited in (5.4). The only cases not covered by Lemmas 4 and 5 are the ones in (5.4b) and (5.4d). On the other hand, the only cases left to consider according to Remark 3 are the cases where $h_2 = 6$ and $m_2 = 2$, respectively $h_2 = m_2 = 2$. Both cases belong to (5.4b). In the first case, we have $p = 5m/2$, while in the second case we have $p = 15m/2$. In particular, m must be divisible by 2. In the first case, the action of ϕ^p is the same as the one in (4.12). With the help of the *GAP* package `CHEVIE` [12], one finds that each of the 5 (complex) reflections in G_{20} which are less than the (chosen) Coxeter element is a valid choice for w_i, and each of these choices gives rise to $m/2$

elements in $NC^m(G_{20})$ since the index i ranges from 1 to $m/2$. On the other hand, if $p = 15m/2$, then the action of ϕ^p is the same as the one in (4.11). By Lemma 6, every element of $NC(W)$ is fixed under conjugation by c^5, and, thus, on elements fixed by ϕ^p, the action of ϕ^p reduces to the one in the first case.

Hence, in total, we obtain $1 + 5\frac{m}{2} = \frac{5m+2}{2}$ elements in $\operatorname{Fix}_{NC^m(G_{20})}(\phi^p)$, which agrees with the limit in (5.4b).

Case $G_{23} = H_3$. The degrees are $2, 6, 10$, and hence we have

$$\operatorname{Cat}^m(H_3;q) = \frac{[10m+10]_q\,[10m+6]_q\,[10m+2]_q}{[10]_q\,[6]_q\,[2]_q}.$$

Let ζ be a $10m$-th root of unity. The following cases on the right-hand side of (3.3) do occur:

$$\lim_{q\to\zeta}\operatorname{Cat}^m(H_3;q) = m+1, \quad \text{if } \zeta = \zeta_{10}, \zeta_5, \tag{5.5a}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(H_3;q) = \tfrac{10m+6}{6}, \quad \text{if } \zeta = \zeta_6, \zeta_3,\ 3 \mid m, \tag{5.5b}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(H_3;q) = \operatorname{Cat}^m(H_3), \quad \text{if } \zeta = -1 \text{ or } \zeta = 1, \tag{5.5c}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(H_3;q) = 1, \quad \text{otherwise.} \tag{5.5d}$$

We must now prove that the left-hand side of (3.3) in each case agrees with the values exhibited in (5.5). The only cases not covered by Lemmas 4 and 5 are the ones in (5.5b) and (5.5d). By Lemma 3, we are free to choose $p = 5m/3$ if $\zeta = \zeta_6$, respectively $p = 10m/3$ if $\zeta = \zeta_3$. In both cases, m must be divisible by 3.

We start with the case that $p = 5m/3$. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) = (*; c^2 w_{\frac{m}{3}+1} c^{-2}, c^2 w_{\frac{m}{3}+2} c^{-2}, \dots, c^2 w_m c^{-2}, c w_1 c^{-1}, \dots, c w_{\frac{m}{3}} c^{-1}\big). \tag{5.6}$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^2 w_{\frac{m}{3}+i} c^{-2}, \quad i = 1, 2, \dots, \tfrac{2m}{3}, \tag{5.7a}$$

$$w_i = c w_{i-\frac{2m}{3}} c^{-1}, \quad i = \tfrac{2m}{3}+1, \tfrac{2m}{3}+2, \dots, m. \tag{5.7b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$: either all the w_i's are equal to ε, or there is an i with $1 \le i \le \frac{m}{3}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{3}}) = \ell_T(w_{i+\frac{2m}{3}}) = 1.$$

Writing t_1, t_2, t_3 for $w_i, w_{i+\frac{m}{3}}, w_{i+\frac{2m}{3}}$, in that order, the equations (5.7) reduce to

$$t_1 = c^2 t_2 c^{-2}, \tag{5.8a}$$

$$t_2 = c^2 t_3 c^{-2}, \tag{5.8b}$$

$$t_3 = c t_1 c^{-1}. \tag{5.8c}$$

One of these equations is in fact superfluous: if we substitute (5.8b) and (5.8c) in (5.8a), then we obtain $t_1 = c^5 t_1 c^{-5}$ which is automatically satisfied due to Lemma 6 with $d = 2$.

Since $(w_0; w_1, \dots, w_m) \in NC^m(H_3)$, we must have $t_1 t_2 t_3 = c$. Combining this with (5.8), we infer that

$$t_1 (c^{-2} t_1 c^2)(c t_1 c^{-1}) = c. \tag{5.9}$$

With the help of Stembridge's *Maple* package `coxeter` [29], one obtains five solutions for t_1 in this equation:

$$t_1 \in \big\{[2],\ [3],\ [2,1,2],\ [1,2,3,2,1],\ [1,3,2,1,2,1,3]\big\}. \tag{5.10}$$

Here we have used the short notation of `coxeter`: if $\{s_1, s_2, s_3\}$ is a simple system of generators of H_3, corresponding to the Dynkin diagram displayed in Figure 1, then $[j_1, j_2, \dots, j_k]$ stands for the element $s_{j_1} s_{j_2} \dots s_{j_k}$.



FIGURE 1. The Dynkin diagram for H_3

We claim that each of the above five solutions gives rise to $m/3$ elements of $\operatorname{Fix}_{NC^m(H_3)}(\phi^p)$. Indeed, given t_1, the elements t_2 and t_3 can be computed by (5.8a) and (5.8c), and there are $m/3$ possibilities to choose the index i for w_i.

In total, we obtain $1 + 5\frac{m}{3} = \frac{10m+6}{6}$ elements in $\operatorname{Fix}_{NC^m(H_3)}(\phi^p)$, which agrees with the limit in (5.5b).

In the case that $p = 10m/3$, we infer from (4.1) that

$$\begin{multline}\phi^p\big((w_0; w_1, \dots, w_m)\big)\\ = \big(*; c^4 w_{\frac{2m}{3}+1} c^{-4}, c^4 w_{\frac{2m}{3}+2} c^{-4}, \dots, c^4 w_m c^{-4}, c^3 w_1 c^{-3}, \dots, c^3 w_{\frac{2m}{3}} c^{-3}\big). \tag{5.11}\end{multline}$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^4 w_{\frac{2m}{3}+i} c^{-4}, \quad i = 1, 2, \dots, \tfrac{m}{3}, \tag{5.12a}$$

$$w_i = c^3 w_{i-\frac{m}{3}} c^{-3}, \quad i = \tfrac{m}{3}+1, \tfrac{m}{3}+2, \dots, m. \tag{5.12b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$: either all the w_i's are equal to ε, or there is an i with $1 \le i \le \frac{m}{3}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{3}}) = \ell_T(w_{i+\frac{2m}{3}}) = 1.$$

Writing t_1, t_2, t_3 for $w_i, w_{i+\frac{m}{3}}, w_{i+\frac{2m}{3}}$, in that order, the equations (5.12) reduce to

$$t_1 = c^4 t_3 c^{-4}, \tag{5.13a}$$

$$t_2 = c^3 t_1 c^{-3}, \tag{5.13b}$$

$$t_3 = c^3 t_2 c^{-3}. \tag{5.13c}$$

One of these equations is in fact superfluous: if we substitute (5.13b) and (5.13c) in (5.13a), then we obtain $t_1 = c^{10} t_1 c^{-10}$ which is automatically satisfied since $c^{10} = \varepsilon$.

Since $(w_0; w_1, \dots, w_m) \in NC^m(H_3)$, we must have $t_1 t_2 t_3 = c$. Combining this with (5.13), we infer that

$$t_1 (c^3 t_1 c^{-3})(c^{-4} t_1 c^4) = c. \tag{5.14}$$

Using that $c^5 t_1 c^{-5} = t_1$, due to Lemma 6 with $d = 2$, we see that this equation is equivalent with (5.9). Therefore, we are facing exactly the same enumeration problem here as for $p = 5m/3$, and, consequently, the number of solutions to (5.14) is the same, namely $\frac{5m+3}{3}$, as required.

Finally, we turn to (5.5d). By Remark 3, the only choices for h_2 and m_2 to be considered are $h_2 = 1$ and $m_2 = 3$, $h_2 = m_2 = 2$, respectively $h_2 = 2$ and $m_2 = 3$.

These correspond to the choices $p = 10m/3$, $p = 5m/2$, respectively $p = 5m/3$, out of which only $p = 5m/2$ has not yet been discussed and belongs to the current case. The corresponding action of ϕ^p is given by (4.12). A computation with Stembridge's *Maple* package `coxeter` [29] finds no solution. Hence, the left-hand side of (3.3) is equal to 1, as required.

Case G_{24}. The degrees are $4, 6, 14$, and hence we have

$$\mathrm{Cat}^m(G_{24}; q) = \frac{[14m + 14]_q\,[14m + 6]_q\,[14m + 4]_q}{[14]_q\,[6]_q\,[4]_q}.$$

Let ζ be a $14m$-th root of unity. The following cases on the right-hand side of (3.3) do occur:

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_{24}; q) = m + 1, \quad \text{if } \zeta = \zeta_{14}, \zeta_7, \tag{5.15a}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_{24}; q) = \tfrac{7m+3}{3}, \quad \text{if } \zeta = \zeta_6, \zeta_3,\ 3 \mid m, \tag{5.15b}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_{24}; q) = \tfrac{7m+2}{2}, \quad \text{if } \zeta = \zeta_4,\ 2 \mid m, \tag{5.15c}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_{24}; q) = \mathrm{Cat}^m(G_{24}), \quad \text{if } \zeta = -1 \text{ or } \zeta = 1, \tag{5.15d}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_{24}; q) = 1, \quad \text{otherwise.} \tag{5.15e}$$

We must now prove that the left-hand side of (3.3) in each case agrees with the values exhibited in (5.15). The only cases not covered by Lemmas 4 and 5 are the ones in (5.15b), (5.15c), and (5.15e).

We first consider (5.15b). By Lemma 3, we are free to choose $p = 7m/3$ if $\zeta = \zeta_6$, respectively $p = 14m/3$ if $\zeta = \zeta_3$. In both cases, m must be divisible by 3.

We start with the case that $p = 7m/3$. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^3 w_{\frac{2m}{3}+1} c^{-3}, c^3 w_{\frac{2m}{3}+2} c^{-3}, \dots, c^3 w_m c^{-3}, c^2 w_1 c^{-2}, \dots, c^2 w_{\frac{2m}{3}} c^{-2}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^3 w_{\frac{2m}{3}+i} c^{-3}, \quad i = 1, 2, \dots, \tfrac{m}{3}, \tag{5.16a}$$

$$w_i = c^2 w_{i-\frac{m}{3}} c^{-2}, \quad i = \tfrac{m}{3} + 1, \tfrac{m}{3} + 2, \dots, m. \tag{5.16b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$: either all the w_i's are equal to ε, or there is an i with $1 \le i \le \frac{m}{3}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{3}}) = \ell_T(w_{i+\frac{2m}{3}}) = 1.$$

Writing t_1, t_2, t_3 for $w_i, w_{i+\frac{m}{3}}, w_{i+\frac{2m}{3}}$, in that order, the equations (5.16) reduce to

$$t_1 = c^3 t_3 c^{-3}, \tag{5.17a}$$

$$t_2 = c^2 t_1 c^{-2}, \tag{5.17b}$$

$$t_3 = c^2 t_2 c^{-2}. \tag{5.17c}$$

One of these equations is in fact superfluous: if we substitute (5.17b) and (5.17c) in (5.17a), then we obtain $t_1 = c^7 t_1 c^{-7}$ which is automatically satisfied due to Lemma 6 with $d = 2$.

Since $(w_0; w_1, \dots, w_m) \in NC^m(G_{24})$, we must have $t_1t_2t_3 = c$. Combining this with (5.17), we infer that

$$t_1(c^2t_1c^{-2})(c^4t_1c^{-4}) = c. \tag{5.18}$$

With the help of the *GAP* package `CHEVIE` [12], one obtains 7 solutions for t_1 in this equation:

$$t_1 \in \{[1], [2], [3], [15], [16], [19], [21]\}, \tag{5.19}$$

each of them giving rise to $m/3$ elements of $\operatorname{Fix}_{NC^m(G_{24})}(\phi^p)$ since i ranges from 1 to $m/3$. Here we have used the short notation of `CHEVIE`: $[j_1, j_2, \dots, j_k]$ stands for the element $r_{j_1}r_{j_2}\dots r_{j_k}$, where r_i is the i-th (complex) reflection corresponding to the i-th root in the internal ordering of the roots of G_{24} in `CHEVIE`.

In total, we obtain $1 + 7\frac{m}{3} = \frac{7m+3}{3}$ elements in $\operatorname{Fix}_{NC^m(G_{24})}(\phi^p)$, which agrees with the limit in (5.15b).

In the case that $p = 14m/3$, we infer from (4.1) that

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = (*; c^5w_{\frac{m}{3}+1}c^{-5}, c^5w_{\frac{m}{3}+2}c^{-5}, \dots, c^5w_mc^{-5}, c^4w_1c^{-4}, \dots, c^4w_{\frac{m}{3}}c^{-4}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^5w_{\frac{m}{3}+i}c^{-5}, \quad i = 1, 2, \dots, \tfrac{2m}{3}, \tag{5.20a}$$

$$w_i = c^4w_{i-\frac{2m}{3}}c^{-4}, \quad i = \tfrac{2m}{3}+1, \tfrac{2m}{3}+2, \dots, m. \tag{5.20b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$: either all the w_i's are equal to ε, or there is an i with $1 \le i \le \frac{m}{3}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{3}}) = \ell_T(w_{i+\frac{2m}{3}}) = 1.$$

Writing t_1, t_2, t_3 for $w_i, w_{i+\frac{m}{3}}, w_{i+\frac{2m}{3}}$, in that order, the equations (5.20) reduce to

$$t_1 = c^5t_2c^{-5}, \tag{5.21a}$$

$$t_2 = c^5t_3c^{-5}, \tag{5.21b}$$

$$t_3 = c^4t_1c^{-4}. \tag{5.21c}$$

One of these equations is in fact superfluous: if we substitute (5.21b) and (5.21c) in (5.21a), then we obtain $t_1 = c^{14}t_1c^{-14}$ which is automatically satisfied since $c^{14} = \varepsilon$.

Since $(w_0; w_1, \dots, w_m) \in NC^m(G_{24})$, we must have $t_1t_2t_3 = c$. Combining this with (5.21), we infer that

$$t_1(c^9t_1c^{-9})(c^{-4}t_1c^4) = c. \tag{5.22}$$

Using that $c^7t_1c^{-7} = t_1$, due to Lemma 6 with $d = 2$, we see that this equation is equivalent with (5.18). Therefore, we are facing exactly the same enumeration problem here as for $p = 7m/3$, and, consequently, the number of solutions to (5.22) is the same, namely $\frac{7m+3}{3}$, as required.

Our next case is (5.15c). By Lemma 3, we are free to choose $p = 7m/2$. In particular, m must be divisible by 2. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = (*; c^4w_{\frac{m}{2}+1}c^{-4}, c^4w_{\frac{m}{2}+2}c^{-4}, \dots, c^4w_mc^{-4}, c^3w_1c^{-3}, \dots, c^3w_{\frac{m}{2}}c^{-3}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^4 w_{\frac{m}{2}+i} c^{-4}, \quad i = 1, 2, \dots, \tfrac{m}{2}, \tag{5.23a}$$

$$w_i = c^3 w_{i-\frac{m}{2}} c^{-3}, \quad i = \tfrac{m}{2}+1, \tfrac{m}{2}+2, \dots, m. \tag{5.23b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$: either all the w_i's are equal to ε, or there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 1.$$

Writing t_1, t_2 for $w_i, w_{i+\frac{m}{2}}$, in that order, the equations (5.23) reduce to

$$t_1 = c^4 t_2 c^{-4}, \tag{5.24a}$$

$$t_2 = c^3 t_1 c^{-3}. \tag{5.24b}$$

One of these equations is in fact superfluous: if we substitute (5.24b) in (5.24a), then we obtain $t_1 = c^7 t_1 c^{-7}$ which is automatically satisfied due to Lemma 6 with $d = 2$.

Since $(w_0; w_1, \dots, w_m) \in NC^m(G_{24})$, we must have $t_1 t_2 \le_T c$, where ℓ_T is the partial order defined in (2.1). Combining this with (5.24), we infer that

$$t_1(c^3 t_1 c^{-3}) \le_T c. \tag{5.25}$$

With the help of the *GAP* package `CHEVIE` [12], one obtains 7 solutions for t_1 in this equation:

$$t_1 \in \big\{[5],\ [6],\ [7],\ [9],\ [12],\ [29],\ [32]\big\}, \tag{5.26}$$

each of them giving rise to $m/2$ elements of $\mathrm{Fix}_{NC^m(G_{24})}(\phi^p)$ since i ranges from 1 to $m/2$. Here we have used again the short notation of `CHEVIE` referring to the internal ordering of the roots of G_{24} in `CHEVIE`.

In total, we obtain $1 + 7\frac{m}{2} = \frac{7m+2}{2}$ elements in $\mathrm{Fix}_{NC^m(G_{24})}(\phi^p)$, which agrees with the limit in (5.15c).

Finally, we turn to (5.15e). By Remark 3, the only choices for h_2 and m_2 to be considered are $h_2 = 1$ and $m_2 = 3$, $h_2 = m_2 = 2$, respectively $h_2 = 2$ and $m_2 = 3$. These correspond to the choices $p = 14m/3$, $p = 7m/2$, respectively $p = 7m/3$, all of which have already been discussed as they do not belong to (5.15e). Hence, (3.3) must necessarily hold, as required.

Case G_{25}. The degrees are $6, 9, 12$, and hence we have

$$\mathrm{Cat}^m(G_{25}; q) = \frac{[12m+12]_q\,[12m+9]_q\,[12m+6]_q}{[12]_q\,[9]_q\,[6]_q}.$$

Let ζ be a $12m$-th root of unity. The following cases on the right-hand side of (3.3) do occur:

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_{25}; q) = m+1, \quad \text{if } \zeta = \zeta_{12}, \zeta_4, \tag{5.27a}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_{25}; q) = \tfrac{4m+3}{3}, \quad \text{if } \zeta = \zeta_9,\ 3 \mid m, \tag{5.27b}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_{25}; q) = (m+1)(2m+1), \quad \text{if } \zeta = \zeta_6, -1 \tag{5.27c}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_{25}; q) = \mathrm{Cat}^m(G_{25}), \quad \text{if } \zeta = \zeta_3, 1, \tag{5.27d}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_{25}; q) = 1, \quad \text{otherwise.} \tag{5.27e}$$

We must now prove that the left-hand side of (3.3) in each case agrees with the values exhibited in (5.27). The only cases not covered by Lemmas 4 and 5 are the ones in (5.27b) and (5.27e).

We first consider (5.27b). By Lemma 3, we are free to choose $p = 4m/3$. In particular, m must be divisible by 3. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^2 w_{\frac{2m}{3}+1} c^{-2}, c^2 w_{\frac{2m}{3}+2} c^{-2}, \dots, c^2 w_m c^{-2}, c w_1 c^{-1}, \dots, c w_{\frac{2m}{3}} c^{-1}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^2 w_{\frac{2m}{3}+i} c^{-2}, \quad i = 1, 2, \dots, \tfrac{m}{3}, \tag{5.28a}$$

$$w_i = c w_{i-\frac{m}{3}} c^{-1}, \quad i = \tfrac{m}{3}+1, \tfrac{m}{3}+2, \dots, m. \tag{5.28b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$: either all the w_i's are equal to ε, or there is an i with $1 \le i \le \frac{m}{3}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{3}}) = \ell_T(w_{i+\frac{2m}{3}}) = 1.$$

Writing t_1, t_2, t_3 for $w_i, w_{i+\frac{m}{3}}, w_{i+\frac{2m}{3}}$, in that order, the equations (5.28) reduce to

$$t_1 = c^2 t_3 c^{-2}, \tag{5.29a}$$

$$t_2 = c t_1 c^{-1}, \tag{5.29b}$$

$$t_3 = c t_2 c^{-1}. \tag{5.29c}$$

One of these equations is in fact superfluous: if we substitute (5.29b) and (5.29c) in (5.29a), then we obtain $t_1 = c^4 t_1 c^{-4}$ which is automatically satisfied due to Lemma 6 with $d = 3$.

Since $(w_0; w_1, \dots, w_m) \in NC^m(G_{25})$, we must have $t_1 t_2 t_3 = c$. Combining this with (5.29), we infer that

$$t_1 (c t_1 c^{-1})(c^2 t_1 c^{-2}) = c. \tag{5.30}$$

With the help of the *GAP* package `CHEVIE` [12], one obtains four solutions for t_1 in this equation:

$$t_1 \in \big\{[1], [2], [3], [14]\big\}, \tag{5.31}$$

each of them giving rise to $m/3$ elements of $\mathrm{Fix}_{NC^m(G_{25})}(\phi^p)$ since i ranges from 1 to $m/3$. Here we have used again the short notation of `CHEVIE` referring to the internal ordering of the roots of G_{25} in `CHEVIE`.

In total, we obtain $1 + 4\frac{m}{3} = \frac{4m+3}{3}$ elements in $\mathrm{Fix}_{NC^m(G_{25})}(\phi^p)$, which agrees with the limit in (5.27b).

Finally, we turn to (5.27e). By Remark 3, the only choice for h_2 and m_2 to be considered are $h_2 = m_2 = 3$. This corresponds to the choice $p = 4m/3$, which has already been discussed as they do not belong to (5.27e). Hence, (3.3) must necessarily hold, as required.

Case G_{26}. The degrees are $6, 12, 18$, and hence we have

$$\mathrm{Cat}^m(G_{26}; q) = \frac{[18m+18]_q\,[18m+12]_q\,[18m+6]_q}{[18]_q\,[12]_q\,[6]_q}.$$

Let ζ be a $14m$-th root of unity. The following cases on the right-hand side of (3.3) do occur:

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{26};q)=m+1,\quad \text{if } \zeta=\zeta_{18},\zeta_9, \tag{5.32a}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{26};q)=\tfrac{3m+2}{2},\quad \text{if } \zeta=\zeta_{12},\zeta_4,\ 2\mid m, \tag{5.32b}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{26};q)=\operatorname{Cat}^m(G_{26}),\quad \text{if } \zeta=\zeta_6,\zeta_3,-1,1, \tag{5.32c}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{26};q)=1,\quad \text{otherwise.} \tag{5.32d}$$

We must now prove that the left-hand side of (3.3) in each case agrees with the values exhibited in (5.32). The only cases not covered by Lemmas 4 and 5 are the ones in (5.32b) and (5.32d).

We first consider (5.32b). By Lemma 3, we are free to choose $p=3m/2$ if $\zeta=\zeta_{12}$, respectively $p=9m/2$ if $\zeta=\zeta_4$. In both cases, m must be divisible by 2.

We start with the case that $p=3m/2$. From (4.1), we infer

$$\phi^p\big((w_0;w_1,\dots,w_m)\big)=(*;c^2w_{\frac{m}{2}+1}c^{-2},c^2w_{\frac{m}{2}+2}c^{-2},\dots,c^2w_mc^{-2},cw_1c^{-1},\dots,cw_{\frac{m}{2}}c^{-1}\big).$$

Supposing that $(w_0;w_1,\dots,w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i=c^2w_{\frac{m}{2}+i}c^{-2},\quad i=1,2,\dots,\tfrac{m}{2}, \tag{5.33a}$$

$$w_i=cw_{i-\frac{m}{2}}c^{-1},\quad i=\tfrac{m}{2}+1,\tfrac{m}{2}+2,\dots,m. \tag{5.33b}$$

There are two distinct possibilities for choosing the w_i's, $1\le i\le m$: either all the w_i's are equal to ε, or there is an i with $1\le i\le\frac{m}{2}$ such that

$$\ell_T(w_i)=\ell_T(w_{i+\frac{m}{2}})=1.$$

Writing t_1,t_2 for $w_i,w_{i+\frac{m}{2}}$, in that order, the equations (5.33) reduce to

$$t_1=c^2t_2c^{-2}, \tag{5.34a}$$

$$t_2=c^1t_1c^{-1}. \tag{5.34b}$$

One of these equations is in fact superfluous: if we substitute (5.34b) in (5.34a), then we obtain $t_1=c^3t_1c^{-3}$ which is automatically satisfied due to Lemma 6 with $d=6$.

Since $(w_0;w_1,\dots,w_m)\in NC^m(G_{26})$, we must have $t_1t_2\le_T c$. Combining this with (5.34), we infer that

$$t_1(ct_1c^{-1})\le_T c. \tag{5.35}$$

With the help of the *GAP* package `CHEVIE` [12], one obtains three solutions for t_1 in this equation:

$$t_1\in\big\{[2],\ [3],\ [12]\big\},$$

each of them giving rise to $m/2$ elements of $\operatorname{Fix}_{NC^m(G_{26})}(\phi^p)$ since i ranges from 1 to $m/2$. Here we have again used the short notation of `CHEVIE` referring to the internal ordering of the roots of G_{26} in `CHEVIE`.

In total, we obtain $1+3\frac{m}{2}=\frac{3m+2}{2}$ elements in $\operatorname{Fix}_{NC^m(G_{26})}(\phi^p)$, which agrees with the limit in (5.32b).

In the case that $p=9m/2$, we infer from (4.1) that

$$\phi^p\big((w_0;w_1,\dots,w_m)\big)$$
$$=(*;c^5w_{\frac{m}{2}+1}c^{-5},c^5w_{\frac{m}{2}+2}c^{-5},\dots,c^5w_mc^{-5},c^4w_1c^{-4},\dots,c^4w_{\frac{m}{2}}c^{-4}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^5 w_{\frac{m}{2}+i} c^{-5}, \quad i = 1, 2, \dots, \tfrac{m}{2}, \tag{5.36a}$$

$$w_i = c^4 w_{i-\frac{m}{2}} c^{-4}, \quad i = \tfrac{m}{2}+1, \tfrac{m}{2}+2, \dots, m. \tag{5.36b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$: either all the w_i's are equal to ε, or there is an i with $1 \le i \le \frac{m}{3}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 1.$$

Writing t_1, t_2 for $w_i, w_{i+\frac{m}{2}}$, in that order, the equations (5.36) reduce to

$$t_1 = c^5 t_2 c^{-5}, \tag{5.37a}$$

$$t_2 = c^4 t_1 c^{-4}. \tag{5.37b}$$

One of these equations is in fact superfluous: if we substitute (5.37b) in (5.37a), then we obtain $t_1 = c^9 t_1 c^{-9}$ which is automatically satisfied due to Lemma 6 with $d = 2$. Since $(w_0; w_1, \dots, w_m) \in NC^m(G_{26})$, we must have $t_1 t_2 \le_T c$. Combining this with (5.37), we infer that

$$t_1(c^4 t_1 c^{-4}) \le_T c. \tag{5.38}$$

Using that $c^3 t_1 c^{-3} = t_1$, due to Lemma 6 with $d = 6$, we see that this equation is equivalent with (5.35). Therefore, we are facing exactly the same enumeration problem here as for $p = 3m/2$, and, consequently, the number of solutions to (5.38) is the same, namely $\frac{3m+2}{2}$, as required.

Finally, we turn to (5.32d). By Remark 3, the only choices for h_2 and m_2 to be considered are $h_2 = 6$ and $m_2 = 2$, respectively $h_2 = m_2 = 2$. These correspond to the choices $p = 3m/2$, respectively $p = 9m/2$, all of which have already been discussed as they do not belong to (5.32d). Hence, (3.3) must necessarily hold, as required.

Case G_{27}. The degrees are $6, 12, 30$, and hence we have

$$\operatorname{Cat}^m(G_{27}; q) = \frac{[30m+30]_q\,[30m+12]_q\,[30m+6]_q}{[30]_q\,[12]_q\,[6]_q}.$$

Let ζ be a $14m$-th root of unity. The following cases on the right-hand side of (3.3) do occur:

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{27}; q) = m+1, \quad \text{if } \zeta = \zeta_{30}, \zeta_{15}, \zeta_{10}, \zeta_5, \tag{5.39a}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{27}; q) = \tfrac{5m+2}{2}, \quad \text{if } \zeta = \zeta_{12}, \zeta_4,\ 2 \mid m, \tag{5.39b}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{27}; q) = \operatorname{Cat}^m(G_{27}), \quad \text{if } \zeta = \zeta_6, \zeta_3, -1, 1, \tag{5.39c}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{27}; q) = 1, \quad \text{otherwise.} \tag{5.39d}$$

We must now prove that the left-hand side of (3.3) in each case agrees with the values exhibited in (5.39). The only cases not covered by Lemmas 4 and 5 are the ones in (5.39b) and (5.39d).

We first consider (5.39b). By Lemma 3, we are free to choose $p = 5m/2$ if $\zeta = \zeta_{12}$, respectively $p = 15m/2$ if $\zeta = \zeta_4$. In both cases, m must be divisible by 2.

We start with the case that $p = 5m/2$. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^3 w_{\frac{m}{2}+1} c^{-3}, c^3 w_{\frac{m}{2}+2} c^{-3}, \dots, c^3 w_m c^{-3}, c^2 w_1 c^{-2}, \dots, c^2 w_{\frac{m}{2}} c^{-2}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^3 w_{\frac{m}{2}+i} c^{-3}, \quad i = 1, 2, \dots, \tfrac{m}{2}, \tag{5.40a}$$

$$w_i = c^2 w_{i-\frac{m}{2}} c^{-2}, \quad i = \tfrac{m}{2}+1, \tfrac{m}{2}+2, \dots, m. \tag{5.40b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$: either all the w_i's are equal to ε, or there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 1.$$

Writing t_1, t_2 for $w_i, w_{i+\frac{m}{2}}$, in that order, the equations (5.40) reduce to

$$t_1 = c^3 t_2 c^{-3}, \tag{5.41a}$$

$$t_2 = c^2 t_1 c^{-2}. \tag{5.41b}$$

One of these equations is in fact superfluous: if we substitute (5.41b) in (5.41a), then we obtain $t_1 = c^5 t_1 c^{-5}$ which is automatically satisfied due to Lemma 6 with $d = 6$.

Since $(w_0; w_1, \dots, w_m) \in NC^m(G_{27})$, we must have $t_1 t_2 \le_T c$. Combining this with (5.41), we infer that

$$t_1(c^2 t_1 c^{-2}) \le_T c. \tag{5.42}$$

With the help of the *GAP* package `CHEVIE` [12], one obtains five solutions for t_1 in this equation:

$$t_1 \in \big\{[1], [2], [15], [16], [28]\big\},$$

each of them giving rise to $m/2$ elements of $\mathrm{Fix}_{NC^m(G_{27})}(\phi^p)$ since i ranges from 1 to $m/2$. Here we have used the short notation of `CHEVIE` referring to the internal ordering of the roots of G_{27} in `CHEVIE`.

In total, we obtain $1 + 5\frac{m}{2} = \frac{5m+2}{2}$ elements in $\mathrm{Fix}_{NC^m(G_{27})}(\phi^p)$, which agrees with the limit in (5.39b).

In the case that $p = 15m/2$, we infer from (4.1) that

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^8 w_{\frac{m}{2}+1} c^{-8}, c^8 w_{\frac{m}{2}+2} c^{-8}, \dots, c^8 w_m c^{-8}, c^7 w_1 c^{-7}, \dots, c^7 w_{\frac{m}{2}} c^{-7}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^8 w_{\frac{m}{2}+i} c^{-8}, \quad i = 1, 2, \dots, \tfrac{m}{2}, \tag{5.43a}$$

$$w_i = c^7 w_{i-\frac{m}{2}} c^{-7}, \quad i = \tfrac{m}{2}+1, \tfrac{m}{2}+2, \dots, m. \tag{5.43b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$: either all the w_i's are equal to ε, or there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 1.$$

Writing t_1, t_2 for $w_i, w_{i+\frac{m}{2}}$, in that order, the equations (5.43) reduce to

$$t_1 = c^8 t_2 c^{-8}, \tag{5.44a}$$

$$t_2 = c^7 t_1 c^{-7}. \tag{5.44b}$$

One of these equations is in fact superfluous: if we substitute (5.44b) in (5.44a), then we obtain $t_1 = c^{15}t_1c^{-15}$ which is automatically satisfied due to Lemma 6 with $d = 2$.

Since $(w_0; w_1, \dots, w_m) \in NC^m(G_{27})$, we must have $t_1t_2 \le_T c$. Combining this with (5.44), we infer that

$$t_1(c^7t_1c^{-7}) \le_T c. \tag{5.45}$$

Using that $c^5t_1c^{-5} = t_1$, due to Lemma 6 with $d = 6$, we see that this equation is equivalent with (5.42). Therefore, we are facing exactly the same enumeration problem here as for $p = 5m/2$, and, consequently, the number of solutions to (5.45) is the same, namely $\frac{5m+2}{2}$, as required.

Finally, we turn to (5.39d). By Remark 3, the only choices for h_2 and m_2 to be considered are $h_2 = 6$ and $m_2 = 3$, $h_2 = 6$ and $m_2 = 2$, $h_2 = m_2 = 3$, respectively $h_2 = m_2 = 2$. These correspond to the choices $p = 5m/3$, $5m/2$, $10m/3$, respectively $15m/2$, out of which only $p = 5m/3$ and $p = 10m/3$ have not yet been discussed and belong to the current case. If $p = 5m/3$, the corresponding action of ϕ^p is given by (5.6), so that we have to solve for t_1 with $\ell_T(t_1) = 1$ in the equation (5.9). A computation with the help of the *GAP* package `CHEVIE` [12] finds no solution. If $p = 10m/3$, the corresponding action of ϕ^p is given by (5.11), so that we have to solve for t_1 with $\ell_T(t_1)$ in the equation (5.14). Using that $c^5t_1c^{-5} = t_1$, due to Lemma 6 with $d = 6$, we see that this equation is equivalent with the one in (5.9). Hence, in both cases, the left-hand side of (3.3) is equal to 1, as required.

CASE $G_{28} = F_4$. The degrees are $2, 6, 8, 12$, and hence we have

$$\operatorname{Cat}^m(F_4; q) = \frac{[12m+12]_q\,[12m+8]_q\,[12m+6]_q\,[12m+2]_q}{[12]_q\,[8]_q\,[6]_q\,[2]_q}.$$

Let ζ be a $12m$-th root of unity. The following cases on the right-hand side of (3.3) do occur:

$$\lim_{q\to\zeta} \operatorname{Cat}^m(F_4; q) = m+1, \quad \text{if } \zeta = \zeta_{12}, \tag{5.46a}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(F_4; q) = \tfrac{3m+2}{2}, \quad \text{if } \zeta = \zeta_8,\ 2 \mid m, \tag{5.46b}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(F_4; q) = (m+1)(2m+1), \quad \text{if } \zeta = \zeta_6, \zeta_3, \tag{5.46c}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(F_4; q) = \tfrac{(m+1)(3m+2)}{2}, \quad \text{if } \zeta = \zeta_4, \tag{5.46d}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(F_4; q) = \operatorname{Cat}^m(F_4), \quad \text{if } \zeta = -1 \text{ or } \zeta = 1, \tag{5.46e}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(F_4; q) = 1, \quad \text{otherwise}. \tag{5.46f}$$

We must now prove that the left-hand side of (3.3) in each case agrees with the values exhibited in (5.46). The only cases not covered by Lemmas 4 and 5 are the ones in (5.46b) and (5.46f). By Lemma 3, we are free to choose $p = 3m/2$. In particular, m must be divisible by 2. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) = (*; c^2w_{\frac{m}{2}+1}c^{-2}, c^2w_{\frac{m}{2}+2}c^{-2}, \dots, c^2w_mc^{-2}, cw_1c^{-1}, \dots, cw_{\frac{m}{2}}c^{-1}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^2 w_{\frac{m}{2}+i} c^{-2}, \quad i = 1, 2, \dots, \tfrac{m}{2}, \tag{5.47a}$$

$$w_i = c w_{i-\frac{m}{2}} c^{-1}, \quad i = \tfrac{m}{2}+1, \tfrac{m}{2}+2, \dots, m. \tag{5.47b}$$

There are four distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 2, \tag{5.48a}$$

and all other w_j's are equal to ε,

(iii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 1, \tag{5.48b}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(iv) there are i_1 and i_2 with $1 \le i_1 < i_2 \le \frac{m}{2}$ such that

$$\ell_T(w_{i_1}) = \ell_T(w_{i_2}) = \ell_T(w_{i_1+\frac{m}{2}}) = \ell_T(w_{i_2+\frac{m}{2}}) = 1, \tag{5.48c}$$

and all other w_j's are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(F_4)$, we must have $w_i w_{i+\frac{m}{2}} \le_T c$, respectively $w_{i_1} w_{i_2} w_{i_1+\frac{m}{2}} w_{i_2+\frac{m}{2}} = c$. Together with Equations (5.47)–(5.48), this implies that

$$w_i = c^3 w_i c^{-3} \quad \text{and} \quad w_i(c w_i c^{-1}) = c, \tag{5.49}$$

respectively that

$$w_i = c^3 w_i c^{-3}, \quad w_i(c w_i c^{-1}) \le_T c, \quad \text{and} \quad \ell_T(w_i) = 1, \tag{5.50}$$

respectively that

$$w_{i_1} = c^3 w_{i_1} c^{-3}, \quad w_{i_1}(c w_{i_1} c^{-1}) \le_T c, \quad \text{and} \quad \ell_T(w_{i_1}) = 1. \tag{5.51}$$

With the help of Stembridge's *Maple* package `coxeter` [29], one obtains three solutions for w_i in (5.49):

$$w_i \in \big\{[1,2,3,4,3,2],\ [2,3],\ [1,3,2,1,3,4]\big\},$$

where we have again used the short notation of `coxeter`, $\{s_1, s_2, s_3, s_4\}$ being a simple system of generators of F_4, corresponding to the Dynkin diagram displayed in Figure 2. Each of the above solutions for w_i gives rise to $m/2$ elements of $\mathrm{Fix}_{NC^m(F_4)}(\phi^p)$ since i ranges from 1 to $m/2$.



FIGURE 2. The Dynkin diagram for F_4

There are no solutions for w_i in (5.50) and for w_{i_1} in (5.51).

In total, we obtain $1 + 3\frac{m}{2} = \frac{3m+2}{2}$ elements in $\mathrm{Fix}_{NC^m(F_4)}(\phi^p)$, which agrees with the limit in (5.46b).

Finally, we turn to (5.46f). By Remark 3, the are no choices for h_2 and m_2 to be considered. Hence, the left-hand side of (3.3) is equal to 1, as required.

CASE G_{29}. The degrees are $4, 8, 12, 20$, and hence we have

$$\operatorname{Cat}^m(G_{29};q)=\frac{[20m+20]_q\,[20m+12]_q\,[20m+8]_q\,[20m+4]_q}{[20]_q\,[12]_q\,[8]_q\,[4]_q}.$$

Let ζ be a $20m$-th root of unity. The following cases on the right-hand side of (3.3) do occur:

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{29};q)=m+1,\quad \text{if } \zeta=\zeta_{20},\zeta_{10},\zeta_5, \tag{5.52a}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{29};q)=\tfrac{5m+3}{3},\quad \text{if } \zeta=\zeta_{12},\zeta_6,\zeta_3,\ 3\mid m, \tag{5.52b}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{29};q)=\tfrac{5m+2}{2},\quad \text{if } \zeta=\zeta_8,\ 2\mid m, \tag{5.52c}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{29};q)=\operatorname{Cat}^m(G_{29}),\quad \text{if } \zeta=\zeta_4,-1,1, \tag{5.52d}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{29};q)=1,\quad \text{otherwise.} \tag{5.52e}$$

We must now prove that the left-hand side of (3.3) in each case agrees with the values exhibited in (5.52). The only cases not covered by Lemmas 4 and 5 are the ones in (5.52b), (5.52c), and (5.52e).

We begin with the case in (5.52b). By Lemma 3, we are free to choose $p=5m/3$ if $\zeta=\zeta_{12}$, we are free to choose $p=10m/3$ if $\zeta=\zeta_6$, we are free to choose $p=20m/3$ if $\zeta=\zeta_3$. In particular, in all three cases, m must be divisible by 3.

We start with the case that $p=5m/3$. From (4.1), we infer

$$\phi^p\big((w_0;w_1,\dots,w_m)\big)=(*;c^2w_{\frac{m}{3}+1}c^{-2},c^2w_{\frac{m}{3}+2}c^{-2},\dots,c^2w_mc^{-2},cw_1c^{-1},\dots,cw_{\frac{m}{3}}c^{-1}\big).$$

Supposing that $(w_0;w_1,\dots,w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i=c^2w_{\frac{m}{3}+i}c^{-2},\quad i=1,2,\dots,\tfrac{2m}{3}, \tag{5.53a}$$

$$w_i=cw_{i-\frac{2m}{3}}c^{-1},\quad i=\tfrac{2m}{3}+1,\tfrac{2m}{3}+2,\dots,m. \tag{5.53b}$$

There are two distinct possibilities for choosing the w_i's, $1\le i\le m$:

(i) all the w_i's are equal to ε (and $w_0=c$),

(ii) there is an i with $1\le i\le\frac{m}{3}$ such that

$$\ell_T(w_i)=\ell_T(w_{i+\frac{m}{3}})=\ell_T(w_{i+\frac{2m}{3}})=1, \tag{5.54a}$$

and the other w_j's, $1\le j\le m$, are equal to ε.

Moreover, since $(w_0;w_1,\dots,w_m)\in NC^m(G_{29})$, we must have $w_iw_{i+\frac{m}{3}}w_{i+\frac{2m}{3}}\le_T c$. Together with Equations (5.53)–(5.54), this implies that

$$w_i=c^5w_ic^{-5}\quad\text{and}\quad w_i(c^3w_ic^{-3})(cw_ic^{-1})=c. \tag{5.55}$$

With the help of the *GAP* package `CHEVIE` [12], one obtains five solutions for w_i in (5.55):

$$w_i\in\big\{[1],\,[2],\,[8],\,[25],\,[31]\big\},$$

where we have again used the short notation of `CHEVIE` referring to the internal ordering of the roots of G_{29} in `CHEVIE`. Each of the above solutions for w_i gives rise to $m/3$ elements of $\operatorname{Fix}_{NC^m(G_{29})}(\phi^p)$ since i ranges from 1 to $m/3$.

In total, we obtain $1+5\frac{m}{3}=\frac{5m+3}{3}$ elements in $\operatorname{Fix}_{NC^m(G_{29})}(\phi^p)$, which agrees with the limit in (5.52b).

In the case that $p = 10m/3$, we infer from (4.1) that

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^4 w_{\frac{2m}{3}+1} c^{-4}, c^4 w_{\frac{2m}{3}+2} c^{-4}, \dots, c^4 w_m c^{-4}, c^3 w_1 c^{-3}, \dots, c^3 w_{\frac{2m}{3}} c^{-3}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^4 w_{\frac{2m}{3}+i} c^{-4}, \quad i = 1, 2, \dots, \tfrac{m}{3}, \tag{5.56a}$$

$$w_i = c^3 w_{i-\frac{m}{3}} c^{-3}, \quad i = \tfrac{m}{3}+1, \tfrac{m}{3}+2, \dots, m. \tag{5.56b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$: either all the w_i's are equal to ε, or there is an i with $1 \le i \le \frac{m}{3}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{3}}) = \ell_T(w_{i+\frac{2m}{3}}) = 1.$$

Writing t_1, t_2, t_3 for $w_i, w_{i+\frac{m}{3}}, w_{i+\frac{2m}{3}}$, in that order, the equations (5.56) reduce to

$$t_1 = c^4 t_3 c^{-4}, \tag{5.57a}$$

$$t_2 = c^3 t_1 c^{-3}, \tag{5.57b}$$

$$t_3 = c^3 t_2 c^{-3}. \tag{5.57c}$$

One of these equations is in fact superfluous: if we substitute (5.57b) and (5.57c) in (5.57a), then we obtain $t_1 = c^{10} t_1 c^{-10}$ which is automatically satisfied due to Lemma 6 with $d = 2$.

Since $(w_0; w_1, \dots, w_m) \in NC^m(G_{29})$, we must have $t_1 t_2 t_3 \le_T c$. Combining this with (5.57), we infer that

$$t_1 (c^3 t_1 c^{-3})(c^6 t_1 c^{-6}) \le_T c. \tag{5.58}$$

Using that $c^5 t_1 c^{-5} = t_1$, due to Lemma 6 with $d = 4$, we see that this equation is equivalent with (5.55). Therefore, we are facing exactly the same enumeration problem here as for $p = 5m/3$, and, consequently, the number of solutions to (5.58) is the same, namely $\frac{5m+3}{3}$, as required.

In the case that $p = 20m/3$, we infer from (4.1) that

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^7 w_{\frac{m}{3}+1} c^{-7}, c^7 w_{\frac{m}{3}+2} c^{-7}, \dots, c^7 w_m c^{-7}, c^6 w_1 c^{-6}, \dots, c^6 w_{\frac{m}{3}} c^{-6}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^7 w_{\frac{m}{3}+i} c^{-7}, \quad i = 1, 2, \dots, \tfrac{2m}{3}, \tag{5.59a}$$

$$w_i = c^6 w_{i-\frac{2m}{3}} c^{-6}, \quad i = \tfrac{2m}{3}+1, \tfrac{2m}{3}+2, \dots, m. \tag{5.59b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$: either all the w_i's are equal to ε, or there is an i with $1 \le i \le \frac{m}{3}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{3}}) = \ell_T(w_{i+\frac{2m}{3}}) = 1.$$

Writing t_1, t_2, t_3 for $w_i, w_{i+\frac{m}{3}}, w_{i+\frac{2m}{3}}$, in that order, the equations (5.59) reduce to

$$t_1 = c^7 t_2 c^{-7}, \tag{5.60a}$$

$$t_2 = c^7 t_3 c^{-7}, \tag{5.60b}$$

$$t_3 = c^6 t_1 c^{-6}. \tag{5.60c}$$

One of these equations is in fact superfluous: if we substitute (5.60b) and (5.60c) in (5.60a), then we obtain $t_1 = c^{20} t_1 c^{-20}$ which is automatically satisfied since $c^{20} = \varepsilon$.

Since $(w_0; w_1, \dots, w_m) \in NC^m(G_{29})$, we must have $t_1 t_2 t_3 \le_T c$. Combining this with (5.60), we infer that

$$t_1 (c^{13} t_1 c^{-13})(c^6 t_1 c^{-6}) \le_T c. \tag{5.61}$$

Using that $c^5 t_1 c^{-5} = t_1$, due to Lemma 6 with $d = 4$, we see that this equation is equivalent with (5.55). Therefore, we are facing exactly the same enumeration problem here as for $p = 5m/3$, and, consequently, the number of solutions to (5.61) is the same, namely $\frac{5m+3}{3}$, as required.

Next we discuss the case in (5.52c). By Lemma 3, we are free to choose $p = 5m/2$. In particular, m must be divisible by 2. From (4.1), we infer

$$\begin{aligned}\phi^p\big((w_0; w_1, \dots, w_m)\big)\\ &= (*; c^3 w_{\frac{m}{2}+1} c^{-3}, c^3 w_{\frac{m}{2}+2} c^{-3}, \dots, c^3 w_m c^{-3}, c^2 w_1 c^{-2}, \dots, c^2 w_{\frac{m}{2}} c^{-2}).\end{aligned}$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^3 w_{\frac{m}{2}+i} c^{-3}, \quad i = 1, 2, \dots, \tfrac{m}{2}, \tag{5.62a}$$

$$w_i = c^2 w_{i-\frac{m}{2}} c^{-2}, \quad i = \tfrac{m}{2}+1, \tfrac{m}{2}+2, \dots, m. \tag{5.62b}$$

There are four distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 2, \tag{5.63a}$$

and all other w_j's are equal to ε,

(iii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 1, \tag{5.63b}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(iv) there are i_1 and i_2 with $1 \le i_1 < i_2 \le \frac{m}{2}$ such that

$$\ell_T(w_{i_1}) = \ell_T(w_{i_2}) = \ell_T(w_{i_1+\frac{m}{2}}) = \ell_T(w_{i_2+\frac{m}{2}}) = 1, \tag{5.63c}$$

and all other w_j's are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(G_{29})$, we must have $w_i w_{i+\frac{m}{2}} \le_T c$, respectively $w_{i_1} w_{i_2} w_{i_1+\frac{m}{2}} w_{i_2+\frac{m}{2}} = c$. Together with Equations (5.62)–(5.63), this implies that

$$w_i = c^3 w_i c^{-3} \quad \text{and} \quad w_i (c^2 w_i c^{-2}) = c, \tag{5.64}$$

respectively that

$$w_i = c^3 w_i c^{-3}, \quad w_i (c^2 w_i c^{-2}) \le_T c, \quad \text{and} \quad \ell_T(w_i) = 1, \tag{5.65}$$

respectively that

$$\begin{aligned} w_{i_1} = c^3 w_{i_1} c^{-3}, \quad w_{i_2} = c^3 w_{i_2} c^{-3},\\ w_{i_1} w_{i_2} (c^2 w_{i_1} c^{-2})(c^2 w_{i_2} c^{-2}) = c, \quad \text{and} \quad \ell_T(w_{i_1}) = \ell_T(w_{i_2}) = 1. \end{aligned} \tag{5.66}$$

With the help of the *GAP* package `CHEVIE` [12], one obtains five solutions for w_i in (5.65):

$$w_i \in \{[4], [9], [14], [27], [32]\},$$

where we have again used the short notation of `CHEVIE` referring to the internal ordering of the roots of G_{29} in `CHEVIE`. Each of these solutions for w_i gives rise to $m/2$ elements of $\operatorname{Fix}_{NC^m(G_{29})}(\phi^p)$ since i ranges from 1 to $m/2$.

There are no solutions for w_i in (5.64) and for (w_{i_1}, w_{i_2}) in (5.66).

In total, we obtain $1 + 5\frac{m}{2} = \frac{5m+2}{2}$ elements in $\operatorname{Fix}_{NC^m(G_{29})}(\phi^p)$, which agrees with the limit in (5.52c).

Finally, we turn to (5.52e). By Remark 3, the only choices for h_2 and m_2 to be considered are $h_2 = 1$ and $m_2 = 3$, $h_2 = 2$ and $m_2 = 3$, $h_2 = 4$ and $m_2 = 2$, $h_2 = 4$ and $m_2 = 3$, respectively $h_2 = m_2 = 4$. These correspond to the choices $p = 20m/3$, $p = 10m/3$, $p = 5m/2$, $p = 5m/3$, respectively $p = 5m/4$, out of which only $p = 5m/4$ has not yet been discussed and belongs to the current case. The corresponding action of ϕ^p is given by

$$\begin{aligned}\phi^p\big((w_0; w_1, \dots, w_m)\big)\\ &= \big(*; c^2 w_{\frac{3m}{4}+1} c^{-2}, c^2 w_{\frac{3m}{4}+2} c^{-2}, \dots, c^2 w_m c^{-2}, c w_1 c^{-1}, \dots, c w_{\frac{3m}{4}} c^{-1}\big),\end{aligned}$$

so that we have to solve

$$t_1 (c t_1 c^{-1})(c^2 t_1 c^{-2})(c^3 t_1 c^{-3}) = c$$

for t_1 with $\ell_T(t_1)$. A computation with the help of the *GAP* package `CHEVIE` [12] finds no solution. Hence, the left-hand side of (3.3) is equal to 1, as required.

Case $G_{30} = H_4$. The degrees are $2, 12, 20, 30$, and hence we have

$$\operatorname{Cat}^m(H_4; q) = \frac{[30m+30]_q\,[30m+20]_q\,[30m+12]_q\,[30m+2]_q}{[30]_q\,[20]_q\,[12]_q\,[2]_q}.$$

Let ζ be a $30m$-th root of unity. The following cases on the right-hand side of (3.3) do occur:

$$\lim_{q\to\zeta} \operatorname{Cat}^m(H_4; q) = m+1, \quad \text{if } \zeta = \zeta_{30}, \zeta_{15}, \tag{5.67a}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(H_4; q) = \tfrac{3m+2}{2}, \quad \text{if } \zeta = \zeta_{20},\ 2 \mid m, \tag{5.67b}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(H_4; q) = \tfrac{5m+2}{2}, \quad \text{if } \zeta = \zeta_{12},\ 2 \mid m, \tag{5.67c}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(H_4; q) = \tfrac{(m+1)(3m+2)}{2}, \quad \text{if } \zeta = \zeta_{10}, \zeta_5, \tag{5.67d}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(H_4; q) = \tfrac{(m+1)(5m+2)}{2}, \quad \text{if } \zeta = \zeta_6, \zeta_3, \tag{5.67e}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(H_4; q) = \tfrac{(3m+2)(5m+2)}{4}, \quad \text{if } \zeta = \zeta_4,\ 2 \mid m, \tag{5.67f}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(H_4; q) = \operatorname{Cat}^m(H_4), \quad \text{if } \zeta = -1 \text{ or } \zeta = 1, \tag{5.67g}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(H_4; q) = 1, \quad \text{otherwise.} \tag{5.67h}$$

We must now prove that the left-hand side of (3.3) in each case agrees with the values exhibited in (5.67). The only cases not covered by Lemmas 4 and 5 are the ones in (5.67b), (5.67c), (5.67f), and (5.67h).

We begin with the case in (5.67b). By Lemma 3, we are free to choose $p = 3m/2$. In particular, m must be divisible by 2. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) = (*; c^2 w_{\frac{m}{2}+1} c^{-2}, c^2 w_{\frac{m}{2}+2} c^{-2}, \dots, c^2 w_m c^{-2}, c w_1 c^{-1}, \dots, c w_{\frac{m}{2}} c^{-1}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^2 w_{\frac{m}{2}+i} c^{-2}, \quad i = 1, 2, \dots, \tfrac{m}{2}, \tag{5.68a}$$

$$w_i = c w_{i-\frac{m}{2}} c^{-1}, \quad i = \tfrac{m}{2}+1, \tfrac{m}{2}+2, \dots, m. \tag{5.68b}$$

There are four distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 2, \tag{5.69a}$$

and all other w_j's are equal to ε,

(iii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 1, \tag{5.69b}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(iv) there are i_1 and i_2 with $1 \le i_1 < i_2 \le \frac{m}{2}$ such that

$$\ell_T(w_{i_1}) = \ell_T(w_{i_2}) = \ell_T(w_{i_1+\frac{m}{2}}) = \ell_T(w_{i_2+\frac{m}{2}}) = 1, \tag{5.69c}$$

and all other w_j's are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(H_4)$, we must have $w_i w_{i+\frac{m}{2}} \le_T c$, respectively $w_{i_1} w_{i_2} w_{i_1+\frac{m}{2}} w_{i_2+\frac{m}{2}} = c$. Together with Equations (5.68)–(5.69), this implies that

$$w_i = c^3 w_i c^{-3} \quad \text{and} \quad w_i(c w_i c^{-1}) = c, \tag{5.70}$$

respectively that

$$w_i = c^3 w_i c^{-3}, \quad w_i(c w_i c^{-1}) \le_T c, \quad \text{and} \quad \ell_T(w_i) = 1, \tag{5.71}$$

respectively that

$$w_{i_1} = c^3 w_{i_1} c^{-3}, \quad w_{i_1}(c w_{i_1} c^{-1}) \le_T c, \quad \text{and} \quad \ell_T(w_{i_1}) = 1. \tag{5.72}$$

With the help of Stembridge's *Maple* package `coxeter` [29], one obtains three solutions for w_i in (5.70):

$$w_i \in \big\{[1,2,3,4,3,2,1,2], [2,1,2,3], [1,3,2,1,2,1,3,4]\big\},$$

where we have again used the short notation of `coxeter`, $\{s_1, s_2, s_3, s_4\}$ being a simple system of generators of H_4, corresponding to the Dynkin diagram displayed in Figure 3. Each of the above solutions for w_i gives rise to $m/2$ elements of $\mathrm{Fix}_{NC^m(H_4)}(\phi^p)$ since i ranges from 1 to $m/2$.



FIGURE 3. The Dynkin diagram for H_4

There are no solutions for w_i in (5.71) and for w_{i_1} in (5.72).

In total, we obtain $1 + 3\frac{m}{2} = \frac{3m+2}{2}$ elements in $\mathrm{Fix}_{NC^m(H_4)}(\phi^p)$, which agrees with the limit in (5.67b).

Next we discuss the case in (5.67c). By Lemma 3, we are free to choose $p = 5m/2$. In particular, m must be divisible by 2. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^3 w_{\frac{m}{2}+1} c^{-3}, c^3 w_{\frac{m}{2}+2} c^{-3}, \dots, c^3 w_m c^{-3}, c^2 w_1 c^{-2}, \dots, c^2 w_{\frac{m}{2}} c^{-2}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^3 w_{\frac{m}{2}+i} c^{-3}, \quad i = 1, 2, \dots, \tfrac{m}{2}, \tag{5.73a}$$

$$w_i = c^2 w_{i-\frac{m}{2}} c^{-2}, \quad i = \tfrac{m}{2}+1, \tfrac{m}{2}+2, \dots, m. \tag{5.73b}$$

There are four distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 2, \tag{5.74a}$$

and all other w_j's are equal to ε,

(iii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 1, \tag{5.74b}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(iv) there are i_1 and i_2 with $1 \le i_1 < i_2 \le \frac{m}{2}$ such that

$$\ell_T(w_{i_1}) = \ell_T(w_{i_2}) = \ell_T(w_{i_1+\frac{m}{2}}) = \ell_T(w_{i_2+\frac{m}{2}}) = 1, \tag{5.74c}$$

and all other w_j's are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(H_4)$, we must have $w_i w_{i+\frac{m}{2}} \le_T c$, respectively $w_{i_1} w_{i_2} w_{i_1+\frac{m}{2}} w_{i_2+\frac{m}{2}} = c$. Together with Equations (5.73)–(5.74), this implies that

$$w_i = c^3 w_i c^{-3} \quad \text{and} \quad w_i (c^2 w_i c^{-2}) = c, \tag{5.75}$$

respectively that

$$w_i = c^3 w_i c^{-3}, \quad w_i (c^2 w_i c^{-2}) \le_T c, \quad \text{and} \quad \ell_T(w_i) = 1, \tag{5.76}$$

respectively that

$$w_{i_1} = c^3 w_{i_1} c^{-3}, \quad w_{i_1} (c^2 w_{i_1} c^{-2}) \le_T c, \quad \text{and} \quad \ell_T(w_{i_1}) = 1. \tag{5.77}$$

With the help of Stembridge's *Maple* package `coxeter` [29], one obtains five solutions for w_i in (5.75):

$$w_i \in \big\{[1,3,2,1,2,1,3,2], [1,2,1,4,3,2,1,2,1,3,2,1,4,3], \\ [2,1,2,3,2,1,2,4], [2,1,2,1,4,3,2,1,2,1,3,4], [1,2,3,2,1,4,3,2,1,2,1,3]\big\},$$

where we used again `coxeter`'s short notation, $\{s_1, s_2, s_3, s_4\}$ being a simple system of generators of H_4, corresponding to the Dynkin diagram displayed in Figure 3. Each of these solutions for w_i gives rise to $m/2$ elements of $\mathrm{Fix}_{NC^m(H_4)}(\phi^p)$ since i ranges from 1 to $m/2$.

There are no solutions for w_i in (5.76) and for w_{i_1} in (5.77).

In total, we obtain $1 + 5\frac{m}{2} = \frac{5m+2}{2}$ elements in $\mathrm{Fix}_{NC^m(H_4)}(\phi^p)$, which agrees with the limit in (5.67c).

Finally we discuss the case in (5.67f). By Lemma 3, we are free to choose $p = 15m/2$. In particular, m must be divisible by 2. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^8 w_{\frac{m}{2}+1} c^{-8}, c^8 w_{\frac{m}{2}+2} c^{-8}, \dots, c^8 w_m c^{-8}, c^7 w_1 c^{-7}, \dots, c^7 w_{\frac{m}{2}} c^{-7}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^8 w_{\frac{m}{2}+i} c^{-8}, \quad i = 1, 2, \dots, \tfrac{m}{2}, \tag{5.78a}$$

$$w_i = c^7 w_{i-\frac{m}{2}} c^{-7}, \quad i = \tfrac{m}{2}+1, \tfrac{m}{2}+2, \dots, m. \tag{5.78b}$$

There are four distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 2, \tag{5.79a}$$

and all other w_j's are equal to ε,

(iii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 1, \tag{5.79b}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(iv) there are i_1 and i_2 with $1 \le i_1 < i_2 \le \frac{m}{2}$ such that

$$\ell_T(w_{i_1}) = \ell_T(w_{i_2}) = \ell_T(w_{i_1+\frac{m}{2}}) = \ell_T(w_{i_2+\frac{m}{2}}) = 1, \tag{5.79c}$$

and all other w_j's are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(H_4)$, we must have $w_i w_{i+\frac{m}{2}} \le_T c$, respectively $w_{i_1} w_{i_2} w_{i_1+\frac{m}{2}} w_{i_2+\frac{m}{2}} = c$. Together with Equations (5.78)–(5.79), this implies that

$$w_i = c^{15} w_i c^{-15} \quad \text{and} \quad w_i(c^7 w_i c^{-7}) = c, \tag{5.80}$$

respectively that

$$w_i = c^{15} w_i c^{-15}, \quad w_i(c^7 w_i c^{-7}) = c, \quad \text{and} \quad \ell_T(w_i) = 1, \tag{5.81}$$

respectively that

$$w_{i_1} = c^{15} w_{i_1} c^{-15}, \quad w_{i_2} = c^{15} w_{i_2} c^{-15}, \quad w_{i_1} w_{i_2}(c^7 w_{i_1} c^{-7})(c^7 w_{i_2} c^{-7}) = c. \tag{5.82}$$

Here, the first equations in both (5.80) and (5.81), and the first two equations in (5.82) are automatically satisfied due to Lemma 6 with $d = 2$.

With the help of Stembridge's *Maple* package `coxeter` [29], one obtains eight solutions for w_i in (5.80):

$$w_i \in \big\{[1,3,2,1,2,1,3,2],\ [1,2,3,4,3,2,1,2],\ [2,1,2,3], \\ [1,2,1,4,3,2,1,2,1,3,2,1,4,3],\ [1,3,2,1,2,1,3,4],\ [2,1,2,3,2,1,2,4], \\ [2,1,2,1,4,3,2,1,2,1,3,4],\ [1,2,3,2,1,4,3,2,1,2,1,3]\big\}, \tag{5.83}$$

where $\{s_1, s_2, s_3, s_4\}$ is a simple system of generators of H_4, corresponding to the Dynkin diagram displayed in Figure 3, and each of them gives rise to $m/2$ elements of

$\mathrm{Fix}_{NC^m(H_4)}(\phi^p)$ since i ranges from 1 to $m/2$. Furthermore, one obtains 15 solutions for w_i in (5.81):

$$\begin{aligned}w_i \in \big\{&[2],\ [3],\ [4],\ [2,1,2],\ [3,2,1,2,3],\ [1,3,2,1,2,1,3],\ [2,1,2,3,2,1,2],\ [1,2,3,4,3,2,1],\\ &[2,1,3,2,1,2,1,3,2],\ [1,4,3,2,1,2,1,3,4],\ [2,1,2,3,4,3,2,1,2],\\ &[1,2,1,4,3,2,1,2,1,3,2,1,4],\ [1,3,2,1,2,3,4,3,2,1,2,1,3],\\ &[2,1,2,1,4,3,2,1,2,1,3,2,1,2,4],\ [1,3,2,1,4,3,2,1,2,1,3,2,1,4,3]\big\},\end{aligned}$$

each of them giving rise to $m/2$ elements of $\mathrm{Fix}_{NC^m(H_4)}(\phi^p)$ since i ranges from 1 to $m/2$, and one obtains 30 pairs (w_{i_1}, w_{i_2}) of solutions in (5.82):

$$\begin{aligned}(w_{i_1}, w_{i_2}) \in \big\{&([2],\ [2,1,3,2,1,2,1,3,2]),\ ([2],\ [2,1,2,3,4,3,2,1,2]),\ ([3],\ [3,2,1,2,3]),\\ &([3],\ [1,3,2,1,4,3,2,1,2,1,3,2,1,4,3]),\ ([4],\ [1,4,3,2,1,2,1,3,4]),\\ &([4],\ [2,1,2,3,4,3,2,1,2]),\ ([2,1,2],\ [3]),\ ([2,1,2],\ [1,4,3,2,1,2,1,3,4]),\\ &([3,2,1,2,3],\ [2,1,3,2,1,2,1,3,2]),\ ([3,2,1,2,3],\ [1,3,2,1,2,3,4,3,2,1,2,1,3]),\\ &([1,3,2,1,2,1,3],\ [2]),\ ([1,3,2,1,2,1,3],\ [4]),\ ([2,1,2,3,2,1,2],\ [4]),\\ &([2,1,2,3,2,1,2],\ [2,1,2]),\ ([1,2,3,4,3,2,1],\ [2]),\ ([1,2,3,4,3,2,1],\ [3,2,1,2,3]),\\ &([2,1,3,2,1,2,1,3,2],\ [1,3,2,1,2,1,3]),\ ([2,1,3,2,1,2,1,3,2],\ [2,1,2,3,2,1,2]),\\ &([1,4,3,2,1,2,1,3,4],\ [2,1,2,1,4,3,2,1,2,1,3,2,1,2,4]),\\ &([1,4,3,2,1,2,1,3,4],\ [1,3,2,1,4,3,2,1,2,1,3,2,1,4,3]),\\ &([2,1,2,3,4,3,2,1,2],\ [2,1,2,3,2,1,2]),\\ &([2,1,2,3,4,3,2,1,2],\ [2,1,2,1,4,3,2,1,2,1,3,2,1,2,4]),\\ &([1,2,1,4,3,2,1,2,1,3,2,1,4],\ [3]),\\ &([1,2,1,4,3,2,1,2,1,3,2,1,4],\ [1,2,3,4,3,2,1]),\\ &([1,3,2,1,2,3,4,3,2,1,2,1,3],\ [1,3,2,1,2,1,3]),\\ &([1,3,2,1,2,3,4,3,2,1,2,1,3],\ [1,2,3,4,3,2,1]),\\ &([2,1,2,1,4,3,2,1,2,1,3,2,1,2,4],\ [2,1,2]),\\ &([2,1,2,1,4,3,2,1,2,1,3,2,1,2,4],\ [1,2,1,4,3,2,1,2,1,3,2,1,4]),\\ &([1,3,2,1,4,3,2,1,2,1,3,2,1,4,3],\ [1,2,1,4,3,2,1,2,1,3,2,1,4]),\\ &([1,3,2,1,4,3,2,1,2,1,3,2,1,4,3],\ [1,3,2,1,2,3,4,3,2,1,2,1,3])\big\},\end{aligned} \tag{5.84}$$

each of them giving rise to $\binom{m/2}{2}$ elements of $\mathrm{Fix}_{NC^m(H_4)}(\phi^p)$ since $1 \le i_1 < i_2 \le \frac{m}{2}$.

In total, we obtain $1+(15+8)\frac{m}{2}+30\binom{m/2}{2} = \frac{(3m+2)(5m+2)}{4}$ elements in $\mathrm{Fix}_{NC^m(H_4)}(\phi^p)$, which agrees with the limit in (5.67f).

Finally, we turn to (5.67h). By Remark 3, the only choices for h_2 and m_2 to be considered are $h_2 = 2$ and $m_2 = 4$, respectively $h_2 = m_2 = 2$. These correspond to the choices $p = 15m/2$, respectively $p = 15m/4$, out of which only $p = 15m/4$ has not yet been discussed and belongs to the current case. The corresponding action of ϕ^p is given by

$$\begin{aligned}&\phi^p\big((w_0; w_1, \dots, w_m)\big)\\ &\qquad= \big(*; c^4 w_{\frac{m}{4}+1} c^{-4}, c^4 w_{\frac{m}{4}+2} c^{-4}, \dots, c^4 w_m c^{-4}, c^3 w_1 c^{-3}, \dots, c^3 w_{\frac{m}{4}} c^{-3}\big),\end{aligned}$$

so that we have to solve

$$t_1(c^{11}t_1c^{-11})(c^7t_1c^{-7})(c^3t_1c^{-3}) = c$$

for t_1 with $\ell_T(t_1)$. A computation with Stembridge's *Maple* package `coxeter` [29] finds no solution. Hence, the left-hand side of (3.3) is equal to 1, as required.

CASE G_{32}. The degrees are $12, 18, 24, 30$, and hence we have

$$\operatorname{Cat}^m(G_{32};q) = \frac{[30m+30]_q\,[30m+24]_q\,[30m+18]_q\,[30m+12]_q}{[30]_q\,[24]_q\,[18]_q\,[12]_q}.$$

Let ζ be a $30m$-th root of unity. The following cases on the right-hand side of (3.3) do occur:

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{32};q) = m+1, \quad \text{if } \zeta=\zeta_{30},\zeta_{15},\zeta_{10},\zeta_5, \tag{5.85a}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{32};q) = \tfrac{5m+4}{4}, \quad \text{if } \zeta=\zeta_{24},\zeta_8,\ 4\mid m, \tag{5.85b}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{32};q) = \tfrac{5m+3}{3}, \quad \text{if } \zeta=\zeta_{18},\zeta_9,\ 3\mid m, \tag{5.85c}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{32};q) = \tfrac{(5m+4)(5m+2)}{8}, \quad \text{if } \zeta=\zeta_{12},\zeta_4,\ 2\mid m, \tag{5.85d}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{32};q) = \operatorname{Cat}^m(G_{32}), \quad \text{if } \zeta=\zeta_6,\zeta_3,-1,1, \tag{5.85e}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{32};q) = 1, \quad \text{otherwise.} \tag{5.85f}$$

We must now prove that the left-hand side of (3.3) in each case agrees with the values exhibited in (5.85). The only cases not covered by Lemmas 4 and 5 are the ones in (5.85b), (5.85c), (5.85d), and (5.85f).

We begin with the case in (5.85b). By Lemma 3, we are free to choose $p = 5m/4$ if $\zeta=\zeta_{24}$, and we are free to choose $p = 15m/4$ if $\zeta=\zeta_8$. In particular, in all both cases, m must be divisible by 4.

We start with the case that $p = 5m/4$. From (4.1), we infer

$$\begin{aligned}\phi^p\big((w_0;w_1,\dots,w_m)\big)\\ &= \big(*;c^2w_{\frac{3m}{4}+1}c^{-2},c^2w_{\frac{3m}{4}+2}c^{-2},\dots,c^2w_mc^{-2},cw_1c^{-1},\dots,cw_{\frac{3m}{4}}c^{-1}\big).\end{aligned}$$

Supposing that $(w_0;w_1,\dots,w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^2w_{\frac{3m}{4}+i}c^{-2}, \quad i=1,2,\dots,\tfrac{m}{4}, \tag{5.86a}$$

$$w_i = cw_{i-\frac{m}{4}}c^{-1}, \quad i=\tfrac{m}{4}+1,\tfrac{m}{4}+2,\dots,m. \tag{5.86b}$$

There are two distinct possibilities for choosing the w_i's, $1\le i\le m$:

(i) all the w_i's are equal to ε (and $w_0=c$),
(ii) there is an i with $1\le i\le\frac{m}{4}$ such that

$$\ell_T(w_i)=\ell_T(w_{i+\frac{m}{4}})=\ell_T(w_{i+\frac{2m}{4}})=\ell_T(w_{i+\frac{3m}{4}})=1, \tag{5.87a}$$

and the other w_j's, $1\le j\le m$, are equal to ε.

Moreover, since $(w_0;w_1,\dots,w_m)\in NC^m(G_{32})$, we must have

$$w_iw_{i+\frac{m}{4}}w_{i+\frac{2m}{4}}w_{i+\frac{3m}{4}}=c.$$

Together with Equations (5.86)–(5.87), this implies that

$$w_i = c^5 w_i c^{-5} \quad \text{and} \quad w_i (c w_i c^{-1})(c^2 w_i c^{-2})(c^3 w_i c^{-3}) = c. \tag{5.88}$$

With the help of the *GAP* package `CHEVIE` [12], one obtains five solutions for w_i in (5.88):

$$w_i \in \big\{[1],\ [2],\ [3],\ [4],\ [27]\big\}, \tag{5.89}$$

where we have again used the short notation of `CHEVIE` referring to the internal ordering of the roots of G_{32} in `CHEVIE`. Each of the above solutions for w_i gives rise to $m/4$ elements of $\operatorname{Fix}_{NC^m(G_{32})}(\phi^p)$ since i ranges from 1 to $m/4$.

In total, we obtain $1 + 5\frac{m}{4} = \frac{5m+4}{4}$ elements in $\operatorname{Fix}_{NC^m(G_{32})}(\phi^p)$, which agrees with the limit in (5.85b).

In the case that $p = 15m/4$, we infer from (4.1) that

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^4 w_{\frac{m}{4}+1} c^{-4}, c^4 w_{\frac{m}{4}+2} c^{-4}, \dots, c^4 w_m c^{-4}, c^3 w_1 c^{-3}, \dots, c^3 w_{\frac{m}{4}} c^{-3}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^4 w_{\frac{m}{4}+i} c^{-4}, \quad i = 1, 2, \dots, \tfrac{3m}{4}, \tag{5.90a}$$

$$w_i = c^3 w_{i-\frac{3m}{4}} c^{-3}, \quad i = \tfrac{3m}{4}+1, \tfrac{3m}{4}+2, \dots, m. \tag{5.90b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$: either all the w_i's are equal to ε, or there is an i with $1 \le i \le \frac{m}{4}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{4}}) = \ell_T(w_{i+\frac{2m}{4}}) = \ell_T(w_{i+\frac{3m}{4}}) = 1.$$

Writing t_1, t_2, t_3, t_4 for $w_i, w_{i+\frac{m}{4}}, w_{i+\frac{2m}{4}}, w_{i+\frac{3m}{4}}$, in that order, the equations (5.90) reduce to

$$t_1 = c^4 t_2 c^{-4}, \tag{5.91a}$$

$$t_2 = c^4 t_3 c^{-4}, \tag{5.91b}$$

$$t_3 = c^4 t_4 c^{-4}, \tag{5.91c}$$

$$t_4 = c^3 t_1 c^{-3}. \tag{5.91d}$$

One of these equations is in fact superfluous: if we substitute (5.91b)–(5.91d) in (5.91a), then we obtain $t_1 = c^{15} t_1 c^{-15}$ which is automatically satisfied due to Lemma 6 with $d = 2$.

Since $(w_0; w_1, \dots, w_m) \in NC^m(G_{32})$, we must have $t_1 t_2 t_3 t_4 = c$. Combining this with (5.91), we infer that

$$t_1 (c^{11} t_1 c^{-11})(c^7 t_1 c^{-7})(c^3 t_1 c^{-3}) = c. \tag{5.92}$$

Using that $c^5 t_1 c^{-5} = t_1$, due to Lemma 6 with $d = 6$, we see that this equation is equivalent with (5.88). Therefore, we are facing exactly the same enumeration problem here as for $p = 5m/4$, and, consequently, the number of solutions to (5.92) is the same, namely $\frac{5m+4}{4}$, as required.

Next we consider the case in (5.85c). By Lemma 3, we are free to choose $p = 5m/3$ if $\zeta = \zeta_{18}$, and we are free to choose $p = 10m/3$ if $\zeta = \zeta_9$. In particular, in both cases, m must be divisible by 3.

We start with the case that $p = 5m/3$. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) = \big(*; c^2 w_{\frac{m}{3}+1} c^{-2}, c^2 w_{\frac{m}{3}+2} c^{-2}, \dots, c^2 w_m c^{-2}, c w_1 c^{-1}, \dots, c w_{\frac{m}{3}} c^{-1}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^2 w_{\frac{m}{3}+i} c^{-2}, \quad i = 1, 2, \dots, \tfrac{2m}{3}, \tag{5.93a}$$

$$w_i = c w_{i-\frac{2m}{3}} c^{-1}, \quad i = \tfrac{2m}{3}+1, \tfrac{2m}{3}+2, \dots, m. \tag{5.93b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{3}}) = \ell_T(w_{i+\frac{2m}{3}}) = 1, \tag{5.94a}$$

and the other w_j's, $1 \le j \le m$, are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(G_{32})$, we must have $w_i w_{i+\frac{m}{3}} w_{i+\frac{2m}{3}} \le_T c$. Together with Equations (5.93)–(5.94), this implies that

$$w_i = c^5 w_i c^{-5} \quad \text{and} \quad w_i (c^3 w_i c^{-3})(c w_i c^{-1}) \le_T c. \tag{5.95}$$

With the help of the *GAP* package `CHEVIE` [12], one obtains three solutions for w_i in (5.95):

$$w_i \in \big\{[1],\ [2],\ [3],\ [4],\ [27]\big\},$$

where we have again used the short notation of `CHEVIE` referring to the internal ordering of the roots of G_{32} in `CHEVIE`. Each of the above solutions for w_i gives rise to $m/3$ elements of $\mathrm{Fix}_{NC^m(G_{32})}(\phi^p)$ since i ranges from 1 to $m/3$.

In total, we obtain $1 + 5\frac{m}{3} = \frac{5m+3}{3}$ elements in $\mathrm{Fix}_{NC^m(G_{32})}(\phi^p)$, which agrees with the limit in (5.85c).

In the case that $p = 10m/3$, we infer from (4.1) that

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^4 w_{\frac{2m}{3}+1} c^{-4}, c^4 w_{\frac{2m}{3}+2} c^{-4}, \dots, c^4 w_m c^{-4}, c^3 w_1 c^{-3}, \dots, c^3 w_{\frac{2m}{3}} c^{-3}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^4 w_{\frac{2m}{3}+i} c^{-4}, \quad i = 1, 2, \dots, \tfrac{m}{3}, \tag{5.96a}$$

$$w_i = c^3 w_{i-\frac{m}{3}} c^{-3}, \quad i = \tfrac{m}{3}+1, \tfrac{m}{3}+2, \dots, m. \tag{5.96b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$: either all the w_i's are equal to ε, or there is an i with $1 \le i \le \frac{m}{3}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{3}}) = \ell_T(w_{i+\frac{2m}{3}}) = 1.$$

Writing t_1, t_2, t_3 for $w_i, w_{i+\frac{m}{3}}, w_{i+\frac{2m}{3}}$, in that order, the equations (5.96) reduce to

$$t_1 = c^4 t_3 c^{-4}, \tag{5.97a}$$

$$t_2 = c^3 t_1 c^{-3}, \tag{5.97b}$$

$$t_3 = c^3 t_2 c^{-3}. \tag{5.97c}$$

One of these equations is in fact superfluous: if we substitute (5.97b) and (5.97c) in (5.97a), then we obtain $t_1 = c^{10} t_1 c^{-10}$ which is automatically satisfied due to Lemma 6 with $d = 2$.

Since $(w_0; w_1, \dots, w_m) \in NC^m(G_{32})$, we must have $t_1 t_2 t_3 \le_T c$. Combining this with (5.97), we infer that

$$t_1 (c^3 t_1 c^{-3})(c^6 t_1 c^{-6}) \le_T c. \tag{5.98}$$

Using that $c^5t_1c^{-5} = t_1$, due to Lemma 6 with $d = 4$, we see that this equation is equivalent with (5.95). Therefore, we are facing exactly the same enumeration problem here as for $p = 5m/3$, and, consequently, the number of solutions to (5.98) is the same, namely $\frac{5m+3}{3}$, as required.

Next we discuss the case in (5.85d). By Lemma 3, we are free to choose $p = 5m/2$ if $\zeta = \zeta_{12}$, and we are free to choose $p = 15m/2$ if $\zeta = \zeta_4$. In particular, m must be divisible by 2. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = (*; c^3w_{\frac{m}{2}+1}c^{-3}, c^3w_{\frac{m}{2}+2}c^{-3}, \dots, c^3w_mc^{-3}, c^2w_1c^{-2}, \dots, c^2w_{\frac{m}{2}}c^{-2}\big). \quad (5.99)$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^3w_{\frac{m}{2}+i}c^{-3}, \quad i = 1, 2, \dots, \tfrac{m}{2}, \quad (5.100a)$$

$$w_i = c^2w_{i-\frac{m}{2}}c^{-2}, \quad i = \tfrac{m}{2}+1, \tfrac{m}{2}+2, \dots, m. \quad (5.100b)$$

There are four distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 2, \quad (5.101a)$$

and all other w_j's are equal to ε,

(iii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 1, \quad (5.101b)$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(iv) there are i_1 and i_2 with $1 \le i_1 < i_2 \le \frac{m}{2}$ such that

$$\ell_T(w_{i_1}) = \ell_T(w_{i_2}) = \ell_T(w_{i_1+\frac{m}{2}}) = \ell_T(w_{i_2+\frac{m}{2}}) = 1, \quad (5.101c)$$

and all other w_j's are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(G_{32})$, we must have $w_iw_{i+\frac{m}{2}} \le_T c$, respectively $w_{i_1}w_{i_2}w_{i_1+\frac{m}{2}}w_{i_2+\frac{m}{2}} = c$. Together with Equations (5.100)–(5.101), this implies that

$$w_i = c^3w_ic^{-3} \quad \text{and} \quad w_i(c^2w_ic^{-2}) = c, \quad (5.102)$$

respectively that

$$w_i = c^3w_ic^{-3}, \quad w_i(c^2w_ic^{-2}) \le_T c, \quad \text{and} \quad \ell_T(w_i) = 1, \quad (5.103)$$

respectively that

$$w_{i_1} = c^3w_{i_1}c^{-3}, \quad w_{i_2} = c^3w_{i_2}c^{-3}, \\ w_{i_1}w_{i_2}(c^2w_{i_1}c^{-2})(c^2w_{i_2}c^{-2}) = c, \quad \text{and} \quad \ell_T(w_{i_1}) = \ell_T(w_{i_2}) = 1. \quad (5.104)$$

With the help of the *GAP* package `CHEVIE` [12], one obtains ten solutions for w_i in (5.102):

$$w_i \in \big\{[3,4],\ [1,20],\ [4,7],\ [1,2],\ [2,3],\ [4,27],\ [3,23],\ [1,23],\ [2,16],\ [12,27]\big\}, \quad (5.105)$$

where we have again used the short notation of `CHEVIE` referring to the internal ordering of the roots of G_{32} in `CHEVIE`, one obtains solutions for w_i in (5.103):

$$w_i \in \big\{[4]\ \ [3]\ \ [20]\ \ [1]\ \ [7]\ \ [2]\ \ [16]\ \ [12]\ \ [27]\ \ [23]\big\},$$

each of them giving rise to $m/2$ elements of $\operatorname{Fix}_{NC^m(G_{32})}(\phi^p)$ since i ranges from 1 to $m/2$, and one obtains 25 pairs (w_{i_1}, w_{i_2}) satisfying (5.104):

$$\begin{multline}(w_{i_1}, w_{i_2}) \in \big\{([4], [20]),\ ([4], [7]),\ ([4], [27]),\ ([3], [4]),\ ([3], [12]),\ ([3], [23]),\ ([20], [3]),\\ ([20], [1]),\ ([1], [20]),\ ([1], [2]),\ ([1], [23]),\ ([7], [4]),\ ([7], [1]),\ ([2], [3]),\ ([2], [7]),\\ ([2], [16]),\ ([16], [4]),\ ([16], [2]),\ ([12], [2]),\ ([12], [27]),\ ([27], [1]),\ ([27], [16]),\\ ([27], [12]),\ ([23], [3]),\ ([23], [27])\big\}, \quad (5.106)\end{multline}$$

each of them giving rise to $\binom{m/2}{2}$ elements of $\operatorname{Fix}_{NC^m(G_{32})}(\phi^p)$ since $1 \le i_1 < i_2 \le \frac{m}{2}$.

In total, we obtain $1 + 20\frac{m}{2} + 25\binom{m/2}{2} = \frac{(5m+4)(5m+2)}{8}$ elements in $\operatorname{Fix}_{NC^m(G_{32})}(\phi^p)$, which agrees with the limit in (5.85d).

If $p = 15m/2$, then, from (4.1), we infer

$$\begin{multline}\phi^p\big((w_0; w_1, \dots, w_m)\big)\\ = \big(*; c^8 w_{\frac{m}{2}+1} c^{-8}, c^8 w_{\frac{m}{2}+2} c^{-8}, \dots, c^8 w_m c^{-8}, c^7 w_1 c^{-7}, \dots, c^7 w_{\frac{m}{2}} c^{-7}\big).\end{multline}$$

Using that $c^5 w c^{-5} = w$ for all $w \in NC(G_{32}$, due to Lemma 6 with $d = 6$, we see that this action is identical with the one in (5.99). Therefore, we are facing exactly the same enumeration problem here as for $p = 5m/2$, and, consequently, the number of elements in $\operatorname{Fix}_{NC^m(G_{32})}(\phi^p)$ is the same, namely $\frac{(5m+4)(5m+2)}{8}$, as required.

Finally, we turn to (5.85f). By Remark 3, the only choices for h_2 and m_2 to be considered are $h_2 = 2$ and $m_2 = 4$, $h_2 = m_2 = 2$, $h_2 = m_2 = 3$, $h_2 = 6$ and $m_2 = 4$, $h_2 = 6$ and $m_2 = 3$, respectively $h_2 = 6$ and $m_2 = 2$. These correspond to the choices $p = 15m/4$, $p = 15m/2$, $p = 10m/3$, $p = 5m/4$, $p = 5m/3$, respectively $p = 5m/2$, all of which have already been discussed as they do not belong to (5.85f). Hence, (3.3) must necessarily hold, as required.

Case G_{33}. The degrees are $4, 6, 10, 12, 18$, and hence we have

$$\operatorname{Cat}^m(G_{33}; q) = \frac{[18m+18]_q\,[18m+12]_q\,[18m+10]_q\,[18m+6]_q\,[18m+4]_q}{[18]_q\,[12]_q\,[10]_q\,[6]_q\,[4]_q}.$$

Let ζ be a $18m$-th root of unity. The following cases on the right-hand side of (3.3) do occur:

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{33}; q) = m+1, \quad \text{if } \zeta = \zeta_{18}, \zeta_9, \tag{5.107a}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{33}; q) = \tfrac{3m+2}{2}, \quad \text{if } \zeta = \zeta_{12},\ 2 \mid m, \tag{5.107b}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{33}; q) = \tfrac{9m+5}{5}, \quad \text{if } \zeta = \zeta_{10}, \zeta_5,\ 5 \mid m, \tag{5.107c}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{33}; q) = \tfrac{(m+1)(3m+2)(3m+1)}{2}, \quad \text{if } \zeta = \zeta_6, \zeta_3, \tag{5.107d}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{33}; q) = \tfrac{(3m+2)(9m+2)}{4}, \quad \text{if } \zeta = \zeta_4,\ 2 \mid m, \tag{5.107e}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{33}; q) = \operatorname{Cat}^m(G_{33}), \quad \text{if } \zeta = -1 \text{ or } \zeta = 1, \tag{5.107f}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{33}; q) = 1, \quad \text{otherwise}. \tag{5.107g}$$

We must now prove that the left-hand side of (3.3) in each case agrees with the values exhibited in (5.107). The only cases not covered by Lemmas 4 and 5 are the ones in (5.107b), (5.107c), (5.107e), and (5.107g).

We begin with the case in (5.107b). By Lemma 3, we are free to choose $p = 3m/2$. In particular, m must be divisible by 2. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) = (*; c^2 w_{\frac{m}{2}+1} c^{-2}, c^2 w_{\frac{m}{2}+2} c^{-2}, \dots, c^2 w_m c^{-2}, c w_1 c^{-1}, \dots, c w_{\frac{m}{2}} c^{-1}).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^2 w_{\frac{m}{2}+i} c^{-2}, \quad i = 1, 2, \dots, \tfrac{m}{2}, \tag{5.108a}$$

$$w_i = c w_{i-\frac{m}{2}} c^{-1}, \quad i = \tfrac{m}{2}+1, \tfrac{m}{2}+2, \dots, m. \tag{5.108b}$$

There are four distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 2, \tag{5.109a}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(iii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 1, \tag{5.109b}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(iv) there are i_1 and i_2 with $1 \le i_1 < i_2 \le \frac{m}{2}$ such that

$$\ell_T(w_{i_1}) = \ell_T(w_{i_2}) = \ell_T(w_{i_1+\frac{m}{2}}) = \ell_T(w_{i_2+\frac{m}{2}}) = 1, \tag{5.109c}$$

and the other w_j's, $1 \le j \le m$, are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(G_{33})$, we must have $w_i w_{i+\frac{m}{2}} \le_T c$, respectively $w_{i_1} w_{i_2} w_{i_1+\frac{m}{2}} w_{i_2+\frac{m}{2}} \le_T c$. Together with Equations (5.108)–(5.109), this implies that

$$w_i = c^3 w_i c^{-3} \quad \text{and} \quad w_i(c w_i c^{-1}) \le_T c, \quad \text{and} \quad \ell_T(w_i) = 2, \tag{5.110}$$

respectively that

$$w_i = c^3 w_i c^{-3}, \quad w_i(c w_i c^{-1}) \le_T c, \quad \text{and} \quad \ell_T(w_i) = 1, \tag{5.111}$$

respectively that

$$w_{i_1} = c^3 w_{i_1} c^{-3}, \quad w_{i_1}(c w_{i_1} c^{-1}) \le_T c, \quad \text{and} \quad \ell_T(w_{i_1}) = 1. \tag{5.112}$$

With the help of the *GAP* package `CHEVIE` [12], one obtains three solutions for w_i in (5.110):

$$w_i \in \big\{[1, 44],\ [2, 4],\ [5, 42]\big\},$$

where we have again used the short notation of `CHEVIE` referring to the internal ordering of the roots of G_{33} in `CHEVIE`. Each of them gives rise to $m/2$ elements of $\mathrm{Fix}_{NC^m(G_{33})}(\phi^p)$ since i ranges from 1 to $m/2$.

There are no solutions to (5.111) and to (5.112).

In total, we obtain $1 + 3\frac{m}{2} = \frac{3m+2}{2}$ elements in $\mathrm{Fix}_{NC^m(G_{33})}(\phi^p)$, which agrees with the limit in (5.107b).

Next we turn to the case in (5.107c). By Lemma 3, we are free to choose $p = 9m/5$ if $\zeta = \zeta_{10}$, and we are free to choose $p = 18m/5$ if $\zeta = \zeta_5$. In particular, in all both cases, m must be divisible by 5.

We start with the case that $p = 9m/5$. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^2 w_{\frac{m}{5}+1} c^{-2}, c^2 w_{\frac{m}{5}+2} c^{-2}, \dots, c^2 w_m c^{-2}, c w_1 c^{-1}, \dots, c w_{\frac{m}{5}} c^{-1}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^2 w_{\frac{m}{5}+i} c^{-2}, \quad i = 1, 2, \dots, \tfrac{4m}{5}, \tag{5.113a}$$
$$w_i = c w_{i-\frac{4m}{5}} c^{-1}, \quad i = \tfrac{4m}{5}+1, \tfrac{4m}{5}+2, \dots, m. \tag{5.113b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),
(ii) there is an i with $1 \le i \le \frac{m}{5}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{5}}) = \ell_T(w_{i+\frac{2m}{5}}) = \ell_T(w_{i+\frac{3m}{5}}) = \ell_T(w_{i+\frac{4m}{5}}) = 1, \tag{5.114a}$$

and the other w_j's, $1 \le j \le m$, are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(G_{33})$, we must have

$$w_i w_{i+\frac{m}{5}} w_{i+\frac{2m}{5}} w_{i+\frac{3m}{5}} w_{i+\frac{4m}{5}} = c.$$

Together with Equations (5.113)–(5.114), this implies that

$$w_i = c^9 w_i c^{-9} \quad \text{and} \quad w_i (c^7 w_i c^{-7})(c^5 w_i c^{-5})(c^3 w_i c^{-3})(c w_i c^{-1}) = c. \tag{5.115}$$

With the help of the *GAP* package `CHEVIE` [12], one obtains nine solutions for w_i in (5.115):

$$w_i \in \big\{[5], [4], [1], [2], [10], [27], [42], [44], [52]\big\}, \tag{5.116}$$

where we have again used the short notation of `CHEVIE` referring to the internal ordering of the roots of G_{33} in `CHEVIE`. Each of the above solutions for w_i gives rise to $m/5$ elements of $\operatorname{Fix}_{NC^m(G_{33})}(\phi^p)$ since i ranges from 1 to $m/5$.

In total, we obtain $1 + 9\frac{m}{5} = \frac{9m+5}{5}$ elements in $\operatorname{Fix}_{NC^m(G_{33})}(\phi^p)$, which agrees with the limit in (5.107c).

In the case that $p = 18m/5$, we infer from (4.1) that

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^4 w_{\frac{2m}{5}+1} c^{-4}, c^4 w_{\frac{2m}{5}+2} c^{-4}, \dots, c^4 w_m c^{-4}, c^3 w_1 c^{-3}, \dots, c^3 w_{\frac{2m}{5}} c^{-3}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^4 w_{\frac{2m}{5}+i} c^{-4}, \quad i = 1, 2, \dots, \tfrac{3m}{5}, \tag{5.117a}$$
$$w_i = c^3 w_{i-\frac{3m}{5}} c^{-3}, \quad i = \tfrac{3m}{5}+1, \tfrac{3m}{5}+2, \dots, m. \tag{5.117b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$: either all the w_i's are equal to ε, or there is an i with $1 \le i \le \frac{m}{5}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{5}}) = \ell_T(w_{i+\frac{2m}{5}}) = \ell_T(w_{i+\frac{3m}{5}}) = \ell_T(w_{i+\frac{4m}{5}}) = 1.$$

Writing t_1, t_2, t_3, t_4, t_5 for $w_i, w_{i+\frac{m}{5}}, w_{i+\frac{2m}{5}}, w_{i+\frac{3m}{5}}, w_{i+\frac{4m}{5}}$, in that order, the equations (5.117) reduce to

$$t_1 = c^4 t_3 c^{-4}, \tag{5.118a}$$
$$t_2 = c^4 t_4 c^{-4}, \tag{5.118b}$$
$$t_3 = c^4 t_5 c^{-4}, \tag{5.118c}$$
$$t_4 = c^3 t_1 c^{-3}, \tag{5.118d}$$
$$t_5 = c^3 t_2 c^{-3}. \tag{5.118e}$$

One of these equations is in fact superfluous: if we substitute (5.118b)–(5.118e) in (5.118a), then we obtain $t_1 = c^{18} t_1 c^{-18}$ which is automatically satisfied since $c^{18} = \varepsilon$.

Since $(w_0; w_1, \dots, w_m) \in NC^m(G_{33})$, we must have $t_1 t_2 t_3 t_4 t_5 = c$. Combining this with (5.118), we infer that

$$t_1 (c^7 t_1 c^{-7})(c^{14} t_1 c^{-14})(c^3 t_1 c^{-3})(c^{10} t_1 c^{-10}) = c. \tag{5.119}$$

Using that $c^9 t_1 c^{-9} = t_1$, due to Lemma 6 with $d = 2$, we see that this equation is equivalent with (5.115). Therefore, we are facing exactly the same enumeration problem here as for $p = 9m/5$, and, consequently, the number of solutions to (5.119) is the same, namely $\frac{9m+5}{5}$, as required.

Next we consider the case in (5.107e). By Lemma 3, we are free to choose $p = 9m/2$. In particular, m must be divisible by 2. From (4.1), we infer

$$\begin{aligned}&\phi^p\big((w_0; w_1, \dots, w_m)\big)\\ &\qquad = \big(*; c^5 w_{\frac{m}{2}+1} c^{-5}, c^5 w_{\frac{m}{2}+2} c^{-5}, \dots, c^5 w_m c^{-5}, c^4 w_1 c^{-4}, \dots, c^4 w_{\frac{m}{2}} c^{-4}\big).\end{aligned}$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^5 w_{\frac{m}{2}+i} c^{-5}, \quad i = 1, 2, \dots, \tfrac{m}{2}, \tag{5.120a}$$
$$w_i = c^4 w_{i-\frac{m}{2}} c^{-4}, \quad i = \tfrac{m}{2}+1, \tfrac{m}{2}+2, \dots, m. \tag{5.120b}$$

There are four distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{2}$ such that
$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 2, \tag{5.121a}$$
and the other w_j's, $1 \le j \le m$, are equal to ε,

(iii) there is an i with $1 \le i \le \frac{m}{2}$ such that
$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 1, \tag{5.121b}$$
and the other w_j's, $1 \le j \le m$, are equal to ε,

(iv) there are i_1 and i_2 with $1 \le i_1 < i_2 \le \frac{m}{2}$ such that
$$\ell_T(w_{i_1}) = \ell_T(w_{i_2}) = \ell_T(w_{i_1+\frac{m}{2}}) = \ell_T(w_{i_2+\frac{m}{2}}) = 1, \tag{5.121c}$$
and the other w_j's, $1 \le j \le m$, are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(G_{33})$, we must have $w_i w_{i+\frac{m}{2}} \le_T c$, respectively $w_{i_1} w_{i_2} w_{i_1+\frac{m}{2}} w_{i_2+\frac{m}{2}} \le_T c$. Together with Equations (5.120)–(5.121), this implies that

$$w_i = c^9 w_i c^{-9} \quad \text{and} \quad w_i (c^4 w_i c^{-4}) \le_T c, \quad \text{and} \quad \ell_T(w_i) = 2, \tag{5.122}$$

respectively that

$$w_i = c^9 w_i c^{-9}, \quad w_i(c^4 w_i c^{-4}) \le_T c, \quad \text{and} \quad \ell_T(w_i) = 1, \tag{5.123}$$

respectively that

$$\begin{multline} w_{i_1} = c^9 w_{i_1} c^{-9}, \quad w_{i_2} = c^9 w_{i_2} c^{-9}, \\ w_{i_1} w_{i_2} (c^4 w_{i_1} c^{-4})(c^4 w_{i_2} c^{-4}) \le_T c, \quad \text{and} \quad \ell_T(w_{i_1}) = \ell_T(w_{i_2}) = 1. \end{multline} \tag{5.124}$$

With the help of the *GAP* package `CHEVIE` [12], one obtains 21 solutions for w_i in (5.122):

$$\begin{multline} w_i \in \big\{[4,5],\ [1,20],\ [5,7],\ [1,2],\ [2,4],\ [10,259],\ [27,208],\ [27,44],\ [4,39],\ [5,42],\ [1,39], \\ [5,52],\ [10,208],\ [1,44],\ [42,113],\ [4,113],\ [2,27],\ [10,42],\ [2,49],\ [52,61],\ [44,123]\big\}, \end{multline}$$

where we have again used the short notation of `CHEVIE` referring to the internal ordering of the roots of G_{33} in `CHEVIE`, one obtains 18 solutions for w_i in (5.123):

$$\begin{multline} w_i \in \big\{[5],\ [4],\ [20],\ [208],\ [259],\ [1],\ [2],\ [7],\ [10],\ [27], \\ [39],\ [42],\ [49],\ [44],\ [52],\ [113],\ [61],\ [123]\big\}, \end{multline}$$

each of them giving rise to $m/2$ elements of $\operatorname{Fix}_{NC^m(G_{33})}(\phi^p)$ since i ranges from 1 to $m/2$, and one obtains 54 pairs (w_{i_1}, w_{i_2}) satisfying (5.124):

$$\begin{gathered} (w_{i_1}, w_{i_2}) \in \big\{([5],[20]),\ ([5],[7]),\ ([5],[42]),\ ([5],[52]),\ ([4],[5]),\ ([4],[10]),\ ([4],[39]),\ ([4],[113]),\ ([20],[4]), \\ ([20],[1]),\ ([208],[27]),\ ([208],[52]),\ ([259],[10]),\ ([259],[27]),\ ([1],[20]),\ ([1],[2]),\ ([1],[39]),\ ([1],[44]),\ ([2],[4]), \\ ([2],[7]),\ ([2],[27]),\ ([2],[49]),\ ([7],[5]),\ ([7],[1]),\ ([10],[208]),\ ([10],[259]),\ ([10],[2]),\ ([10],[42]),\ ([27],[5]), \\ ([27],[208]),\ ([27],[44]),\ ([27],[61]),\ ([39],[4]),\ ([39],[42]),\ ([42],[1]),\ ([42],[27]),\ ([42],[113]),\ ([42],[123]), \\ ([49],[5]),\ ([49],[2]),\ ([44],[4]),\ ([44],[259]),\ ([44],[52]),\ ([44],[123]),\ ([52],[1]),\ ([52],[10]),\ ([52],[49]), \\ ([52],[61]),\ ([113],[42]),\ ([113],[44]),\ ([61],[2]),\ ([61],[52]),\ ([123],[10]),\ ([123],[44])\big\}, \end{gathered}$$

each of them giving rise to $\binom{m/2}{2}$ elements of $\operatorname{Fix}_{NC^m(G_{33})}(\phi^p)$ since $1 \le i_1 < i_2 \le m$.

In total, we obtain $1+(21+18)\frac{m}{2}+54\binom{m/2}{2} = \frac{(3m+2)(9m+2)}{4}$ elements in $\operatorname{Fix}_{NC^m(G_{33})}(\phi^p)$, which agrees with the limit in (5.107e).

Finally, we turn to (5.107g). By Remark 3, the only choices for h_2 and m_2 to be considered are $h_2 = 1$ and $m_2 = 5$, $h_2 = 2$ and $m_2 = 5$, $h_2 = 2$ and $m_2 = 4$, respectively $h_2 = m_2 = 2$. These correspond to the choices $p = 18m/5$, $p = 9m/5$, $p = 9m/4$, respectively $p = 9m/2$, out of which only $p = 9m/4$ has not yet been discussed and belongs to the current case. The corresponding action of ϕ^p is given by

$$\begin{multline} \phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^3 w_{\frac{3m}{4}+1} c^{-3}, c^3 w_{\frac{3m}{4}+2} c^{-3}, \dots, c^3 w_m c^{-3}, c^2 w_1 c^{-2}, \dots, c^2 w_{\frac{3m}{4}} c^{-2}\big), \end{multline}$$

so that we have to solve

$$t_1 (c^2 t_1 c^{-2})(c^4 t_1 c^{-4})(c^6 t_1 c^{-6}) \le_T c$$

for t_1 with $\ell_T(t_1)$. A computation with the help of the *GAP* package `CHEVIE` [12] finds no solution. Hence, the left-hand side of (3.3) is equal to 1, as required.

Case G_{34}. The degrees are $6, 12, 18, 24, 30, 42$, and hence we have

$$\operatorname{Cat}^m(G_{34};q) = \frac{[42m+42]_q\,[42m+30]_q\,[42m+24]_q}{[42]_q\,[30]_q\,[24]_q} \times \frac{[42m+18]_q\,[42m+12]_q\,[42m+6]_q}{[18]_q\,[12]_q\,[6]_q}.$$

Let ζ be a $42m$-th root of unity. The following cases on the right-hand side of (3.3) do occur:

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{34};q) = m+1, \quad \text{if } \zeta=\zeta_{42},\zeta_{21},\zeta_{14},\zeta_7, \tag{5.125a}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{34};q) = \tfrac{7m+5}{5}, \quad \text{if } \zeta=\zeta_{30},\zeta_{15},\zeta_{10},\zeta_5,\ 5\mid m, \tag{5.125b}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{34};q) = \tfrac{7m+4}{4}, \quad \text{if } \zeta=\zeta_{24},\zeta_8,\ 4\mid m, \tag{5.125c}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{34};q) = \tfrac{7m+3}{3}, \quad \text{if } \zeta=\zeta_{18},\zeta_9,\ 3\mid m, \tag{5.125d}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{34};q) = \tfrac{(7m+4)(7m+2)}{8}, \quad \text{if } \zeta=\zeta_{12},\zeta_4,\ 2\mid m, \tag{5.125e}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{34};q) = \operatorname{Cat}^m(G_{34}), \quad \text{if } \zeta=\zeta_6,\zeta_3,-1,1, \tag{5.125f}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{34};q) = 1, \quad \text{otherwise.} \tag{5.125g}$$

We must now prove that the left-hand side of (3.3) in each case agrees with the values exhibited in (5.125). The only cases not covered by Lemmas 4 and 5 are the ones in (5.125b), (5.125c), (5.125d), (5.125e), and (5.125g).

We begin with the case in (5.125b). By Lemma 3, we are free to choose $p = 7m/5$ if $\zeta=\zeta_{30}$, we are free to choose $p=14m/5$ if $\zeta=\zeta_{15}$, we are free to choose $p=21m/5$ if $\zeta=\zeta_{10}$, and we are free to choose $p=42m/5$ if $\zeta=\zeta_5$. In particular, in all cases, m must be divisible by 5.

We start with the case that $p=7m/5$. From (4.1), we infer

$$\phi^p\big((w_0;w_1,\dots,w_m)\big) = \big(*;c^2w_{\frac{3m}{5}+1}c^{-2},c^2w_{\frac{3m}{5}+2}c^{-2},\dots,c^2w_mc^{-2},cw_1c^{-1},\dots,cw_{\frac{3m}{5}}c^{-1}\big).$$

Supposing that $(w_0;w_1,\dots,w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^2w_{\frac{3m}{5}+i}c^{-2}, \quad i=1,2,\dots,\tfrac{2m}{5}, \tag{5.126a}$$

$$w_i = cw_{i-\frac{2m}{5}}c^{-1}, \quad i=\tfrac{2m}{5}+1,\tfrac{2m}{5}+2,\dots,m. \tag{5.126b}$$

There are two distinct possibilities for choosing the w_i's, $1\le i\le m$:

(i) all the w_i's are equal to ε (and $w_0=c$),
(ii) there is an i with $1\le i\le\frac{m}{5}$ such that

$$\ell_T(w_i)=\ell_T(w_{i+\frac{m}{5}})=\ell_T(w_{i+\frac{2m}{5}})=\ell_T(w_{i+\frac{3m}{5}})=\ell_T(w_{i+\frac{4m}{5}})=1, \tag{5.127a}$$

and the other w_j's, $1\le j\le m$, are equal to ε.

Moreover, since $(w_0;w_1,\dots,w_m)\in NC^m(G_{34})$, we must have

$$w_iw_{i+\frac{m}{5}}w_{i+\frac{2m}{5}}w_{i+\frac{3m}{5}}w_{i+\frac{4m}{5}} \le_T c.$$

Together with Equations (5.126)–(5.127), this implies that

$$w_i = c^7 w_i c^{-7} \quad \text{and} \quad w_i(c^4 w_i c^{-4})(c w_i c^{-1})(c^5 w_i c^{-5})(c^2 w_i c^{-2}) \le_T c. \tag{5.128}$$

With the help of the *GAP* package `CHEVIE` [12], one obtains seven solutions for w_i in (5.128):

$$w_i \in \big\{[4],\ [5],\ [6],\ [2],\ [11],\ [44],\ [63],\ [74]\big\},$$

where we have again used the short notation of `CHEVIE` referring to the internal ordering of the roots of G_{34} in `CHEVIE`. Each of the above solutions for w_i gives rise to $m/5$ elements of $\operatorname{Fix}_{NC^m(G_{34})}(\phi^p)$ since i ranges from 1 to $m/5$.

In total, we obtain $1 + 7\frac{m}{5} = \frac{7m+5}{5}$ elements in $\operatorname{Fix}_{NC^m(G_{34})}(\phi^p)$, which agrees with the limit in (5.125b).

In the case that $p = 14m/5$, we infer from (4.1) that

$$\begin{aligned}\phi^p\big((w_0; w_1, \dots, w_m)\big)&\\ &= (*; c^3 w_{\frac{m}{5}+1} c^{-3}, c^3 w_{\frac{m}{5}+2} c^{-3}, \dots, c^3 w_m c^{-3}, c^2 w_1 c^{-2}, \dots, c^2 w_{\frac{m}{5}} c^{-2}\big).\end{aligned}$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^3 w_{\frac{m}{5}+i} c^{-3}, \quad i = 1, 2, \dots, \tfrac{4m}{5}, \tag{5.129a}$$

$$w_i = c^2 w_{i-\frac{4m}{5}} c^{-2}, \quad i = \tfrac{4m}{5}+1, \tfrac{4m}{5}+2, \dots, m. \tag{5.129b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$: either all the w_i's are equal to ε, or there is an i with $1 \le i \le \frac{m}{5}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{5}}) = \ell_T(w_{i+\frac{2m}{5}}) = \ell_T(w_{i+\frac{3m}{5}}) = \ell_T(w_{i+\frac{4m}{5}}) = 1.$$

Writing t_1, t_2, t_3, t_4, t_5 for $w_i, w_{i+\frac{m}{5}}, w_{i+\frac{2m}{5}}, w_{i+\frac{3m}{5}}, w_{i+\frac{4m}{5}}$, in that order, the equations (5.129) reduce to

$$t_1 = c^3 t_2 c^{-3}, \tag{5.130a}$$

$$t_2 = c^3 t_3 c^{-3}, \tag{5.130b}$$

$$t_3 = c^3 t_4 c^{-3}, \tag{5.130c}$$

$$t_4 = c^3 t_5 c^{-3}, \tag{5.130d}$$

$$t_5 = c^2 t_1 c^{-2}. \tag{5.130e}$$

One of these equations is in fact superfluous: if we substitute (5.130b)–(5.130e) in (5.130a), then we obtain $t_1 = c^{14} t_1 c^{-14}$ which is automatically satisfied due to Lemma 6 with $d = 3$.

Since $(w_0; w_1, \dots, w_m) \in NC^m(G_{34})$, we must have $t_1 t_2 t_3 t_4 t_5 \le_T c$. Combining this with (5.130), we infer that

$$t_1(c^{11} t_1 c^{-11})(c^8 t_1 c^{-8})(c^5 t_1 c^{-5})(c^2 t_1 c^{-2}) \le_T c. \tag{5.131}$$

Using that $c^7 t_1 c^{-7} = t_1$, due to Lemma 6 with $d = 6$, we see that this equation is equivalent with (5.128). Therefore, we are facing exactly the same enumeration problem here as for $p = 7m/5$, and, consequently, the number of solutions to (5.131) is the same, namely $\frac{7m+5}{5}$, as required.

In the case that $p = 21m/5$, we infer from (4.1) that

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = (*; c^5 w_{\frac{4m}{5}+1} c^{-5}, c^5 w_{\frac{4m}{5}+2} c^{-5}, \dots, c^5 w_m c^{-5}, c^4 w_1 c^{-4}, \dots, c^4 w_{\frac{4m}{5}} c^{-4}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^5 w_{\frac{4m}{5}+i} c^{-5}, \quad i = 1, 2, \dots, \tfrac{m}{5}, \tag{5.132a}$$

$$w_i = c^4 w_{i-\frac{m}{5}} c^{-4}, \quad i = \tfrac{m}{5}+1, \tfrac{m}{5}+2, \dots, m. \tag{5.132b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$: either all the w_i's are equal to ε, or there is an i with $1 \le i \le \frac{m}{5}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{5}}) = \ell_T(w_{i+\frac{2m}{5}}) = \ell_T(w_{i+\frac{3m}{5}}) = \ell_T(w_{i+\frac{4m}{5}}) = 1.$$

Writing t_1, t_2, t_3, t_4, t_5 for $w_i, w_{i+\frac{m}{5}}, w_{i+\frac{2m}{5}}, w_{i+\frac{3m}{5}}, w_{i+\frac{4m}{5}}$, in that order, the equations (5.132) reduce to

$$t_1 = c^5 t_5 c^{-5}, \tag{5.133a}$$

$$t_2 = c^4 t_1 c^{-4}, \tag{5.133b}$$

$$t_3 = c^4 t_2 c^{-4}, \tag{5.133c}$$

$$t_4 = c^4 t_3 c^{-4}, \tag{5.133d}$$

$$t_5 = c^4 t_4 c^{-4}. \tag{5.133e}$$

One of these equations is in fact superfluous: if we substitute (5.133b)–(5.133e) in (5.133a), then we obtain $t_1 = c^{21} t_1 c^{-21}$ which is automatically satisfied due to Lemma 6 with $d = 6$.

Since $(w_0; w_1, \dots, w_m) \in NC^m(G_{34})$, we must have $t_1 t_2 t_3 t_4 t_5 \le_T c$. Combining this with (5.133), we infer that

$$t_1 (c^4 t_1 c^{-4})(c^8 t_1 c^{-8})(c^{12} t_1 c^{-12})(c^{16} t_1 c^{-16}) \le_T c. \tag{5.134}$$

Using that $c^7 t_1 c^{-7} = t_1$, due to Lemma 6 with $d = 6$, we see that this equation is equivalent with (5.128). Therefore, we are facing exactly the same enumeration problem here as for $p = 7m/5$, and, consequently, the number of solutions to (5.134) is the same, namely $\frac{7m+5}{5}$, as required.

In the case that $p = 42m/5$, we infer from (4.1) that

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = (*; c^9 w_{\frac{3m}{5}+1} c^{-9}, c^9 w_{\frac{3m}{5}+2} c^{-9}, \dots, c^9 w_m c^{-9}, c^8 w_1 c^{-8}, \dots, c^8 w_{\frac{3m}{5}} c^{-8}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^9 w_{\frac{3m}{5}+i} c^{-9}, \quad i = 1, 2, \dots, \tfrac{2m}{5}, \tag{5.135a}$$

$$w_i = c^8 w_{i-\frac{2m}{5}} c^{-8}, \quad i = \tfrac{2m}{5}+1, \tfrac{2m}{5}+2, \dots, m. \tag{5.135b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$: either all the w_i's are equal to ε, or there is an i with $1 \le i \le \frac{m}{5}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{5}}) = \ell_T(w_{i+\frac{2m}{5}}) = \ell_T(w_{i+\frac{3m}{5}}) = \ell_T(w_{i+\frac{4m}{5}}) = 1.$$

Writing t_1, t_2, t_3, t_4, t_5 for $w_i, w_{i+\frac{m}{5}}, w_{i+\frac{2m}{5}}, w_{i+\frac{3m}{5}}, w_{i+\frac{4m}{5}}$, in that order, the equations (5.135) reduce to

$$t_1 = c^9 t_4 c^{-9}, \tag{5.136a}$$

$$t_2 = c^9 t_5 c^{-9}, \tag{5.136b}$$

$$t_3 = c^8 t_1 c^{-8}, \tag{5.136c}$$

$$t_4 = c^8 t_2 c^{-8}, \tag{5.136d}$$

$$t_5 = c^8 t_3 c^{-8}. \tag{5.136e}$$

One of these equations is in fact superfluous: if we substitute (5.136b)–(5.136e) in (5.136a), then we obtain $t_1 = c^{42} t_1 c^{-42}$ which is automatically satisfied since $c^{42} = \varepsilon$.

Since $(w_0; w_1, \dots, w_m) \in NC^m(G_{34})$, we must have $t_1 t_2 t_3 t_4 t_5 \le_T c$. Combining this with (5.136), we infer that

$$t_1 (c^{25} t_1 c^{-25})(c^8 t_1 c^{-8})(c^{33} t_1 c^{-33})(c^{16} t_1 c^{-16}) \le_T c. \tag{5.137}$$

Using that $c^7 t_1 c^{-7} = t_1$, due to Lemma 6 with $d = 6$, we see that this equation is equivalent with (5.128). Therefore, we are facing exactly the same enumeration problem here as for $p = 7m/5$, and, consequently, the number of solutions to (5.137) is the same, namely $\frac{7m+5}{5}$, as required.

Next we consider the case in (5.125c). By Lemma 3, we are free to choose $p = 7m/4$ if $\zeta = \zeta_{24}$, and we are free to choose $p = 21m/4$ if $\zeta = \zeta_8$. In both cases, m must be divisible by 4.

We start with the case that $p = 7m/4$. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) = (*; c^2 w_{\frac{m}{4}+1} c^{-2}, c^2 w_{\frac{m}{4}+2} c^{-2}, \dots, c^2 w_m c^{-2}, c w_1 c^{-1}, \dots, c w_{\frac{m}{4}} c^{-1}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^2 w_{\frac{m}{4}+i} c^{-2}, \quad i = 1, 2, \dots, \tfrac{3m}{4}, \tag{5.138a}$$

$$w_i = c w_{i-\frac{3m}{4}} c^{-1}, \quad i = \tfrac{3m}{4}+1, \tfrac{3m}{4}+2, \dots, m. \tag{5.138b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{4}}) = \ell_T(w_{i+\frac{2m}{4}}) = \ell_T(w_{i+\frac{3m}{4}}) = 1, \tag{5.139a}$$

and the other w_j's, $1 \le j \le m$, are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(G_{34})$, we must have

$$w_i w_{i+\frac{m}{4}} w_{i+\frac{2m}{4}} w_{i+\frac{3m}{4}} \le_T c.$$

Together with Equations (5.138)–(5.139), this implies that

$$w_i = c^7 w_i c^{-7} \quad \text{and} \quad w_i (c^5 w_i c^{-5})(c^3 w_i c^{-3})(c w_i c^{-1}) \le_T c. \tag{5.140}$$

With the help of the *GAP* package `CHEVIE` [12], one obtains seven solutions for w_i in (5.140):

$$w_i \in \big\{[1],\ [2],\ [28],\ [34],\ [61],\ [46],\ [168]\big\},$$

where we have again used the short notation of `CHEVIE` referring to the internal ordering of the roots of G_{34} in `CHEVIE`. Each of them gives rise to $m/4$ elements of $\mathrm{Fix}_{NC^m(G_{34})}(\phi^p)$ since i ranges from 1 to $m/4$.

In total, we obtain $1 + 7\frac{m}{4} = \frac{7m+4}{4}$ elements in $\operatorname{Fix}_{NC^m(G_{34})}(\phi^p)$, which agrees with the limit in (5.125c).

If $p = 21m/4$, then, from (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^6 w_{\frac{3m}{4}+1} c^{-6}, c^6 w_{\frac{3m}{4}+2} c^{-6}, \dots, c^6 w_m c^{-6}, c^5 w_1 c^{-5}, \dots, c^5 w_{\frac{3m}{4}} c^{-5}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^6 w_{\frac{3m}{4}+i} c^{-6}, \quad i = 1, 2, \dots, \tfrac{m}{4}, \tag{5.141a}$$

$$w_i = c^5 w_{i-\frac{3m}{4}} c^{-5}, \quad i = \tfrac{m}{4}+1, \tfrac{m}{4}+2, \dots, m. \tag{5.141b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{4}}) = \ell_T(w_{i+\frac{2m}{4}}) = \ell_T(w_{i+\frac{3m}{4}}) = 1, \tag{5.142a}$$

and the other w_j's, $1 \le j \le m$, are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(G_{34})$, we must have

$$w_i w_{i+\frac{m}{4}} w_{i+\frac{2m}{4}} w_{i+\frac{3m}{4}} \le_T c.$$

Together with Equations (5.141)–(5.142), this implies that

$$w_i = c^{21} w_i c^{-21} \quad \text{and} \quad w_i (c^5 w_i c^{-5})(c^{10} w_i c^{-10})(c^{15} w_i c^{-15}) \le_T c. \tag{5.143}$$

Using that $c^7 t_1 c^{-7} = t_1$, due to Lemma 6 with $d = 6$, we see that this equation is equivalent with (5.140). Therefore, we are facing exactly the same enumeration problem here as for $p = 7m/4$, and, consequently, the number of solutions to (5.137) is the same, namely $\frac{7m+4}{4}$, as required.

Our next case is the case in (5.125d). By Lemma 3, we are free to choose $p = 7m/3$ if $\zeta = \zeta_{18}$, and we are free to choose $p = 14m/3$ if $\zeta = \zeta_9$. In both cases, m must be divisible by 3.

We start with the case that $p = 7m/3$. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^3 w_{\frac{2m}{3}+1} c^{-3}, c^3 w_{\frac{2m}{3}+2} c^{-3}, \dots, c^3 w_m c^{-3}, c^2 w_1 c^{-2}, \dots, c^2 w_{\frac{2m}{3}} c^{-2}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^3 w_{\frac{2m}{3}+i} c^{-3}, \quad i = 1, 2, \dots, \tfrac{m}{3}, \tag{5.144a}$$

$$w_i = c^2 w_{i-\frac{m}{3}} c^{-2}, \quad i = \tfrac{m}{3}+1, \tfrac{m}{3}+2, \dots, m. \tag{5.144b}$$

There are four distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{3}}) = \ell_T(w_{i+\frac{2m}{3}}) = 2, \tag{5.145a}$$

and all other w_j's are equal to ε,

(iii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{3}}) = \ell_T(w_{i+\frac{2m}{3}}) = 1, \tag{5.145b}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(iv) there are i_1 and i_2 with $1 \le i_1 < i_2 \le \frac{m}{2}$ such that

$$\ell_T(w_{i_1}) = \ell_T(w_{i_2}) = \ell_T(w_{i_1+\frac{m}{3}}) = \ell_T(w_{i_2+\frac{m}{3}}) = \ell_T(w_{i_1+\frac{2m}{3}}) = \ell_T(w_{i_2+\frac{2m}{3}}) = 1, \tag{5.145c}$$

and all other w_j's are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(G_{34})$, we must have $w_i w_{i+\frac{m}{3}} w_{i+\frac{2m}{3}} \le_T c$, respectively $w_{i_1} w_{i_2} w_{i_1+\frac{m}{3}} w_{i_2+\frac{m}{3}} w_{i_1+\frac{2m}{3}} w_{i_2+\frac{2m}{3}} = c$. Together with Equations (5.144)–(5.145), this implies that

$$w_i = c^7 w_i c^{-7} \quad \text{and} \quad w_i(c^2 w_i c^{-2})(c^4 w_i c^{-4}) = c, \tag{5.146}$$

respectively that

$$w_i = c^7 w_i c^{-7}, \quad w_i(c^2 w_i c^{-2})(c^4 w_i c^{-4}) \le_T c, \quad \text{and} \quad \ell_T(w_i) = 1, \tag{5.147}$$

respectively that

$$\begin{aligned} w_{i_1} = c^7 w_{i_1} c^{-7}, \quad & w_{i_2} = c^7 w_{i_2} c^{-7}, \\ & \text{and} \quad w_{i_1} w_{i_2} (c^2 w_{i_1} c^{-2})(c^2 w_{i_2} c^{-2})(c^4 w_{i_1} c^{-4})(c^4 w_{i_2} c^{-4}) = c. \end{aligned} \tag{5.148}$$

With the help of the *GAP* package `CHEVIE` [12], one obtains 21 solutions for w_i in (5.147):

$$w_i \in \big\{ [4] \ \ [5] \ \ [6] \ \ [11] \ \ [44] \ \ [63] \ \ [74] \big\},$$

where we have again used the short notation of `CHEVIE` referring to the internal ordering of the roots of G_{34} in `CHEVIE`. Each of them gives rise to $m/3$ elements of $\mathrm{Fix}_{NC^m(G_{34})}(\phi^p)$ since i ranges from 1 to $m/3$.

There are no solutions w_i in (5.146) and for (w_{i_1}, w_{i_2}) in (5.148).

In total, we obtain $1 + 7\frac{m}{3} = \frac{7m+3}{3}$ elements in $\mathrm{Fix}_{NC^m(G_{34})}(\phi^p)$, which agrees with the limit in (5.125d).

In the case that $p = 14m/3$, we infer from (4.1) that

$$\begin{aligned} \phi^p&\big((w_0; w_1, \dots, w_m)\big) \\ &= (*; c^5 w_{\frac{m}{3}+1} c^{-5}, c^5 w_{\frac{m}{3}+2} c^{-5}, \dots, c^5 w_m c^{-5}, c^4 w_1 c^{-4}, \dots, c^4 w_{\frac{m}{3}} c^{-4}\big). \end{aligned}$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^5 w_{\frac{m}{3}+i} c^{-5}, \quad i = 1, 2, \dots, \tfrac{2m}{3}, \tag{5.149a}$$

$$w_i = c^4 w_{i-\frac{2m}{3}} c^{-4}, \quad i = \tfrac{2m}{3}+1, \tfrac{2m}{3}+2, \dots, m. \tag{5.149b}$$

There are four distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{3}}) = \ell_T(w_{i+\frac{2m}{3}}) = 2, \tag{5.150a}$$

and all other w_j's are equal to ε,

(iii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{3}}) = \ell_T(w_{i+\frac{2m}{3}}) = 1, \tag{5.150b}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(iv) there are i_1 and i_2 with $1 \le i_1 < i_2 \le \frac{m}{2}$ such that

$$\ell_T(w_{i_1}) = \ell_T(w_{i_2}) = \ell_T(w_{i_1+\frac{m}{3}}) = \ell_T(w_{i_2+\frac{m}{3}}) = \ell_T(w_{i_1+\frac{2m}{3}}) = \ell_T(w_{i_2+\frac{2m}{3}}) = 1, \tag{5.150c}$$

and all other w_j's are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(G_{34})$, we must have $w_i w_{i+\frac{m}{3}} w_{i+\frac{2m}{3}} \le_T c$, respectively $w_{i_1} w_{i_2} w_{i_1+\frac{m}{3}} w_{i_2+\frac{m}{3}} w_{i_1+\frac{2m}{3}} w_{i_2+\frac{2m}{3}} = c$. Together with Equations (5.149)–(5.150), this implies that

$$w_i = c^{14} w_i c^{-14} \quad \text{and} \quad w_i (c^9 w_i c^{-9})(c^4 w_i c^{-4}) = c, \tag{5.151}$$

respectively that

$$w_i = c^{14} w_i c^{-14}, \quad w_i (c^9 w_i c^{-9})(c^4 w_i c^{-4}) \le_T c, \quad \text{and} \quad \ell_T(w_i) = 1, \tag{5.152}$$

respectively that

$$\begin{aligned} w_{i_1} = c^{14} w_{i_1} c^{-14}, \quad w_{i_2} = c^{14} w_{i_2} c^{-14},& \\ \text{and} \quad w_{i_1} w_{i_2} (c^9 w_{i_1} c^{-9})(c^9 w_{i_2} c^{-9})(c^4 w_{i_1} c^{-4})(c^4 w_{i_2} c^{-4}) = c.& \end{aligned} \tag{5.153}$$

Using that $c^7 w c^{-7} = w$ for all $w \in NC(G_{34}$, due to Lemma 6 with $d = 6$, we see that this equation is equivalent with (5.146). Therefore, we are facing exactly the same enumeration problem here as for $p = 7m/3$, and, consequently, the number of solutions to (5.151) is the same, namely $\frac{7m+3}{3}$, as required.

Next we consider the case in (5.125e). By Lemma 3, we are free to choose $p = 7m/2$ if $\zeta = \zeta_{12}$, and we are free to choose $p = 21m/2$ if $\zeta = \zeta_4$. In both cases, m must be divisible by 2.

We begin with the case that $p = 7m/2$. From (4.1), we infer

$$\begin{aligned} &\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ &\quad = (*; c^4 w_{\frac{m}{2}+1} c^{-4}, c^4 w_{\frac{m}{2}+2} c^{-4}, \dots, c^4 w_m c^{-4}, c^3 w_1 c^{-3}, \dots, c^3 w_{\frac{m}{2}} c^{-3}\big). \end{aligned} \tag{5.154}$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^4 w_{\frac{m}{2}+i} c^{-4}, \quad i = 1, 2, \dots, \tfrac{m}{2}, \tag{5.155a}$$

$$w_i = c^3 w_{i-\frac{m}{2}} c^{-3}, \quad i = \tfrac{m}{2}+1, \tfrac{m}{2}+2, \dots, m. \tag{5.155b}$$

There are several distinct possibilities for choosing the w_i's, $1 \le i \le m$, which we summarise as follows:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$1 \le \ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) \le 3, \tag{5.156a}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(iii) there are i_1 and i_2 with $1 \le i_1 < i_2 \le \frac{m}{2}$ such that

$$\ell_1 = \ell_T(w_{i_1}) = \ell_T(w_{i_1+\frac{m}{2}}) \ge 1, \quad \ell_2 = \ell_T(w_{i_2}) == \ell_T(w_{i_2+\frac{m}{2}}) \ge 1, \quad \ell_1 + \ell_2 \le 3, \tag{5.156b}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(iv) there are i_1, i_2, i_3 with $1 \le i_1 < i_2 < i_3 \le \frac{m}{2}$ such that

$$\ell_T(w_{i_1}) = \ell_T(w_{i_2}) = \ell_T(w_{i_3}) = \ell_T(w_{i_1+\frac{m}{2}}) = \ell_T(w_{i_2+\frac{m}{2}}) = \ell_T(w_{i_3+\frac{m}{2}}) = 1, \tag{5.156c}$$

and all other w_j's are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(G_{34})$, we must have $w_i w_{i+\frac{m}{2}} \le_T c$, respectively $w_{i_1} w_{i_2} w_{i_1+\frac{m}{2}} w_{i_2+\frac{m}{2}} \le_T c$, respectively

$$w_{i_1} w_{i_2} w_{i_3} w_{i_1+\frac{m}{2}} w_{i_2+\frac{m}{2}} w_{i_3+\frac{m}{2}} = c$$

. Together with Equations (5.155)–(5.156), this implies that

$$w_i = c^7 w_i c^{-7} \quad \text{and} \quad w_i(c^4 w_i c^{-4}) \le_T c, \tag{5.157}$$

respectively that

$$w_{i_1} = c^7 w_{i_1} c^{-7}, \quad w_{i_2} = c^7 w_{i_2} c^{-7}, \quad \text{and} \quad w_{i_1} w_{i_2} (c^4 w_{i_1} c^{-4})(c^4 w_{i_2} c^{-4}) \le_T c, \tag{5.158}$$

respectively that

$$\begin{aligned} w_{i_1} = c^7 w_{i_1} c^{-7}, \quad w_{i_2} = c^7 w_{i_2} c^{-7}, \quad w_{i_3} = c^7 w_{i_3} c^{-7},& \\ \text{and} \quad w_{i_1} w_{i_2} w_{i_3} (c^4 w_{i_1} c^{-4})(c^4 w_{i_2} c^{-4})(c^4 w_{i_3} c^{-4}) = c.& \end{aligned} \tag{5.159}$$

With the help of the *GAP* package `CHEVIE` [12], one obtains 14 solutions for w_i in (5.157) with $\ell_T(w_i) = 1$:

$$w_i \in \big\{[14],\ [35],\ [1],\ [2],\ [10],\ [24],\ [28],\ [34],\ [39],\ [56],\ [61],\ [46],\ [168],\ [105]\big\},$$

where we have again used the short notation of `CHEVIE` referring to the internal ordering of the roots of G_{34} in `CHEVIE`, one obtains 21 solutions for w_i in (5.157) with $\ell_T(w_i) = 2$:

$$\begin{aligned} w_i \in \big\{&[1,14],\ [1,35],\ [1,24],\ [1,34],\ [14,61],\ [2,39],\ [10,34],\ [2,56],\ [2,35],\ [14,46],\ [35,168],\\ &[34,56],\ [2,61],\ [28,56],\ [10,28],\ [10,61],\ [34,105],\ [28,105],\ [24,61],\ [39,168],\ [24,46]\big\}, \end{aligned}$$

each of them giving rise to $m/2$ elements of $\mathrm{Fix}_{NC^m(G_{34})}(\phi^p)$ since i ranges from 1 to $m/2$, and one obtains 49 pairs (w_{i_1}, w_{i_2}) satisfying (5.158):

$$\begin{aligned} (w_{i_1}, w_{i_2}) \in \big\{&([14],[1]),\ ([14],[61]),\ ([14],[46]),\ ([35],[1]),\ ([35],[46]),\ ([35],[168]),\ ([1],[14]),\ ([1],[35]),\ ([1],[24]),\\ &([1],[34]),\ ([2],[35]),\ ([2],[39]),\ ([2],[56]),\ ([2],[61]),\ ([10],[28]),\ ([10],[34]),\ ([10],[61]),\ ([24],[28]),\ ([24],[61]),\\ &([24],[46]),\ ([28],[1]),\ ([28],[10]),\ ([28],[56]),\ ([28],[105]),\ ([34],[39]),\ ([34],[56]),\ ([34],[46]),\ ([34],[105]),\\ &([39],[1]),\ ([39],[2]),\ ([39],[168]),\ ([56],[2]),\ ([56],[34]),\ ([56],[168]),\ ([61],[10]),\ ([61],[24]),\ ([61],[168]),\\ &([61],[105]),\ ([46],[14]),\ ([46],[2]),\ ([46],[10]),\ ([46],[24]),\ ([168],[14]),\ ([168],[35]),\ ([168],[28]),\\ &([168],[39]),\ ([105],[2]),\ ([105],[28]),\ ([105],[34])\big\}, \end{aligned}$$

each of them giving rise to $\binom{m/2}{2}$ elements of $\mathrm{Fix}_{NC^m(G_{34})}(\phi^p)$ since $1 \le i_1 < i_2 \le m$.

There are no solutions for w_i with $\ell_T(w_i) = 3$ in (5.157), and hence no solutions for (w_{i_1}, w_{i_2}) with $\ell_T(w_{i_1}) + \ell_T(w_{i_2}) = 3$ in (5.158), and no solutions for $(w_{i_1}, w_{i_2}, w_{i_3})$ in (5.159).

In total, we obtain $1 + (14+21)\frac{m}{2} + 49\binom{m/2}{2} = \frac{(7m+2)(7m+4)}{8}$ elements in $\mathrm{Fix}_{NC^m(G_{34})}(\phi^p)$, which agrees with the limit in (5.125e).

If $p = 21m/2$, from (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big)\\ = \big(*; c^{11}w_{\frac{m}{2}+1}c^{-11}, c^{11}w_{\frac{m}{2}+2}c^{-11}, \dots, c^{11}w_m c^{-11}, c^{10}w_1c^{-10}, \dots, c^{10}w_{\frac{m}{2}}c^{-10}\big).$$

Using that $c^7wc^{-7} = w$ for all $w \in NC(G_{34}$, due to Lemma 6 with $d = 6$, we see that this action is identical with the one in (5.154). Therefore, we are facing exactly the same enumeration problem here as for $p = 7m/2$, and, consequently, the number of elements in $\operatorname{Fix}_{NC^m(G_{34})}(\phi^p)$ is the same, namely $\frac{(7m+2)(7m+4)}{8}$, as required.

Finally, we turn to (5.125g). By Remark 3, the only choices for h_2 and m_2 to be considered are $h_2 = 1$ and $m_2 = 5$, $h_2 = 2$ and $m_2 = 5$, $h_2 = 2$ and $m_2 = 4$, $h_2 = m_2 = 2$, $h_2 = 3$ and $m_2 = 5$, $h_2 = m_2 = 3$, $h_2 = 6$ and $m_2 = 6$, $h_2 = 6$ and $m_2 = 5$, $h_2 = 6$ and $m_2 = 4$, $h_2 = 6$ and $m_2 = 3$, respectively $h_2 = 6$ and $m_2 = 2$, . These correspond to the choices $p = 42m/5$, $p = 21m/5$, $p = 21m/4$, $p = 21m/2$, $p = 14m/3$, $p = 14m/5$, $p = 7m/6$, $p = 7m/5$, $p = 7m/4$, $p = 7m/3$, respectively $p = 7m/2$, out of which only $p = 7m/6$ has not yet been discussed and belongs to the current case. The corresponding action of ϕ^p is given by

$$\phi^p\big((w_0; w_1, \dots, w_m)\big)\\ = \big(*; c^2w_{\frac{5m}{6}+1}c^{-2}, c^2w_{\frac{5m}{6}+2}c^{-2}, \dots, c^2w_mc^{-2}, cw_1c^{-1}, \dots, cw_{\frac{5m}{6}}c^{-1}\big),$$

so that we have to solve

$$t_1(ct_1c^{-1})(c^2t_1c^{-2})(c^3t_1c^{-3})(c^4t_1c^{-4})(c^5t_1c^{-5}) = c.$$

A computation with the help of the *GAP* package `CHEVIE` [12] finds no solution. Hence, the left-hand side of (3.3) is equal to 1, as required.

Case $G_{35} = E_6$. The degrees are $2, 5, 6, 8, 9, 12$, and hence we have

$$\operatorname{Cat}^m(E_6; q) = \frac{[12m+12]_q\,[12m+9]_q\,[12m+8]_q\,[12m+6]_q\,[12m+5]_q\,[12m+2]_q}{[12]_q\,[9]_q\,[8]_q\,[6]_q\,[5]_q\,[2]_q}.$$

Let ζ be a $12m$-th root of unity. The following cases on the right-hand side of (3.3) do occur:

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_6;q)=m+1,\quad \text{if } \zeta=\zeta_{12}, \tag{5.160a}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_6;q)=\tfrac{4m+3}{3},\quad \text{if } \zeta=\zeta_{9},\ 3\mid m, \tag{5.160b}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_6;q)=\tfrac{3m+2}{2},\quad \text{if } \zeta=\zeta_{8},\ 2\mid m, \tag{5.160c}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_6;q)=(m+1)(2m+1),\quad \text{if } \zeta=\zeta_{6}, \tag{5.160d}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_6;q)=\tfrac{12m+5}{5},\quad \text{if } \zeta=\zeta_{5},\ 5\mid m, \tag{5.160e}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_6;q)=\tfrac{(m+1)(3m+2)}{2},\quad \text{if } \zeta=\zeta_{4}, \tag{5.160f}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_6;q)=\tfrac{(m+1)(4m+3)(2m+1)}{3},\quad \text{if } \zeta=\zeta_{3}, \tag{5.160g}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_6;q)=\tfrac{(m+1)(3m+2)(2m+1)(6m+1)}{2},\quad \text{if } \zeta=-1, \tag{5.160h}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_6;q)=\operatorname{Cat}^m(E_6),\quad \text{if } \zeta=1, \tag{5.160i}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_6;q)=1,\quad \text{otherwise.} \tag{5.160j}$$

We must now prove that the left-hand side of (3.3) in each case agrees with the values exhibited in (5.160). The only cases not covered by Lemmas 4 and 5 are the ones in (5.160b), (5.160c), (5.160e), and (5.160j).

We begin with the case in (5.160b). By Lemma 3, we are free to choose $p=4m/3$. In particular, m must be divisible by 3. From (4.1), we infer

$$\phi^p\big((w_0;w_1,\dots,w_m)\big)\\ =\big(*;c^2w_{\frac{2m}{3}+1}c^{-2},c^2w_{\frac{2m}{3}+2}c^{-2},\dots,c^2w_mc^{-2},cw_1c^{-1},\dots,cw_{\frac{2m}{3}}c^{-1}\big).$$

Supposing that $(w_0;w_1,\dots,w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i=c^2w_{\frac{2m}{3}+i}c^{-2},\quad i=1,2,\dots,\tfrac{m}{3}, \tag{5.161a}$$

$$w_i=cw_{i-\frac{m}{3}}c^{-1},\quad i=\tfrac{m}{3}+1,\tfrac{m}{3}+2,\dots,m. \tag{5.161b}$$

There are four distinct possibilities for choosing the w_i's, $1\le i\le m$:

(i) all the w_i's are equal to ε (and $w_0=c$),

(ii) there is an i with $1\le i\le\frac{m}{3}$ such that

$$\ell_T(w_i)=\ell_T(w_{i+\frac{m}{3}})=\ell_T(w_{i+\frac{2m}{3}})=2, \tag{5.162a}$$

and all other w_j's are equal to ε,

(iii) there is an i with $1\le i\le\frac{m}{3}$ such that

$$\ell_T(w_i)=\ell_T(w_{i+\frac{m}{3}})=\ell_T(w_{i+\frac{2m}{3}})=1, \tag{5.162b}$$

and the other w_j's, $1\le j\le m$, are equal to ε,

(iv) there are i_1 and i_2 with $1\le i_1<i_2\le\frac{m}{3}$ such that

$$\ell_T(w_{i_1})=\ell_T(w_{i_2})=\ell_T(w_{i_1+\frac{m}{3}})=\ell_T(w_{i_2+\frac{m}{3}})=\ell_T(w_{i_1+\frac{2m}{3}})=\ell_T(w_{i_2+\frac{2m}{3}})=1, \tag{5.162c}$$

and all other w_j's are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(E_6)$, we must have $w_i w_{i+\frac{m}{3}} w_{i+\frac{2m}{3}} \le_T c$, respectively $w_{i_1} w_{i_2} w_{i_1+\frac{m}{3}} w_{i_2+\frac{m}{3}} w_{i_1+\frac{2m}{3}} w_{i_2+\frac{2m}{3}} = c$. Together with Equations (5.161)–(5.162), this implies that

$$w_i = c^4 w_i c^{-4} \quad \text{and} \quad w_i(c w_i c^{-1})(c^2 w_i c^{-2}) = c, \tag{5.163}$$

respectively that

$$w_i = c^4 w_i c^{-4}, \quad w_i(c w_i c^{-1})(c^2 w_i c^{-2}) \le_T c, \quad \text{and} \quad \ell_T(w_i) = 1, \tag{5.164}$$

respectively that

$$w_{i_1} = c^4 w_{i_1} c^{-4}, \quad w_{i_1}(c w_{i_1} c^{-1})(c^2 w_{i_1} c^{-2}) \le_T c, \quad \text{and} \quad \ell_T(w_{i_1}) = 1. \tag{5.165}$$

With the help of Stembridge's *Maple* package `coxeter` [29], one obtains four solutions for w_i in (5.163):

$$w_i \in \big\{[1,2,3,4,5,6,5,4,2,3],\ [3,4],\ [2,4,2,5],\ [1,3,4,5,4,3,1,6]\big\},$$

where we have again used the short notation of `coxeter`, $\{s_1, s_2, s_3, s_4, s_5, s_6\}$ being a simple system of generators of E_6, corresponding to the Dynkin diagram displayed in Figure 4. Each of the above solutions for w_i gives rise to $m/3$ elements of $\operatorname{Fix}_{NC^m(E_6)}(\phi^p)$ since i ranges from 1 to $m/3$.



FIGURE 4. The Dynkin diagram for E_6

There are no solutions for w_i in (5.164) and for w_{i_1} in (5.165).

In total, we obtain $1 + 4\frac{m}{3} = \frac{4m+3}{3}$ elements in $\operatorname{Fix}_{NC^m(E_6)}(\phi^p)$, which agrees with the limit in (5.160b).

Next we discuss the case in (5.160c). By Lemma 3, we are free to choose $p = 3m/2$. In particular, m must be divisible by 2. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) = (*; c^2 w_{\frac{m}{2}+1} c^{-2}, c^2 w_{\frac{m}{2}+2} c^{-2}, \dots, c^2 w_m c^{-2}, c w_1 c^{-1}, \dots, c w_{\frac{m}{2}} c^{-1}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^2 w_{\frac{m}{2}+i} c^{-2}, \quad i = 1, 2, \dots, \tfrac{m}{2}, \tag{5.166a}$$

$$w_i = c w_{i-\frac{m}{2}} c^{-1}, \quad i = \tfrac{m}{2}+1, \tfrac{m}{2}+2, \dots, m. \tag{5.166b}$$

There are several distinct possibilities for choosing the w_i's, $1 \le i \le m$, which we summarise as follows:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 3, \tag{5.167a}$$

and all other w_j's are equal to ε,

(iii) there is a j with $1 \le j \le \frac{m}{2}$ such that

$$1 \le \ell_T(w_j) = \ell_T(w_{j+\frac{m}{2}}) \le 2. \tag{5.167b}$$

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(E_6)$, we must have $w_i w_{i+\frac{m}{2}} = c$, respectively $w_j w_{j+\frac{m}{2}} \le_T c$. Together with Equations (5.166)–(5.167), this implies that

$$w_i = c^3 w_i c^{-3} \quad \text{and} \quad w_i(c w_i c^{-1}) = c, \tag{5.168}$$

respectively that

$$w_j = c^3 w_j c^{-3}, \quad w_j(c w_j c^{-1}) \le_T c, \quad \text{and} \quad 1 \le \ell_T(w_j) \le 2. \tag{5.169}$$

With the help of Stembridge's *Maple* package `coxeter` [29], one obtains three solutions for w_i in (5.168):

$$w_i \in \big\{[1,3,4,3,5],\ [1,2,3,4,3,5,6,5,4,3,1],\ [2,3,4,5,4,2,6]\big\},$$

where we used again `coxeter`'s short notation, $\{s_1, s_2, s_3, s_4, s_5, s_6\}$ being a simple system of generators of E_6, corresponding to the Dynkin diagram displayed in Figure 4. Each of these solutions for w_i gives rise to $m/2$ elements of $\operatorname{Fix}_{NC^m(E_6)}(\phi^p)$ since i ranges from 1 to $m/2$.

There are no solutions for w_j in (5.169).

In total, we obtain $1 + 3\frac{m}{2} = \frac{3m+2}{2}$ elements in $\operatorname{Fix}_{NC^m(E_6)}(\phi^p)$, which agrees with the limit in (5.160c).

Finally we discuss the case in (5.160e). By Lemma 3, we are free to choose $p = 12m/5$. In particular, m must be divisible by 5. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^3 w_{\frac{3m}{5}+1} c^{-3}, c^3 w_{\frac{3m}{5}+2} c^{-3}, \dots, c^3 w_m c^{-3}, c^2 w_1 c^{-2}, \dots, c^2 w_{\frac{3m}{5}} c^{-2}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^3 w_{\frac{3m}{5}+i} c^{-3}, \quad i = 1, 2, \dots, \tfrac{2m}{5}, \tag{5.170a}$$

$$w_i = c^2 w_{i-\frac{2m}{5}} c^{-2}, \quad i = \tfrac{2m}{5}+1, \tfrac{2m}{5}+2, \dots, m. \tag{5.170b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{5}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{5}}) = \ell_T(w_{i+\frac{2m}{5}}) = \ell_T(w_{i+\frac{3m}{5}}) = \ell_T(w_{i+\frac{4m}{5}}) = 1, \tag{5.171}$$

and the other w_j's, $1 \le j \le m$, are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(E_6)$, we must have

$$w_i w_{i+\frac{m}{5}} w_{i+\frac{2m}{5}} w_{i+\frac{3m}{5}} w_{i+\frac{4m}{5}} \le_T c.$$

Together with Equations (5.170)–(5.171), this implies that

$$w_i = c^{12} w_i c^{-12} \quad \text{and} \quad w_i(c^7 w_i c^{-7})(c^2 w_i c^{-2})(c^9 w_i c^{-9})(c^4 w_i c^{-4}) \le_T c. \tag{5.172}$$

Here, the first equation is automatically satisfied since $c^{12} = \varepsilon$.

With the help of Stembridge's *Maple* package `coxeter` [29], one obtains 12 solutions for w_i in (5.172):

$$w_i \in \big\{[1], [3], [4], [5], [6], [2,4,2], [3,4,3], [2,4,5,4,2], [1,3,4,5,4,3,1], \\ [1,3,4,5,6,5,4,3,1], [2,3,4,5,6,5,4,2,3], [1,2,3,4,5,6,5,4,2,3,1]\big\},$$

where $\{s_1, s_2, s_3, s_4, s_5, s_6\}$ is a simple system of generators of E_6, corresponding to the Dynkin diagram displayed in Figure 4, and each of them gives rise to $m/5$ elements of $\mathrm{Fix}_{NC^m(E_6)}(\phi^p)$ since i ranges from 1 to $m/5$.

In total, we obtain $1 + 12\frac{m}{5} = \frac{12m+5}{5}$ elements in $\mathrm{Fix}_{NC^m(E_6)}(\phi^p)$, which agrees with the limit in (5.160e).

Finally, we turn to (5.160j). By Remark 3, the only choices for h_2 and m_2 to be considered are $h_2 = 1$ and $m_2 = 5$ and $h_2 = 2$ and $m_2 = 5$. These correspond to the choices $p = 12m/5$, respectively $p = 6m/5$, out which only $p = 6m/5$ has not yet been discussed and belongs to the current case. The corresponding action of ϕ^p is given by

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^2 w_{\frac{4m}{5}+1} c^{-2}, c^2 w_{\frac{4m}{5}+2} c^{-2}, \dots, c^2 w_m c^{-2}, c w_1 c^{-1}, \dots, c w_{\frac{4m}{5}} c^{-1}\big),$$

so that we have to solve

$$t_1 (c t_1 c^{-1})(c^2 t_1 c^{-2})(c^3 t_1 c^{-3})(c^4 t_1 c^{-4}) \le_T c$$

for t_1 with $\ell_T(t_1) = 1$. A computation with the help of Stembridge's *Maple* package `coxeter` [29] finds no solution. Hence, the left-hand side of (3.3) is equal to 1, as required.

CASE $G_{36} = E_7$. The degrees are $2, 6, 8, 10, 12, 14, 18$, and hence we have

$$\mathrm{Cat}^m(E_7; q) = \frac{[18m+18]_q\,[18m+14]_q\,[18m+12]_q}{[18]_q\,[14]_q\,[12]_q} \\ \times \frac{[18m+10]_q\,[18m+8]_q\,[18m+6]_q\,[18m+2]_q}{[10]_q\,[8]_q\,[6]_q\,[2]_q}.$$

Let ζ be a $18m$-th root of unity. The following cases on the right-hand side of (3.3) do occur:

$$\lim_{q\to\zeta} \mathrm{Cat}^m(E_7; q) = m + 1, \quad \text{if } \zeta = \zeta_{18}, \zeta_9, \tag{5.173a}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(E_7; q) = \tfrac{9m+7}{7}, \quad \text{if } \zeta = \zeta_{14}, \zeta_7,\ 7 \mid m, \tag{5.173b}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(E_7; q) = \tfrac{3m+2}{2}, \quad \text{if } \zeta = \zeta_{12},\ 2 \mid m, \tag{5.173c}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(E_7; q) = \tfrac{9m+5}{5}, \quad \text{if } \zeta = \zeta_{10}, \zeta_5,\ 5 \mid m, \tag{5.173d}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(E_7; q) = \tfrac{9m+4}{4}, \quad \text{if } \zeta = \zeta_8,\ 4 \mid m, \tag{5.173e}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(E_7; q) = \frac{(m+1)(3m+2)(3m+1)}{2}, \quad \text{if } \zeta = \zeta_6, \zeta_3, \tag{5.173f}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(E_7; q) = \frac{(3m+2)(9m+4)}{8}, \quad \text{if } \zeta = \zeta_4,\ 2 \mid m, \tag{5.173g}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(E_7; q) = \mathrm{Cat}^m(E_7), \quad \text{if } \zeta = -1 \text{ or } \zeta = 1, \tag{5.173h}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(E_7; q) = 1, \quad \text{otherwise.} \tag{5.173i}$$

We must now prove that the left-hand side of (3.3) in each case agrees with the values exhibited in (5.173). The only cases not covered by Lemmas 4 and 5 are the ones in (5.173b), (5.173c), (5.173d), (5.173e), (5.173g), and (5.173i).

We begin with the case in (5.173b). By Lemma 3, we are free to choose $p = 9m/7$ if $\zeta = \zeta_{14}$, respectively $p = 18m/7$ if $\zeta = \zeta_7$. In both cases, m must be divisible by 7.

We start with the case that $p = 9m/7$. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^2 w_{\frac{5m}{7}+1} c^{-2}, c^2 w_{\frac{5m}{7}+2} c^{-2}, \dots, c^2 w_m c^{-2}, c w_1 c^{-1}, \dots, c w_{\frac{5m}{7}} c^{-1}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^2 w_{\frac{5m}{7}+i} c^{-2}, \quad i = 1, 2, \dots, \tfrac{2m}{7}, \tag{5.174a}$$

$$w_i = c w_{i-\frac{2m}{7}} c^{-1}, \quad i = \tfrac{2m}{7}+1, \tfrac{2m}{7}+2, \dots, m. \tag{5.174b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{7}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{7}}) = \ell_T(w_{i+\frac{2m}{7}}) = \ell_T(w_{i+\frac{3m}{7}}) \\ = \ell_T(w_{i+\frac{4m}{7}}) = \ell_T(w_{i+\frac{5m}{7}}) = \ell_T(w_{i+\frac{6m}{7}}) = 1, \tag{5.175}$$

and the other w_j's, $1 \le j \le m$, are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(E_7)$, we must have

$$w_i w_{i+\frac{m}{7}} w_{i+\frac{2m}{7}} w_{i+\frac{3m}{7}} w_{i+\frac{4m}{7}} w_{i+\frac{5m}{7}} w_{i+\frac{6m}{7}} = c.$$

Together with Equations (5.174)–(5.175), this implies that

$$w_i = c^9 w_i c^{-9} \quad \text{and} \quad w_i(c^5 w_i c^{-5})(c w_i c^{-1})(c^6 w_i c^{-6})(c^2 w_i c^{-2})(c^7 w_i c^{-7})(c^3 w_i c^{-3}) = c. \tag{5.176}$$

Here, the first equation is automatically satisfied due to Lemma 6 with $d = 2$.

With the help of Stembridge's *Maple* package `coxeter` [29], one obtains 9 solutions for w_i in (5.176):

$$w_i \in \big\{[4],\ [5],\ [6],\ [7],\ [3,4,3],\ [2,4,5,4,2],\ [1,3,4,5,6,5,4,3,1], \\ [2,3,4,5,6,7,6,5,4,2,3],\ [1,2,3,4,5,6,7,6,5,4,2,3,1]\big\}, \tag{5.177}$$

where we have again used the short notation of `coxeter`, $\{s_1, s_2, s_3, s_4, s_5, s_6, s_7\}$ being a simple system of generators of E_7, corresponding to the Dynkin diagram displayed in Figure 5. Each of the above solutions for w_i gives rise to $m/7$ elements of $\mathrm{Fix}_{NC^m(E_7)}(\phi^p)$ since i ranges from 1 to $m/7$.



FIGURE 5. The Dynkin diagram for E_7

In total, we obtain $1 + 9\frac{m}{7} = \frac{9m+7}{7}$ elements in $\mathrm{Fix}_{NC^m(E_7)}(\phi^p)$, which agrees with the limit in (5.173b).

In the case that $p = 18m/7$, we infer from (4.1) that

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = (*; c^3 w_{\frac{3m}{7}+1} c^{-3}, c^3 w_{\frac{3m}{7}+2} c^{-3}, \dots, c^3 w_m c^{-3}, c^2 w_1 c^{-2}, \dots, c^2 w_{\frac{3m}{7}} c^{-2}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^3 w_{\frac{3m}{7}+i} c^{-3}, \quad i = 1, 2, \dots, \tfrac{4m}{7}, \tag{5.178a}$$

$$w_i = c^2 w_{i-\frac{4m}{7}} c^{-2}, \quad i = \tfrac{4m}{7}+1, \tfrac{4m}{7}+2, \dots, m. \tag{5.178b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),
(ii) there is an i with $1 \le i \le \frac{m}{7}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{7}}) = \ell_T(w_{i+\frac{2m}{7}}) = \ell_T(w_{i+\frac{3m}{7}}) \\ = \ell_T(w_{i+\frac{4m}{7}}) = \ell_T(w_{i+\frac{5m}{7}}) = \ell_T(w_{i+\frac{6m}{7}}) = 1, \tag{5.179}$$

and the other w_j's, $1 \le j \le m$, are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(E_7)$, we must have

$$w_i w_{i+\frac{m}{7}} w_{i+\frac{2m}{7}} w_{i+\frac{3m}{7}} w_{i+\frac{4m}{7}} w_{i+\frac{5m}{7}} w_{i+\frac{6m}{7}} = c.$$

Together with Equations (5.178)–(5.179), this implies that

$$w_i = c^{18} w_i c^{-18} \\ \text{and} \quad w_i (c^5 w_i c^{-5})(c^{10} w_i c^{-10})(c^{15} w_i c^{-15})(c^2 w_i c^{-2})(c^7 w_i c^{-7})(c^{12} w_i c^{-12}) = c. \tag{5.180}$$

Here, the first equation is automatically satisfied since $c^{18} = \varepsilon$. Due to Lemma 6 with $d = 2$, we have $c^9 w_i c^{-9} = w_i$, hence also $c^{10} w_i c^{-10} = c w_i c^{-1}$, etc., so that (5.180) reduces to (5.176). Therefore, we are facing exactly the same enumeration problem here as for $p = 9m/7$, and, consequently, the number of solutions to (5.180) is the same, namely $\frac{9m+7}{7}$, as required.

Next we discuss the case in (5.173c). By Lemma 3, we are free to choose $p = 3m/2$. In particular, m must be divisible by 2. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) = (*; c^2 w_{\frac{m}{2}+1} c^{-2}, c^2 w_{\frac{m}{2}+2} c^{-2}, \dots, c^2 w_m c^{-2}, c w_1 c^{-1}, \dots, c w_{\frac{m}{2}} c^{-1}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^2 w_{\frac{m}{2}+i} c^{-2}, \quad i = 1, 2, \dots, \tfrac{m}{2}, \tag{5.181a}$$

$$w_i = c w_{i-\frac{m}{2}} c^{-1}, \quad i = \tfrac{m}{2}+1, \tfrac{m}{2}+2, \dots, m. \tag{5.181b}$$

There are several distinct possibilities for choosing the w_i's, $1 \le i \le m$, which we summarise as follows:

(i) all the w_i's are equal to ε (and $w_0 = c$),
(ii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$1 \le \ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) \le 3. \tag{5.182}$$

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(E_7)$, we must have $w_i w_{i+\frac{m}{2}} \le_T c$. Together with Equations (5.181)–(5.182), this implies that

$$w_i = c^3 w_i c^{-3}, \quad w_i(c w_i c^{-1}) \le_T c, \quad \text{and} \quad 1 \le \ell_T(w_i) \le 3. \tag{5.183}$$

With the help of Stembridge's *Maple* package `coxeter` [29], one obtains three solutions for w_i in (5.183) with $\ell_T(w_i) = 2$:

$$w_i \in \big\{[1,3,4,5,4,3],\ [2,3,4,5,6,5,4,2],\ [1,2,4,2,3,4,5,6,7,6,5,4,3,1]\big\},$$

where we used again `coxeter`'s short notation, $\{s_1, s_2, s_3, s_4, s_5, s_6, s_7\}$ being a simple system of generators of E_7, corresponding to the Dynkin diagram displayed in Figure 5, and none if $\ell_T(w_i) = 1$ or $\ell_T(w_i) = 3$. Each of the solutions for w_i gives rise to $m/2$ elements of $\mathrm{Fix}_{NC^m(E_7)}(\phi^p)$ since i ranges from 1 to $m/2$.

In total, we obtain $1 + 3\frac{m}{2} = \frac{3m+2}{2}$ elements in $\mathrm{Fix}_{NC^m(E_7)}(\phi^p)$, which agrees with the limit in (5.173c).

Next we consider the case in (5.173d). By Lemma 3, we are free to choose $p = 9m/5$ if $\zeta = \zeta_{10}$, respectively $p = 18m/5$ if $\zeta = \zeta_5$. In both cases, m must be divisible by 5.

We start with the case that $p = 9m/5$. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) = (*; c^2 w_{\frac{m}{5}+1} c^{-2}, c^2 w_{\frac{m}{5}+2} c^{-2}, \dots, c^2 w_m c^{-2}, c w_1 c^{-1}, \dots, c w_{\frac{m}{5}} c^{-1}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^2 w_{\frac{m}{5}+i} c^{-2}, \quad i = 1, 2, \dots, \tfrac{4m}{5}, \tag{5.184a}$$

$$w_i = c w_{i-\frac{4m}{5}} c^{-1}, \quad i = \tfrac{4m}{5}+1, \tfrac{4m}{5}+2, \dots, m. \tag{5.184b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{5}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{5}}) = \ell_T(w_{i+\frac{2m}{5}}) = \ell_T(w_{i+\frac{3m}{5}}) = \ell_T(w_{i+\frac{4m}{5}}) = 1, \tag{5.185a}$$

and the other w_j's, $1 \le j \le m$, are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(E_7)$, we must have

$$w_i w_{i+\frac{m}{5}} w_{i+\frac{2m}{5}} w_{i+\frac{3m}{5}} w_{i+\frac{4m}{5}} \le_T c.$$

Together with Equations (5.184)–(5.185), this implies that

$$w_i = c^9 w_i c^{-9} \quad \text{and} \quad w_i(c^7 w_i c^{-7})(c^5 w_i c^{-5})(c^3 w_i c^{-3})(c w_i c^{-1}) \le_T c. \tag{5.186}$$

With the help of Stembridge's *Maple* package `coxeter` [29], one obtains 9 solutions for w_i in (5.186):

$$w_i \in \big\{[4],\ [5],\ [6],\ [7],\ [3,4,3],\ [2,4,5,4,2],\ [1,3,4,5,6,5,4,3,1],$$
$$[2,3,4,5,6,7,6,5,4,2,3],\ [1,2,3,4,5,6,7,6,5,4,2,3,1]\big\},$$

where $\{s_1, s_2, s_3, s_4, s_5, s_6, s_7\}$ is a simple system of generators of E_7, corresponding to the Dynkin diagram displayed in Figure 5, and each of them gives rise to $m/5$ elements of $\mathrm{Fix}_{NC^m(E_7)}(\phi^p)$ since i ranges from 1 to $m/5$.[3]

In total, we obtain $1 + 9\frac{m}{5} = \frac{9m+5}{5}$ elements in $\mathrm{Fix}_{NC^m(E_7)}(\phi^p)$, which agrees with the limit in (5.173d).

---

[3] Miraculously, these are exactly the same solutions as in the case of (5.173b). We have no explanation for this phenomenon.

In the case that $p = 18m/5$, we infer from (4.1) that

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^4 w_{\frac{2m}{5}+1} c^{-4}, c^4 w_{\frac{2m}{5}+2} c^{-4}, \dots, c^4 w_m c^{-4}, c^3 w_1 c^{-3}, \dots, c^3 w_{\frac{2m}{5}} c^{-3}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^4 w_{\frac{2m}{5}+i} c^{-4}, \quad i = 1, 2, \dots, \tfrac{3m}{5}, \tag{5.187a}$$

$$w_i = c^3 w_{i-\frac{3m}{5}} c^{-3}, \quad i = \tfrac{3m}{5}+1, \tfrac{3m}{5}+2, \dots, m. \tag{5.187b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{5}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{5}}) = \ell_T(w_{i+\frac{2m}{5}}) = \ell_T(w_{i+\frac{3m}{5}}) = \ell_T(w_{i+\frac{4m}{5}}) = 1, \tag{5.188a}$$

and the other w_j's, $1 \le j \le m$, are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(E_7)$, we must have

$$w_i w_{i+\frac{m}{5}} w_{i+\frac{2m}{5}} w_{i+\frac{3m}{5}} w_{i+\frac{4m}{5}} \le_T c.$$

Together with Equations (5.187)–(5.188), this implies that

$$w_i = c^{18} w_i c^{-18} \quad \text{and} \quad w_i (c^7 w_i c^{-7})(c^{14} w_i c^{-14})(c^3 w_i c^{-3})(c^{10} w_i c^{-10}) \le_T c. \tag{5.189}$$

Here, the first equation is automatically satisfied since $c^{18} = \varepsilon$. Due to Lemma 6 with $d = 2$, we have $c^9 w_i c^{-9} = w_i$, hence also $c^{14} w_i c^{-14} = c^5 w_i c^{-5}$, etc., so that (5.189) reduces to (5.186). Therefore, we are facing exactly the same enumeration problem here as for $p = 9m/5$, and, consequently, the number of solutions to (5.189) is the same, namely $\frac{9m+5}{5}$, as required.

Our next case is the case in (5.173e). By Lemma 3, we are free to choose $p = 9m/4$. In particular, m must be divisible by 4. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^3 w_{\frac{3m}{4}+1} c^{-3}, c^3 w_{\frac{3m}{4}+2} c^{-3}, \dots, c^3 w_m c^{-3}, c^2 w_1 c^{-2}, \dots, c^2 w_{\frac{3m}{4}} c^{-2}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^3 w_{\frac{3m}{4}+i} c^{-3}, \quad i = 1, 2, \dots, \tfrac{m}{4}, \tag{5.190a}$$

$$w_i = c^2 w_{i-\frac{m}{4}} c^{-2}, \quad i = \tfrac{m}{4}+1, \tfrac{m}{4}+2, \dots, m. \tag{5.190b}$$

There are several distinct possibilities for choosing the w_i's, $1 \le i \le m$, which we summarise as follows:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{4}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{4}}) = \ell_T(w_{i+\frac{2m}{4}}) = \ell_T(w_{i+\frac{3m}{4}}) = 1, \tag{5.191}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(E_7)$, we must have $w_i w_{i+\frac{m}{4}} w_{i+\frac{2m}{4}} w_{i+\frac{3m}{4}} \le_T c$. Together with Equations (5.190)–(5.191), this implies that

$$w_i = c^9 w_i c^{-9} \quad \text{and} \quad w_i (c^2 w_i c^{-2})(c^4 w_i c^{-4})(c^6 w_i c^{-6}) \le_T c. \tag{5.192}$$

Here, the first equation in (5.192) is automatically satisfied due to Lemma 6 with $d = 2$.

With the help of Stembridge's *Maple* package `coxeter` [29], one obtains 9 solutions for w_i in (5.192) with $\ell_T(w_i) = 1$:

$$w_i \in \big\{[1],\ [3],\ [2,4,2],\ [3,4,5,4,3],\ [1,3,4,5,4,3,1],\ [2,4,5,6,5,4,2],\ [2,3,4,5,6,5,4,2,3],\\ [1,3,4,5,6,7,6,5,4,3,1],\ [1,4,2,3,4,5,6,7,6,5,4,2,3,1,4]\big\},$$

where $\{s_1, s_2, s_3, s_4, s_5, s_6, s_7\}$ is a simple system of generators of E_7, corresponding to the Dynkin diagram displayed in Figure 5, and each of them gives rise to $m/4$ elements of $\operatorname{Fix}_{NC^m(E_7)}(\phi^p)$ since i ranges from 1 to $m/4$. Hence, we obtain $1 + 9\frac{m}{4} = \frac{9m+4}{4}$ elements in $\operatorname{Fix}_{NC^m(E_7)}(\phi^p)$, which agrees with the limit in (5.173e).

Finally we discuss the case in (5.173g). By Lemma 3, we are free to choose $p = 9m/2$. In particular, m must be divisible by 2. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big)\\ = \big(*; c^5 w_{\frac{m}{2}+1} c^{-5}, c^5 w_{\frac{m}{2}+2} c^{-5}, \dots, c^5 w_m c^{-5}, c^4 w_1 c^{-4}, \dots, c^4 w_{\frac{m}{2}} c^{-4}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^5 w_{\frac{m}{2}+i} c^{-5}, \quad i = 1, 2, \dots, \tfrac{m}{2}, \tag{5.193a}$$

$$w_i = c^4 w_{i-\frac{m}{2}} c^{-4}, \quad i = \tfrac{m}{2}+1, \tfrac{m}{2}+2, \dots, m. \tag{5.193b}$$

There are several distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 3, \tag{5.194a}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(iii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 2, \tag{5.194b}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(iv) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) = 1, \tag{5.194c}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(v) there are i_1 and i_2 with $1 \le i_1 < i_2 \le \frac{m}{2}$ such that

$$\ell_T(w_{i_1}) = \ell_T(w_{i_2}) = \ell_T(w_{i_1+\frac{m}{2}}) = \ell_T(w_{i_2+\frac{m}{2}}) = 1, \tag{5.194d}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(vi) there are i_1 and i_2 with $1 \le i_1, i_2 \le \frac{m}{2}$ such that

$$\ell_T(w_{i_1}) = \ell_T(w_{i_1+\frac{m}{2}}) = 2 \quad \text{and} \quad \ell_T(w_{i_2}) = \ell_T(w_{i_2+\frac{m}{2}}) = 1, \tag{5.194e}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(vii) there are i_1, i_2, i_3 with $1 \le i_1 < i_2 < i_3 \le \frac{m}{2}$ such that

$$\ell_T(w_{i_1}) = \ell_T(w_{i_2}) = \ell_T(w_{i_3}) = \ell_T(w_{i_1+\frac{m}{2}}) = \ell_T(w_{i_2+\frac{m}{2}}) = \ell_T(w_{i_3+\frac{m}{2}}) = 1, \tag{5.194f}$$

and the other w_j's, $1 \le j \le m$, are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(E_7)$, we must have $w_i w_{i+\frac{m}{2}} \le_T c$, respectively $w_{i_1} w_{i_2} w_{i_1+\frac{m}{2}} w_{i_2+\frac{m}{2}} \le_T c$, respectively

$$w_{i_1} w_{i_2} w_{i_3} w_{i_1+\frac{m}{2}} w_{i_2+\frac{m}{2}} w_{i_3+\frac{m}{2}} \le_T c.$$

Together with Equations (5.193)–(5.194), this implies that

$$w_i = c^9 w_i c^{-9} \quad \text{and} \quad w_i(c^4 w_i c^{-4}) \le_T c, \tag{5.195}$$

respectively that

$$w_{i_1} = c^9 w_{i_1} c^{-9}, \quad w_{i_2} = c^9 w_{i_2} c^{-9}, \quad \text{and} \quad w_{i_1} w_{i_2} (c^4 w_{i_1} c^{-4})(c^4 w_{i_2} c^{-4}) \le_T c, \tag{5.196}$$

respectively that

$$\begin{aligned} w_{i_1} = c^9 w_{i_1} c^{-9}, \quad w_{i_2} = c^9 w_{i_2} c^{-9}, \quad w_{i_3} = c^9 w_{i_3} c^{-9},& \\ \text{and} \quad w_{i_1} w_{i_2} w_{i_3} (c^4 w_{i_1} c^{-4})(c^4 w_{i_2} c^{-4})(c^4 w_{i_3} c^{-4}) \le_T c.& \end{aligned} \tag{5.197}$$

Here, the first equation in (5.195), the first two in (5.196), and the first three in (5.197), are all automatically satisfied due to Lemma 6 with $d = 2$.

With the help of Stembridge's *Maple* package `coxeter` [29], one obtains 9 solutions for w_i in (5.195) with $\ell_T(w_i) = 1$:

$$\begin{aligned} w_i \in \big\{&[1],\ [3],\ [2,4,2],\ [3,4,5,4,3],\ [1,3,4,5,4,3,1],\ [2,4,5,6,5,4,2],\ [2,3,4,5,6,5,4,2,3], \\ &[1,3,4,5,6,7,6,5,4,3,1],\ [1,4,2,3,4,5,6,7,6,5,4,2,3,1,4]\big\}, \end{aligned}$$

where $\{s_1, s_2, s_3, s_4, s_5, s_6, s_7\}$ is a simple system of generators of E_7, corresponding to the Dynkin diagram displayed in Figure 5, and each of them gives rise to $m/2$ elements of $\operatorname{Fix}_{NC^m(E_7)}(\phi^p)$ since i ranges from 1 to $m/2$. Furthermore, one obtains 12 solutions for w_i in (5.195) with $\ell_T(w_i) = 2$:

$$\begin{aligned} w_i \in \big\{&[1,2,4,2],\ [1,3,4,5,4,3],\ [1,2,4,5,6,5,4,2],\ [1,3,1,4,5,4,3,1],\ [2,3,4,5,6,5,4,2], \\ &[1,3,1,4,5,6,7,6,5,4,3,1],\ [2,3,4,3,5,6,5,4,2,3],\ [1,2,4,2,3,4,5,6,7,6,5,4,3,1], \\ &[2,3,4,2,5,4,2,6,5,4,2,3],\ [1,3,1,4,3,5,4,3,1,6,7,6,5,4,3,1], \\ &[1,3,4,2,3,4,5,4,2,6,7,6,5,4,2,3,1,4],\ [1,2,4,2,3,4,5,4,6,5,4,3,7,6,5,4,2,3,1,4]\big\}, \end{aligned}$$

each of them giving rise to $m/2$ elements of $\operatorname{Fix}_{NC^m(E_7)}(\phi^p)$ since i ranges from 1 to $m/2$, and one obtains 27 pairs (w_{i_1}, w_{i_2}) of solutions in (5.196) with $\ell_T(w_{i_1}) = \ell_T(w_{i_2}) = 1$:

$$\begin{gathered}
w_i \in \big\{([1],[2,4,2]),\ ([1],[3,4,5,4,3]),\ ([1],[2,4,5,6,5,4,2]),\ ([3],[1,3,4,5,4,3,1]),\\
([3],[2,4,5,6,5,4,2]),\ ([3],[1,3,4,5,6,7,6,5,4,3,1]),\ ([2,4,2],[1]),\\
([2,4,2],[2,3,4,5,6,5,4,2,3]),\ ([2,4,2],[1,3,4,5,6,7,6,5,4,3,1]),\\
([3,4,5,4,3],[1,3,4,5,4,3,1]),\ ([3,4,5,4,3],[2,3,4,5,6,5,4,2,3]),\\
([3,4,5,4,3],[1,3,4,5,6,7,6,5,4,3,1]),\ ([1,3,4,5,4,3,1],[1]),\\
([1,3,4,5,4,3,1],[3]),\ ([1,3,4,5,4,3,1],[1,4,2,3,4,5,6,7,6,5,4,2,3,1,4]),\\
([2,4,5,6,5,4,2],[1]),\ ([2,4,5,6,5,4,2],[2,3,4,5,6,5,4,2,3]),\\
([2,4,5,6,5,4,2],[1,4,2,3,4,5,6,7,6,5,4,2,3,1,4]),\ ([2,3,4,5,6,5,4,2,3],[3]),\\
([2,3,4,5,6,5,4,2,3],[2,4,2]),\ ([2,3,4,5,6,5,4,2,3],[3,4,5,4,3]),\\
([1,3,4,5,6,7,6,5,4,3,1],[3]),\ ([1,3,4,5,6,7,6,5,4,3,1],[3,4,5,4,3]),\\
([1,3,4,5,6,7,6,5,4,3,1],[1,4,2,3,4,5,6,7,6,5,4,2,3,1,4]),\\
([1,4,2,3,4,5,6,7,6,5,4,2,3,1,4],[2,4,2]),\\
([1,4,2,3,4,5,6,7,6,5,4,2,3,1,4],[1,3,4,5,4,3,1]),\\
([1,4,2,3,4,5,6,7,6,5,4,2,3,1,4],[2,4,5,6,5,4,2])\big\},
\end{gathered}$$

each of them giving rise to $\binom{m/2}{2}$ elements of $\operatorname{Fix}_{NC^m(E_7)}(\phi^p)$ since $1 \le i_1 < i_2 \le \frac{m}{2}$.

There are no solutions for w_i in (5.195) with $\ell_T(w_i) = 3$, and hence there are no solutions for w_{i_1}, w_{i_2} in (5.196) if we are in case (vi), or for $w_{i_1}, w_{i_2}, w_{i_3}$ in (5.197).

In total, we obtain $1+(9+12)\frac{m}{2}+27\binom{m/2}{2} = \frac{(3m+2)(9m+4)}{8}$ elements in $\operatorname{Fix}_{NC^m(E_7)}(\phi^p)$, which agrees with the limit in (5.173g).

Finally, we turn to (5.173i). By Remark 3, the only choices for h_2 and m_2 to be considered are $h_2 = 1$ and $m_2 = 7$, $h_2 = 1$ and $m_2 = 5$, $h_2 = 2$ and $m_2 = 7$, $h_2 = 2$ and $m_2 = 5$, $h_2 = 2$ and $m_2 = 4$, respectively $h_2 = m_2 = 2$. These correspond to the choices $p = 18m/7$, $p = 18m/5$, $p = 9m/7$, $p = 9m/5$, $p = 9m/4$, respectively $p = 9m/2$, all of which have already been discussed as they do not belong to (5.173i). Hence, (3.3) must necessarily hold, as required.

Case $G_{37} = E_8$. The degrees are $2, 8, 12, 14, 18, 20, 24, 30$, and hence we have

$$\operatorname{Cat}^m(E_8; q) = \frac{[30m+30]_q\,[30m+24]_q\,[30m+20]_q\,[30m+18]_q}{[30]_q\,[24]_q\,[20]_q\,[18]_q} \\ \times \frac{[30m+14]_q\,[30m+12]_q\,[30m+8]_q\,[30m+2]_q}{[14]_q\,[12]_q\,[8]_q\,[2]_q}.$$

Let ζ be a $30m$-th root of unity. The following cases on the right-hand side of (3.3) do occur:

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q)=m+1,\quad \text{if } \zeta=\zeta_{30},\zeta_{15}, \tag{5.198a}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q)=\tfrac{5m+4}{4},\quad \text{if } \zeta=\zeta_{24},\ 4\mid m, \tag{5.198b}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q)=\tfrac{3m+2}{2},\quad \text{if } \zeta=\zeta_{20},\ 2\mid m, \tag{5.198c}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q)=\tfrac{5m+3}{3},\quad \text{if } \zeta=\zeta_{18},\zeta_9,\ 3\mid m, \tag{5.198d}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q)=\tfrac{15m+7}{7},\quad \text{if } \zeta=\zeta_{14},\zeta_7,\ 7\mid m, \tag{5.198e}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q)=\tfrac{(5m+4)(5m+2)}{8},\quad \text{if } \zeta=\zeta_{12},\ 2\mid m, \tag{5.198f}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q)=\tfrac{(m+1)(3m+2)}{2},\quad \text{if } \zeta=\zeta_{10},\zeta_5, \tag{5.198g}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q)=\frac{(5m+4)(15m+4)}{16},\quad \text{if } \zeta=\zeta_8,\ 4\mid m, \tag{5.198h}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q)=\frac{(m+1)(5m+4)(5m+3)(5m+2)}{24},\quad \text{if } \zeta=\zeta_6,\zeta_3, \tag{5.198i}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q)=\frac{(5m+4)(3m+2)(5m+2)(15m+4)}{64},\quad \text{if } \zeta=\zeta_4,\ 2\mid m, \tag{5.198j}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q)=\operatorname{Cat}^m(E_8),\quad \text{if } \zeta=-1 \text{ or } \zeta=1, \tag{5.198k}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q)=1,\quad \text{otherwise.} \tag{5.198l}$$

We must now prove that the left-hand side of (3.3) in each case agrees with the values exhibited in (5.198). The only cases not covered by Lemmas 4 and 5 are the ones in (5.198b), (5.198c), (5.198d), (5.198e), (5.198f), (5.198h), (5.198j), and (5.198l).

We begin with the case in (5.198b). By Lemma 3, we are free to choose $p=5m/4$. In particular, m must be divisible by 4. From (4.1), we infer

$$\begin{aligned}\phi^p\big((w_0;w_1,\dots,w_m)\big)&\\ &=\big(*;c^2w_{\frac{3m}{4}+1}c^{-2},c^2w_{\frac{3m}{4}+2}c^{-2},\dots,c^2w_mc^{-2},cw_1c^{-1},\dots,cw_{\frac{3m}{4}}c^{-1}\big).\end{aligned}$$

Supposing that $(w_0;w_1,\dots,w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i=c^2w_{\frac{3m}{4}+i}c^{-2},\quad i=1,2,\dots,\tfrac{m}{4}, \tag{5.199a}$$

$$w_i=cw_{i-\frac{m}{4}}c^{-1},\quad i=\tfrac{m}{4}+1,\tfrac{m}{4}+2,\dots,m. \tag{5.199b}$$

There are several distinct possibilities for choosing the w_i's, $1\le i\le m$, which we summarise as follows:

(i) all the w_i's are equal to ε (and $w_0=c$),
(ii) there is an i with $1\le i\le\frac{m}{4}$ such that

$$1\le\ell_T(w_i)=\ell_T(w_{i+\frac{m}{4}})=\ell_T(w_{i+\frac{2m}{4}})=\ell_T(w_{i+\frac{3m}{4}})\le 2. \tag{5.200}$$

Moreover, since $(w_0;w_1,\dots,w_m)\in NC^m(E_8)$, we must have

$$w_iw_{i+\frac{m}{4}}w_{i+\frac{2m}{4}}w_{i+\frac{3m}{4}}\le_T c.$$

Together with Equations (5.199)–(5.200), this implies that

$$w_i = c^5 w_i c^{-5} \quad \text{and} \quad w_i(cw_ic^{-1})(c^2w_ic^{-2})(c^3w_ic^{-3}) \le_T c. \tag{5.201}$$

With the help of Stembridge's *Maple* package `coxeter` [29], one obtains 5 solutions for w_i in (5.201) with $\ell_T(w_i)=2$:

$$w_i \in \big\{[1,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4],\ [3,4,3,5],\ [2,4,5,4,2,6],\\ [1,3,4,5,6,5,4,3,1,7],\ [2,3,4,5,6,7,6,5,4,2,3,8]\big\},$$

where we have again used the short notation of `coxeter`, $\{s_1,s_2,s_3,s_4,s_5,s_6,s_7,s_8\}$ being a simple system of generators of E_8, corresponding to the Dynkin diagram displayed in Figure 6. Each of the above solutions for w_i gives rise to $m/4$ elements of $\operatorname{Fix}_{NC^m(E_8)}(\phi^p)$ since i ranges from 1 to $m/4$.



FIGURE 6. The Dynkin diagram for E_8

There are no solutions for w_i in (5.201) with $\ell_T(w_i)=1$.

In total, we obtain $1+5\frac{m}{4}=\frac{5m+4}{4}$ elements in $\operatorname{Fix}_{NC^m(E_8)}(\phi^p)$, which agrees with the limit in (5.198b).

Next we discuss the case in (5.198c). By Lemma 3, we are free to choose $p=3m/2$. In particular, m must be divisible by 2. From (4.1), we infer

$$\phi^p\big((w_0;w_1,\dots,w_m)\big) = (*;c^2w_{\frac{m}{2}+1}c^{-2},c^2w_{\frac{m}{2}+2}c^{-2},\dots,c^2w_mc^{-2},cw_1c^{-1},\dots,cw_{\frac{m}{2}}c^{-1}\big).$$

Supposing that $(w_0;w_1,\dots,w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^2 w_{\frac{m}{2}+i}c^{-2}, \quad i=1,2,\dots,\tfrac{m}{2}, \tag{5.202a}$$

$$w_i = cw_{i-\frac{m}{2}}c^{-1}, \quad i=\tfrac{m}{2}+1,\tfrac{m}{2}+2,\dots,m. \tag{5.202b}$$

There are several distinct possibilities for choosing the w_i's, $1\le i\le m$, which we summarise as follows:

(i) all the w_i's are equal to ε (and $w_0=c$),

(ii) there is an i with $1\le i\le \frac{m}{2}$ such that

$$1\le \ell_T(w_i)=\ell_T(w_{i+\frac{m}{2}})\le 4. \tag{5.203}$$

Moreover, since $(w_0;w_1,\dots,w_m)\in NC^m(E_8)$, we must have $w_iw_{i+\frac{m}{2}}\le_T c$. Together with Equations (5.202)–(5.203), this implies that

$$w_i = c^3w_ic^{-3}, \quad w_i(cw_ic^{-1})\le_T c, \quad \text{and} \quad 1\le \ell_T(w_i)\le 4. \tag{5.204}$$

With the help of Stembridge's *Maple* package `coxeter` [29], one obtains three solutions for w_i in (5.204) with $\ell_T(w_i)=4$:

$$w_i \in \big\{[1,2,3,4,2,5,4,2,6,5,7,8,7,6,5,4,2,3],\ [2,3,4,2,5,6,5,4,2,7],\\ [1,2,4,2,3,4,5,4,6,7,6,5,4,3,1,8]\big\},$$

where we used again `coxeter`'s short notation, $\{s_1,s_2,s_3,s_4,s_5,s_6,s_7,s_8\}$ being a simple system of generators of E_8, corresponding to the Dynkin diagram displayed in Figure 6.

Each of these solutions for w_i gives rise to $m/2$ elements of $\mathrm{Fix}_{NC^m(E_8)}(\phi^p)$ since i ranges from 1 to $m/2$. There are no solutions for w_i in (5.204) with $1 \le \ell_T(w_i) \le 3$.

In total, we obtain $1+3\frac{m}{2} = \frac{3m+2}{2}$ elements in $\mathrm{Fix}_{NC^m(E_8)}(\phi^p)$, which agrees with the limit in (5.198c).

Next we consider the case in (5.198d). If $\zeta = \zeta_{18}$, then, by Lemma 3, we are free to choose $p = 5m/3$, whereas, for $\zeta = \zeta_9$, we can choose $p = 10m/3$. In particular, in both cases m must be divisible by 3.

First, let $p = 5m/3$. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) = (*; c^2 w_{\frac{m}{3}+1} c^{-2}, c^2 w_{\frac{m}{3}+2} c^{-2}, \dots, c^2 w_m c^{-2}, c w_1 c^{-1}, \dots, c w_{\frac{m}{3}} c^{-1}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^2 w_{\frac{m}{3}+i} c^{-2}, \quad i = 1, 2, \dots, \tfrac{2m}{3}, \tag{5.205a}$$

$$w_i = c w_{i-\frac{2m}{3}} c^{-1}, \quad i = \tfrac{2m}{3}+1, \tfrac{2m}{3}+2, \dots, m. \tag{5.205b}$$

There several distinct possibilities for choosing the w_i's, $1 \le i \le m$, which we summarise as follows:

(i) all the w_i's are equal to ε (and $w_0 = c$),
(ii) there is an i with $1 \le i \le \frac{m}{3}$ such that

$$1 \le \ell_T(w_i) = \ell_T(w_{i+\frac{m}{3}}) = \ell_T(w_{i+\frac{2m}{3}}) \le 2. \tag{5.206}$$

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(E_8)$, we must have

$$w_i w_{i+\frac{m}{3}} w_{i+\frac{2m}{3}} \le_T c.$$

Together with Equations (5.205)–(5.206), this implies that

$$w_i = c^5 w_i c^{-5} \quad \text{and} \quad w_i (c^3 w_i c^{-3})(c w_i c^{-1}) \le_T c. \tag{5.207}$$

With the help of Stembridge's *Maple* package `coxeter` [29], one obtains five solutions for w_i in (5.207) with $\ell_T(w_i) = 2$:

$$w_i \in \big\{[1,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4],\ [3,4,3,5],\ [2,4,5,4,2,6],\\ [1,3,4,5,6,5,4,3,1,7],\ [2,3,4,5,6,7,6,5,4,2,3,8]\big\},$$

where $\{s_1, s_2, s_3, s_4, s_5, s_6, s_7, s_8\}$ is a simple system of generators of E_8, corresponding to the Dynkin diagram displayed in Figure 6, and each of them gives rise to $m/3$ elements of $\mathrm{Fix}_{NC^m(E_8)}(\phi^p)$ since i ranges from 1 to $m/3$. There are no solutions for w_i in (5.207) with $\ell_T(w_i) = 1$.

In total, we obtain $1+5\frac{m}{3} = \frac{5m+3}{3}$ elements in $\mathrm{Fix}_{NC^m(E_8)}(\phi^p)$, which agrees with the limit in (5.198d).

Now let $p = 10m/3$. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big)\\ = (*; c^4 w_{\frac{2m}{3}+1} c^{-4}, c^4 w_{\frac{2m}{3}+2} c^{-4}, \dots, c^4 w_m c^{-4}, c^3 w_1 c^{-3}, \dots, c^3 w_{\frac{2m}{3}} c^{-3}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^4 w_{\frac{2m}{3}+i} c^{-4}, \quad i = 1, 2, \dots, \tfrac{m}{3}, \tag{5.208a}$$

$$w_i = c^3 w_{i-\frac{m}{3}} c^{-3}, \quad i = \tfrac{m}{3}+1, \tfrac{m}{3}+2, \dots, m. \tag{5.208b}$$

There several distinct possibilities for choosing the w_i's, $1 \le i \le m$, which we summarise as follows:

(i) all the w_i's are equal to ε (and $w_0 = c$),
(ii) there is an i with $1 \le i \le \frac{m}{3}$ such that

$$1 \le \ell_T(w_i) = \ell_T(w_{i+\frac{m}{3}}) = \ell_T(w_{i+\frac{2m}{3}}) \le 2. \tag{5.209}$$

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(E_8)$, we must have

$$w_i w_{i+\frac{m}{3}} w_{i+\frac{2m}{3}} \le_T c.$$

Together with Equations (5.208)–(5.209), this implies that

$$w_i = c^{10} w_i c^{-10} \quad \text{and} \quad w_i (c^3 w_i c^{-3})(c^6 w_i c^{-6}) \le_T c. \tag{5.210}$$

Due to Lemma 6 with $d = 2$, we have $c^{15} w_i c^{-15} = w_i$, hence also $c^5 w_i c^{-5} = w_i$, so that (5.210) reduces to (5.207). Therefore, we are facing exactly the same enumeration problem here as for $p = 5m/3$, and, consequently, the number of solutions to (5.210) is the same, namely $\frac{5m+3}{3}$, as required.

Our next case is the case in (5.198e). If $\zeta = \zeta_{14}$, then, by Lemma 3, we are free to choose $p = 15m/7$, whereas, for $\zeta = \zeta_7$, we can choose $p = 30m/7$. In particular, in both cases m must be divisible by 7.

First, let $p = 15m/7$. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = (*; c^3 w_{\frac{6m}{7}+1} c^{-3}, c^3 w_{\frac{6m}{7}+2} c^{-3}, \dots, c^3 w_m c^{-3}, c^2 w_1 c^{-2}, \dots, c^2 w_{\frac{6m}{7}} c^{-2}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^3 w_{\frac{6m}{7}+i} c^{-3}, \quad i = 1, 2, \dots, \tfrac{m}{7}, \tag{5.211a}$$

$$w_i = c^2 w_{i-\frac{m}{7}} c^{-2}, \quad i = \tfrac{m}{7}+1, \tfrac{m}{7}+2, \dots, m. \tag{5.211b}$$

There are two distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),
(ii) there is an i with $1 \le i \le \frac{m}{7}$ such that

$$\ell_T(w_i) = \ell_T(w_{i+\frac{m}{7}}) = \ell_T(w_{i+\frac{2m}{7}}) = \ell_T(w_{i+\frac{3m}{7}}) \\ = \ell_T(w_{i+\frac{4m}{7}}) = \ell_T(w_{i+\frac{5m}{7}}) = \ell_T(w_{i+\frac{6m}{7}}) = 1, \tag{5.212}$$

and the other w_j's, $1 \le j \le m$, are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(E_8)$, we must have

$$w_i w_{i+\frac{m}{7}} w_{i+\frac{2m}{7}} w_{i+\frac{3m}{7}} w_{i+\frac{4m}{7}} w_{i+\frac{5m}{7}} w_{i+\frac{6m}{7}} \le_T c.$$

Together with the Equations (5.211)–(5.212), this implies that

$$w_i = c^{15} w_i c^{-15} \\ \text{and} \quad w_i (c^2 w_i c^{-2})(c^4 w_i c^{-4})(c^6 w_i c^{-6})(c^8 w_i c^{-8})(c^{10} w_i c^{-10})(c^{12} w_i c^{-12}) \le_T c. \tag{5.213}$$

Here, the first equation is automatically satisfied due to Lemma 6 with $d = 2$.

With the help of Stembridge's *Maple* package `coxeter` [29], one obtains 15 solutions for w_i in (5.213):

$$\begin{aligned}w_i \in \big\{&[4],\ [5],\ [6],\ [7],\ [8],\ [3,4,3],\ [2,4,5,4,2],\ [3,4,5,4,3],\ [2,4,5,6,5,4,2],\\ &[1,3,4,5,6,5,4,3,1],\ [1,3,4,5,6,7,6,5,4,3,1],\\ &[2,3,4,5,6,7,6,5,4,2,3],\ [2,3,4,5,6,7,8,7,6,5,4,2,3],\\ &[1,2,3,4,5,6,7,8,7,6,5,4,2,3,1],\ [1,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4]\big\},\end{aligned}$$

where we have again used the short notation of `coxeter`, $\{s_1,s_2,s_3,s_4,s_5,s_6,s_7,s_8\}$ being a simple system of generators of E_8, corresponding to the Dynkin diagram displayed in Figure 6, and each of them gives rise to $m/7$ elements of $\operatorname{Fix}_{NC^m(E_8)}(\phi^p)$ since i ranges from 1 to $m/7$.

In total, we obtain $1+15\frac{m}{7}=\frac{15m+7}{7}$ elements in $\operatorname{Fix}_{NC^m(E_8)}(\phi^p)$, which agrees with the limit in (5.198e).

Now let $p=30m/7$. From (4.1), we infer

$$\begin{aligned}&\phi^p\big((w_0;w_1,\dots,w_m)\big)\\ &\qquad=\big(*;c^5w_{\frac{5m}{7}+1}c^{-5},c^5w_{\frac{5m}{7}+2}c^{-5},\dots,c^5w_mc^{-5},c^4w_1c^{-4},\dots,c^4w_{\frac{5m}{7}}c^{-4}\big).\end{aligned}$$

Supposing that $(w_0;w_1,\dots,w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i=c^5w_{\frac{5m}{7}+i}c^{-5},\quad i=1,2,\dots,\tfrac{2m}{7},\tag{5.214a}$$

$$w_i=c^4w_{i-\frac{2m}{7}}c^{-4},\quad i=\tfrac{2m}{7}+1,\tfrac{2m}{7}+2,\dots,m.\tag{5.214b}$$

There are two distinct possibilities for choosing the w_i's, $1\le i\le m$:

(i) all the w_i's are equal to ε (and $w_0=c$),
(ii) there is an i with $1\le i\le\frac{m}{7}$ such that

$$\begin{aligned}\ell_T(w_i)=\ell_T(w_{i+\frac{m}{7}})=\ell_T(w_{i+\frac{2m}{7}})=\ell_T(w_{i+\frac{3m}{7}})&\\ =\ell_T(w_{i+\frac{4m}{7}})=\ell_T(w_{i+\frac{5m}{7}})=\ell_T(w_{i+\frac{6m}{7}})=1,&\end{aligned}\tag{5.215}$$

and the other w_j's, $1\le j\le m$, are equal to ε.

Moreover, since $(w_0;w_1,\dots,w_m)\in NC^m(E_8)$, we must have

$$w_iw_{i+\frac{m}{7}}w_{i+\frac{2m}{7}}w_{i+\frac{3m}{7}}w_{i+\frac{4m}{7}}w_{i+\frac{5m}{7}}w_{i+\frac{6m}{7}}\le_T c.$$

Together with the Equations (5.214)–(5.215), this implies that

$$\begin{aligned}&w_i=c^{30}w_ic^{-30}\\ \text{and}\quad &w_i(c^{17}w_ic^{-17})(c^4w_ic^{-4})(c^{21}w_ic^{-21})(c^8w_ic^{-8})(c^{25}w_ic^{-25})(c^{12}w_ic^{-12})\le_T c.\end{aligned}\tag{5.216}$$

Here, the first equation is automatically satisfied since $c^{30}=\varepsilon$. Moreover, due to Lemma 6 with $d=2$, we have $c^{17}w_ic^{-17}=c^2w_ic^{-2}$, etc., so that (5.216) reduces to (5.213). Therefore, we are facing exactly the same enumeration problem here as for $p=15m/7$, and, consequently, the number of solutions to (5.216) is the same, namely $\frac{15m+7}{7}$, as required.

We now turn to the case in (5.198f). By Lemma 3, we are free to choose $p = 5m/2$. In particular, m must be divisible by 2. From (4.1), we infer

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = (*; c^3 w_{\frac{m}{2}+1} c^{-3}, c^3 w_{\frac{m}{2}+2} c^{-3}, \dots, c^3 w_m c^{-3}, c^2 w_1 c^{-2}, \dots, c^2 w_{\frac{m}{2}} c^{-2}\big).$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^3 w_{\frac{m}{2}+i} c^{-3}, \quad i = 1, 2, \dots, \tfrac{m}{2}, \tag{5.217a}$$

$$w_i = c^2 w_{i-\frac{m}{2}} c^{-2}, \quad i = \tfrac{m}{2}+1, \tfrac{m}{2}+2, \dots, m. \tag{5.217b}$$

There are several distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$1 \le \ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) \le 4, \tag{5.218a}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(iii) there are i_1 and i_2 with $1 \le i_1 < i_2 \le \frac{m}{2}$ such that

$$\ell_1 := \ell_T(w_{i_1}) = \ell_T(w_{i_1+\frac{m}{2}}) \ge 1, \quad \ell_2 := \ell_T(w_{i_2}) = \ell_T(w_{i_2+\frac{m}{2}}) \ge 1, \quad \text{and} \quad \ell_1 + \ell_2 \le 4, \tag{5.218b}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(iv) there are i_1, i_2, i_3 with $1 \le i_1 < i_2 < i_3 \le \frac{m}{2}$ such that

$$\ell_1 := \ell_T(w_{i_1}) = \ell_T(w_{i_1+\frac{m}{2}}) \ge 1, \quad \ell_2 := \ell_T(w_{i_2}) = \ell_T(w_{i_2+\frac{m}{2}}) \ge 1, \\ \ell_3 := \ell_T(w_{i_3}) = \ell_T(w_{i_3+\frac{m}{2}}) \ge 1, \quad \text{and} \quad \ell_1 + \ell_2 + \ell_3 \le 4, \tag{5.218c}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(v) there are i_1, i_2, i_3, i_4 with $1 \le i_1 < i_2 < i_3 < i_4 \le \frac{m}{2}$ such that

$$\ell_T(w_{i_1}) = \ell_T(w_{i_2}) = \ell_T(w_{i_3}) = \ell_T(w_{i_4}) \\ = \ell_T(w_{i_1+\frac{m}{2}}) = \ell_T(w_{i_2+\frac{m}{2}}) = \ell_T(w_{i_3+\frac{m}{2}}) = \ell_T(w_{i_4+\frac{m}{2}}) = 1, \tag{5.218d}$$

and all other w_j's are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(E_8)$, we must have $w_i w_{i+\frac{m}{2}} \le_T c$, respectively $w_{i_1} w_{i_2} w_{i_1+\frac{m}{2}} w_{i_2+\frac{m}{2}} \le_T c$, respectively

$$w_{i_1} w_{i_2} w_{i_3} w_{i_1+\frac{m}{2}} w_{i_2+\frac{m}{2}} w_{i_3+\frac{m}{2}} \le_T c,$$

respectively

$$w_{i_1} w_{i_2} w_{i_3} w_{i_4} w_{i_1+\frac{m}{2}} w_{i_2+\frac{m}{2}} w_{i_3+\frac{m}{2}} w_{i_4+\frac{m}{2}} = c.$$

Together with the Equations (5.217)–(5.218), this implies that

$$w_i = c^5 w_i c^{-5} \quad \text{and} \quad w_i(c^2 w_i c^{-2}) \le_T c, \tag{5.219}$$

respectively that

$$w_{i_1} = c^5 w_{i_1} c^{-5}, \quad w_{i_2} = c^5 w_{i_2} c^{-5}, \quad \text{and} \quad w_{i_1} w_{i_2} (c^2 w_{i_1} c^{-2})(c^2 w_{i_2} c^{-2}) \le_T c, \tag{5.220}$$

respectively that

$$w_{i_1} = c^5 w_{i_1} c^{-5}, \quad w_{i_2} = c^5 w_{i_2} c^{-5}, \quad w_{i_3} = c^5 w_{i_3} c^{-5}, \\ \text{and} \quad w_{i_1} w_{i_2} w_{i_3} (c^2 w_{i_1} c^{-2})(c^2 w_{i_2} c^{-2})(c^2 w_{i_3} c^{-2}) \le_T c, \tag{5.221}$$

respectively that

$$\begin{gathered} w_{i_1} = c^5 w_{i_1} c^{-5}, \quad w_{i_2} = c^5 w_{i_2} c^{-5}, \quad w_{i_3} = c^5 w_{i_3} c^{-5}, \quad w_{i_4} = c^5 w_{i_4} c^{-5}, \\ \text{and} \quad w_{i_1} w_{i_2} w_{i_3} w_{i_4} (c^2 w_{i_1} c^{-2})(c^2 w_{i_2} c^{-2})(c^2 w_{i_3} c^{-2})(c^2 w_{i_4} c^{-2}) = c. \end{gathered} \tag{5.222}$$

With the help of Stembridge's *Maple* package `coxeter` [29], one obtains 10 solutions for w_i in (5.219) with $\ell_T(w_i) = 2$:

$$\begin{gathered} w_i \in \big\{[1,4,2,3,4,5,6,7,6,5,4,2,3,4],\ [3,1,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,5], \\ [1,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4],\ [3,4,3,5],\ [2,4,5,4,2,6],\ [1,3,4,5,6,5,4,3,1,7], \\ [2,3,4,5,6,7,6,5,4,2,3,8],\ [2,4,2,3,4,5,6,5,4,3],\ [1,3,4,5,4,2,3,1,4,5,6,7,6,5,4,2], \\ [2,3,1,4,5,6,5,4,2,3,1,4,5,6,7,8,7,6,5,4,3,1]\big\}, \end{gathered} \tag{5.223}$$

and one obtains 10 solutions for w_i in (5.219) with $\ell_T(w_i) = 4$:

$$\begin{gathered} w_i \in \big\{[1,2,3,4,5,6,7,6,5,4,2,3,4,8],\ [1,2,4,2,3,4,5,4,3,6,5,7,6,5,4,2,3,4], \\ [1,2,3,1,4,5,6,7,8,7,6,5,4,2,3,1,4,5], \\ [1,3,1,4,5,4,2,3,4,5,6,5,4,2,7,6,8,7,6,5,4,2,3,1,4,5], \\ [1,2,3,4,2,5,4,2,3,4,6,5,7,6,5,4,3,8,7,6,5,4,2,3,1,4],\ [4,2,3,4,5,6], \\ [2,3,1,4,2,5,4,6,5,4,2,3,1,4,5,6,7,8,7,6,5,4,3,1],\ [1,5,4,2,3,1,4,5,6,7], \\ [3,1,6,5,4,2,3,1,4,3,5,6,7,8],\ [2,3,4,2,3,5,4,6,5,4,2,7,6,5,4,2,3,8]\big\}, \end{gathered} \tag{5.224}$$

where we have again used the short notation of `coxeter`, $\{s_1, s_2, s_3, s_4, s_5, s_6, s_7, s_8\}$ being a simple system of generators of E_8, corresponding to the Dynkin diagram displayed in Figure 6, and each of them gives rise to $m/2$ elements of $\operatorname{Fix}_{NC^m(E_8)}(\phi^p)$ since i ranges from 1 to $m/2$. There are no solutions for w_i in (5.219) with $\ell_T(w_i) = 1$ or $\ell_T(w_i) = 3$.

Consequently, there are no solutions in Cases (iv) and (v), and the only possible solutions occurring in Case (iii) are pairs (w_{i_1}, w_{i_2}) of elements of (5.223) whose product is in (5.224). Another computation using Stembridge's *Maple* package `coxeter` finds the following 25 pairs meeting that description:

$$\begin{gathered} w_i \in \big\{([1,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4],\ [1,4,2,3,4,5,6,7,6,5,4,2,3,4]), \\ ([2,4,5,4,2,6],\ [1,4,2,3,4,5,6,7,6,5,4,2,3,4]), \\ ([3,4,3,5],\ [3,1,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,5]), \\ ([1,3,4,5,6,5,4,3,1,7],\ [3,1,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,5]), \\ ([3,1,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,5],\ [1,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4]), \\ ([2,3,4,5,6,7,6,5,4,2,3,8],\ [1,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4]), \\ ([1,3,4,5,4,2,3,1,4,5,6,7,6,5,4,2],\ [1,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4]), \\ ([1,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4],[3,4,3,5]),\ ([2,4,2,3,4,5,6,5,4,3],\ [3,4,3,5]), \\ ([2,3,1,4,5,6,5,4,2,3,1,4,5,6,7,8,7,6,5,4,3,1],\ [3,4,3,5]), \\ ([1,4,2,3,4,5,6,7,6,5,4,2,3,4],\ [2,4,5,4,2,6]),\ ([3,4,3,5],\ [2,4,5,4,2,6]), \\ ([1,3,4,5,4,2,3,1,4,5,6,7,6,5,4,2],\ [2,4,5,4,2,6]), \\ ([3,1,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,5],\ [1,3,4,5,6,5,4,3,1,7]), \\ ([2,4,5,4,2,6],\ [1,3,4,5,6,5,4,3,1,7]), \\ ([2,3,1,4,5,6,5,4,2,3,1,4,5,6,7,8,7,6,5,4,3,1],\ [1,3,4,5,6,5,4,3,1,7]), \end{gathered}$$

$$([1,4,2,3,4,5,6,7,6,5,4,2,3,4],\ [2,3,4,5,6,7,6,5,4,2,3,8]),$$
$$([1,3,4,5,6,5,4,3,1,7],\ [2,3,4,5,6,7,6,5,4,2,3,8]),$$
$$([2,4,2,3,4,5,6,5,4,3],\ [2,3,4,5,6,7,6,5,4,2,3,8]),\ ([2,4,5,4,2,6],\ [2,4,2,3,4,5,6,5,4,3]),$$
$$([2,3,4,5,6,7,6,5,4,2,3,8],\ [2,4,2,3,4,5,6,5,4,3]),$$
$$([1,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4],\ [1,3,4,5,4,2,3,1,4,5,6,7,6,5,4,2]),$$
$$([1,3,4,5,6,5,4,3,1,7],\ [1,3,4,5,4,2,3,1,4,5,6,7,6,5,4,2]),$$
$$([3,4,3,5],\ [2,3,1,4,5,6,5,4,2,3,1,4,5,6,7,8,7,6,5,4,3,1]),$$
$$([2,3,4,5,6,7,6,5,4,2,3,8],\ [2,3,1,4,5,6,5,4,2,3,1,4,5,6,7,8,7,6,5,4,3,1])\big\},$$

where $\{s_1,s_2,s_3,s_4,s_5,s_6,s_7,s_8\}$ is a simple system of generators of E_8, corresponding to the Dynkin diagram displayed in Figure 6, and each of them gives rise to $\binom{m/2}{2}$ elements of $\operatorname{Fix}_{NC^m(E_8)}(\phi^p)$ since $1\le i_1<i_2\le \frac{m}{2}$.

In total, we obtain

$$1+20\frac{m}{2}+25\binom{m/2}{2}=\frac{(5m+4)(5m+2)}{8}$$

elements in $\operatorname{Fix}_{NC^m(E_8)}(\phi^p)$, which agrees with the limit in (5.198f).

Next we consider the case in (5.198h). By Lemma 3, we are free to choose $p=15m/4$. In particular, m must be divisible by 4. From (4.1), we infer

$$\phi^p\big((w_0;w_1,\dots,w_m)\big)\\ =(*;c^4w_{\frac{m}{4}+1}c^{-4},c^4w_{\frac{m}{4}+2}c^{-4},\dots,c^4w_mc^{-4},c^3w_1c^{-3},\dots,c^3w_{\frac{m}{4}}c^{-3}\big).$$

Supposing that $(w_0;w_1,\dots,w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i=c^4w_{\frac{m}{4}+i}c^{-4},\quad i=1,2,\dots,\tfrac{3m}{4},\tag{5.225a}$$
$$w_i=c^3w_{i-\frac{3m}{4}}c^{-3},\quad i=\tfrac{3m}{4}+1,\tfrac{3m}{4}+2,\dots,m.\tag{5.225b}$$

There are several distinct possibilities for choosing the w_i's, $1\le i\le m$, which we summarise as follows:

(i) all the w_i's are equal to ε (and $w_0=c$),

(ii) there is an i with $1\le i\le\frac{m}{4}$ such that

$$1\le\ell_T(w_i)=\ell_T(w_{i+\frac{m}{4}})=\ell_T(w_{i+\frac{2m}{4}})=\ell_T(w_{i+\frac{3m}{4}})\le 2,\tag{5.226a}$$

and the other w_j's, $1\le j\le m$, are equal to ε,

(iii) there are i_1 and i_2 with $1\le i_1<i_2\le\frac{m}{4}$ such that

$$\ell_T(w_{i_1})=\ell_T(w_{i_2})=\ell_T(w_{i_1+\frac{m}{4}})=\ell_T(w_{i_2+\frac{m}{4}})\\ =\ell_T(w_{i_1+\frac{2m}{4}})=\ell_T(w_{i_2+\frac{2m}{4}})=\ell_T(w_{i_1+\frac{3m}{4}})=\ell_T(w_{i_2+\frac{3m}{4}})=1,\tag{5.226b}$$

and all other w_j are equal to ε.

Moreover, since $(w_0;w_1,\dots,w_m)\in NC^m(E_8)$, we must have

$$w_iw_{i+\frac{m}{4}}w_{i+\frac{2m}{4}}w_{i+\frac{3m}{4}}\le_T c,$$

respectively

$$w_{i_1}w_{i_2}w_{i_1+\frac{m}{4}}w_{i_2+\frac{m}{4}}w_{i_1+\frac{2m}{4}}w_{i_2+\frac{2m}{4}}w_{i_1+\frac{3m}{4}}w_{i_2+\frac{3m}{4}}=c.$$

Together with Equations (5.225)–(5.226), this implies that

$$w_i = c^{15} w_i c^{-15} \quad \text{and} \quad w_i(c^{11} w_i c^{-11})(c^7 w_i c^{-7})(c^3 w_i c^{-3}) \le_T c, \tag{5.227}$$

respectively that

$$\begin{multline} w_{i_1} = c^{15} w_{i_1} c^{-15}, \quad w_{i_1} = c^{15} w_{i_2} c^{-15},\\ \text{and} \quad w_{i_1} w_{i_2} (c^{11} w_{i_1} c^{-11})(c^{11} w_{i_2} c^{-11})(c^7 w_{i_1} c^{-7})(c^7 w_{i_2} c^{-7})(c^3 w_{i_1} c^{-3})(c^3 w_{i_2} c^{-3}) = c. \end{multline} \tag{5.228}$$

Here, the first equation in (5.227) and the first two equations in (5.228) are automatically satisfied due to Lemma 6 with $d = 2$.

With the help of Stembridge's *Maple* package `coxeter` [29], one obtains 30 solutions for w_i in (5.227) with $\ell_T(w_i) = 1$:

$$\begin{gathered} w_i \in \big\{[4],\ [5],\ [6],\ [7],\ [8],\ [3,4,3],\ [4,5,4],\ [5,6,5],\ [6,7,6],\ [7,8,7],\ [2,4,5,4,2],\ [3,4,5,4,3],\\ [2,4,5,6,5,4,2],\ [1,3,4,5,6,5,4,3,1],\ [4,2,3,4,5,4,2,3,4],\ [1,3,4,5,6,7,6,5,4,3,1],\\ [2,3,4,5,6,7,6,5,4,2,3],\ [1,2,3,4,5,6,7,6,5,4,2,3,1],\ [5,4,2,3,4,5,6,5,4,2,3,4,5],\\ [2,3,4,5,6,7,8,7,6,5,4,2,3],\ [1,5,4,2,3,4,5,6,5,4,2,3,1,4,5],\\ [1,2,3,4,5,6,7,8,7,6,5,4,2,3,1],\ [4,2,3,4,5,6,7,8,7,6,5,4,2,3,4],\\ [1,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4],\ [3,1,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3],\\ [1,6,5,4,2,3,4,5,6,7,6,5,4,2,3,1,4,5,6],\ [3,1,6,5,4,2,3,4,5,6,7,6,5,4,2,3,1,4,3,5,6],\\ [4,3,1,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,4],\\ [3,1,7,6,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,6,7],\\ [2,4,3,1,7,6,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,4,2,6,7]\big\}, \end{gathered}$$

one obtains 45 solutions for w_i in (5.227) with $\ell_T(w_i) = 2$ and w_i of type A_1^2 (as a parabolic Coxeter element; see the end of Section 2):

$$\begin{gathered} w_i \in \{[1,2,3,1,4,5,6,5,4,2,3,4,5,7,6,5,4,2,3,1,4,5,8,7,6,5,4,2,3,1],\\ [1,2,3,4,2,3,1,7,6,5,4,2,3,1,4,3,5,4,2,6,7,8,7,6,5,4,2,3,1,4,3,5,4,2,6,7],\\ [1,2,3,4,2,5,4,6,5,4,2,7,6,5,4,2,3,1],\ [1,2,3,4,3,5,4,6,5,4,3,7,6,5,4,2,3,1],\\ [1,2,3,4,5,4,6,5,4,7,8,7,6,5,4,2,3,1],\ [1,2,3,4,5,4,6,7,6,5,4,2,3,1],\\ [1,2,3,4,5,6,5,4,2,3,4,5,7,6,5,4,2,3,4,5,8,7,6,5,4,2,3,1],\ [1,2,6,5,4,2,3,1,7,6,5,4,2,3,1,4,3,5,4,2,6,5,7,6],\\ [1,3,1,4,2,3,4,5,4,6,5,7,6,5,4,2,8,7,6,5,4,2,3,1,4,3],\\ [1,3,1,4,2,7,6,5,4,2,3,1,8,7,6,5,4,2,3,1,4,3,5,4,2,6,5,4,3,1,7,6,8,7],\\ [1,3,4,3,5,4,3,6,5,4,3,1],\ [1,3,4,5,4,6,5,4,7,6,5,4,3,1],\ [1,4,2,3,1,4,3,5,4,3,1,6,7,8,7,6,5,4,2,3,1,4],\\ [1,4,2,3,4,5,6,5,7,6,5,8,7,6,5,4,2,3,1,4],\ [1,4,5,4,2,3,4,5,6,5,4,2,3,1,4,5],\\ [1,5,4,2,3,1,4,5,6,5,4,2,3,1,4,5],\ [1,5,4,2,3,4,5,6,5,4,2,3,1,4,5,8],\\ [1,5,6,5,4,2,3,4,5,6,7,6,5,4,2,3,1,4,5,6],\ [2,3,1,4,2,3,4,5,4,6,5,7,6,5,4,3,8,7,6,5,4,2,3,1,4,3],\\ [2,3,1,6,5,4,2,3,1,4,3,5,6,7,6,5,4,2,3,1,4,3,5,6],\\ [2,3,4,2,3,1,4,7,6,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,4,2,6,7],\\ [2,3,4,2,3,5,4,2,3,6,7,8,7,6,5,4,2,3],\ [2,3,4,5,4,6,5,4,7,6,5,4,2,3],\ [2,3,4,5,6,5,7,6,5,8,7,6,5,4,2,3],\\ [2,4,2,3,4,5,4,3,6,7,8,7,6,5,4,2,3,4],\ [2,4,2,5,4,2,6,5,4,2],\\ [2,4,3,1,6,7,6,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,4,2,6,7],\ [2,4,5,4,2,7,8,7],\\ [3,1,4,2,3,4,5,6,5,7,8,7,6,5,4,2,3,1,4,3],\ [3,1,4,7,6,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,6,7],\\ [3,1,5,6,5,4,2,3,4,5,6,7,6,5,4,2,3,1,4,3,5,6],\ [3,1,6,7,6,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,6,7],\\ [3,4,2,3,1,5,4,2,3,1,4,8,7,6,5,4,2,3,1,4,3,5,4,2,6,5,4,3,7,8],\ [3,4,2,3,4,5,4,2,6,7,8,7,6,5,4,2,3,4], \end{gathered}$$

$$\begin{gathered}[3,4,3,6,7,6],\ [3,4,5,4,3,7,8,7],\ [4,2,3,1,5,4,2,3,1,4,3,5,6,7,8,7,6,5,4,2,3,1,4,3,5,4],\\ [4,2,3,4,5,4,2,3,4,7],\ [4,2,3,4,5,6,7,6,8,7,6,5,4,2,3,4],\ [4,3,1,5,4,2,3,4,5,6,7,6,8,7,6,5,4,2,3,1,4,3,5,4],\\ [4,5,4,2,3,4,5,6,5,4,2,3,4,5],\ [4,5,4,8],\ [4,7,8,7],\\ [5,4,2,3,4,5,6,5,4,2,3,4,5,8],\ [5,4,3,1,6,5,4,2,3,1,4,3,5,4,6,5]\big\}, \qquad (5.229)\end{gathered}$$

and one obtains 20 solutions for w_i in (5.227) with $\ell_T(w_i) = 2$ and w_i of type A_2:

$$\begin{gathered}w_i \in \big\{[1,2,3,1,4,5,6,7,8,7,6,5,4,2,3,1,4,3],\ [1,2,3,4,5,6,7,6,5,4,2,3,1,8],\\ [1,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4],\ [1,2,4,5,4,2,3,4,5,6,5,4,3,1],\\ [1,2,4,5,6,5,4,2,3,4,5,6,7,6,5,4,3,1],\ [1,3,1,4,5,6,5,4,2,3,1,4,5,6,7,6,5,4,2,3],\\ [1,3,1,4,5,6,7,6,5,4,2,3,1,4,5,6,7,8,7,6,5,4,2,3],\ [1,3,4,5,6,5,4,3,1,7],\\ [1,4,2,3,1,4,5,4,6,7,8,7,6,5,4,2,3,1,4,3,5,4],\\ [2,3,4,3,1,5,6,7,6,5,4,2,3,1,4,3,5,6,7,8,7,6,5,4,2,3,1,4],\ [2,3,4,5,6,7,6,5,4,2,3,8],\\ [2,3,4,5,6,7,8,7,6,5,4,2,3,4],\ [2,4,5,4,2,6],\ [3,4,2,3,4,5,4,2],\ [3,4,3,5],\\ [3,4,5,4,2,3,4,5,6,5,4,2],\ [4,5],\ [5,6],\ [6,7],\ [7,8]\big\}, \qquad (5.230)\end{gathered}$$

where $\{s_1, s_2, s_3, s_4, s_5, s_6, s_7, s_8\}$ is a simple system of generators of E_8, corresponding to the Dynkin diagram displayed in Figure 6, and each of them gives rise to $m/4$ elements of $\mathrm{Fix}_{NC^m(E_8)}(\phi^p)$ since i ranges from 1 to $m/4$.

The number of solutions in Case (iii) can be computed from our knowledge of the solutions in Case (ii) according to type, using some elementary counting arguments. Namely, the number of solutions of (5.228) is equal to

$$45 \cdot 2 + 20 \cdot 3 = 150,$$

since an element of type A_1^2 can be decomposed in two ways into a product of two elements of absolute length 1, while for an element of type A_2 this can be done in 3 ways.

In total, we obtain $1 + (30 + 45 + 20)\frac{m}{4} + 150\binom{m/4}{2} = \frac{(5m+4)(15m+4)}{16}$ elements in $\mathrm{Fix}_{NC^m(E_8)}(\phi^p)$, which agrees with the limit in (5.198h).

Finally we discuss the case in (5.198j). By Lemma 3, we are free to choose $p = 15m/2$. In particular, m must be divisible by 2. From (4.1), we infer

$$\begin{multline*}\phi^p\big((w_0; w_1, \dots, w_m)\big)\\ = \big(*; c^8 w_{\frac{m}{2}+1} c^{-8}, c^8 w_{\frac{m}{2}+2} c^{-8}, \dots, c^8 w_m c^{-8}, c^7 w_1 c^{-7}, \dots, c^7 w_{\frac{m}{2}} c^{-7}\big).\end{multline*}$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ϕ^p, we obtain the system of equations

$$w_i = c^8 w_{\frac{m}{2}+i} c^{-8}, \quad i = 1, 2, \dots, \tfrac{m}{2}, \qquad (5.231\text{a})$$

$$w_i = c^7 w_{i-\frac{m}{2}} c^{-7}, \quad i = \tfrac{m}{2}+1, \tfrac{m}{2}+2, \dots, m. \qquad (5.231\text{b})$$

There are several distinct possibilities for choosing the w_i's, $1 \le i \le m$:

(i) all the w_i's are equal to ε (and $w_0 = c$),

(ii) there is an i with $1 \le i \le \frac{m}{2}$ such that

$$1 \le \ell_T(w_i) = \ell_T(w_{i+\frac{m}{2}}) \le 4, \qquad (5.232\text{a})$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(iii) there are i_1 and i_2 with $1 \le i_1 < i_2 \le \frac{m}{2}$ such that

$$\ell_1 := \ell_T(w_{i_1}) = \ell_T(w_{i_1+\frac{m}{2}}) \ge 1, \quad \ell_2 := \ell_T(w_{i_2}) = \ell_T(w_{i_2+\frac{m}{2}}) \ge 1, \quad \text{and} \quad \ell_1 + \ell_2 \le 4, \tag{5.232b}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(iv) there are i_1, i_2, i_3 with $1 \le i_1 < i_2 < i_3 \le \frac{m}{2}$ such that

$$\begin{aligned}\ell_1 := \ell_T(w_{i_1}) = \ell_T(w_{i_1+\frac{m}{2}}) \ge 1, \quad \ell_2 := \ell_T(w_{i_2}) = \ell_T(w_{i_2+\frac{m}{2}}) \ge 1,\\ \ell_3 := \ell_T(w_{i_3}) = \ell_T(w_{i_3+\frac{m}{2}}) \ge 1, \quad \text{and} \quad \ell_1 + \ell_2 + \ell_3 \le 4,\end{aligned} \tag{5.232c}$$

and the other w_j's, $1 \le j \le m$, are equal to ε,

(v) there are i_1, i_2, i_3, i_4 with $1 \le i_1 < i_2 < i_3 < i_4 \le \frac{m}{2}$ such that

$$\begin{aligned}\ell_T(w_{i_1}) = \ell_T(w_{i_2}) = \ell_T(w_{i_3}) = \ell_T(w_{i_4})\\ = \ell_T(w_{i_1+\frac{m}{2}}) = \ell_T(w_{i_2+\frac{m}{2}}) = \ell_T(w_{i_3+\frac{m}{2}}) = \ell_T(w_{i_4+\frac{m}{2}}) = 1,\end{aligned} \tag{5.232d}$$

and all other w_j's are equal to ε.

Moreover, since $(w_0; w_1, \dots, w_m) \in NC^m(E_8)$, we must have $w_i w_{i+\frac{m}{2}} \le_T c$, respectively $w_{i_1} w_{i_2} w_{i_1+\frac{m}{2}} w_{i_2+\frac{m}{2}} \le_T c$, respectively

$$w_{i_1} w_{i_2} w_{i_3} w_{i_1+\frac{m}{2}} w_{i_2+\frac{m}{2}} w_{i_3+\frac{m}{2}} \le_T c,$$

respectively

$$w_{i_1} w_{i_2} w_{i_3} w_{i_4} w_{i_1+\frac{m}{2}} w_{i_2+\frac{m}{2}} w_{i_3+\frac{m}{2}} w_{i_4+\frac{m}{2}} = c.$$

Together with Equations (5.231)–(5.232), this implies that

$$w_i = c^{15} w_i c^{-15} \quad \text{and} \quad w_i(c^7 w_i c^{-7}) \le_T c, \tag{5.233}$$

respectively that

$$w_{i_1} = c^{15} w_{i_1} c^{-15}, \quad w_{i_2} = c^{15} w_{i_2} c^{-15}, \quad \text{and} \quad w_{i_1} w_{i_2} (c^7 w_{i_1} c^{-7})(c^7 w_{i_2} c^{-7}) \le_T c, \tag{5.234}$$

respectively that

$$\begin{aligned}w_{i_1} = c^{15} w_{i_1} c^{-15}, \quad w_{i_2} = c^{15} w_{i_2} c^{-15}, \quad w_{i_3} = c^{15} w_{i_3} c^{-15},\\ \text{and} \quad w_{i_1} w_{i_2} w_{i_3} (c^7 w_{i_1} c^{-7})(c^7 w_{i_2} c^{-7})(c^7 w_{i_3} c^{-7}) \le_T c,\end{aligned} \tag{5.235}$$

respectively that

$$\begin{aligned}w_{i_1} = c^{15} w_{i_1} c^{-15}, \quad w_{i_2} = c^{15} w_{i_2} c^{-15}, \quad w_{i_3} = c^{15} w_{i_3} c^{-15}, \quad w_{i_4} = c^{15} w_{i_4} c^{-15},\\ \text{and} \quad w_{i_1} w_{i_2} w_{i_3} w_{i_4} (c^7 w_{i_1} c^{-7})(c^7 w_{i_2} c^{-7})(c^7 w_{i_3} c^{-7})(c^7 w_{i_4} c^{-7}) = c.\end{aligned} \tag{5.236}$$

Here, the first equation in (5.233), the first two in (5.234), the first three in (5.235), and the first four in (5.236), are all automatically satisfied due to Lemma 6 with $d = 2$.

With the help of Stembridge's *Maple* package `coxeter` [29], one obtains 45 solutions for w_i in (5.233) with $\ell_T(w_i)=1$:

$$\begin{gathered}
w_i \in \big\{[1],\ [3],\ [4],\ [5],\ [6],\ [7],\ [8],\ [2,4,2],\ [3,4,3],\ [4,5,4],\ [5,6,5],\ [6,7,6],\ [7,8,7],\\
[2,4,5,4,2],\ [3,4,5,4,3],\ [1,3,4,5,4,3,1],\ [2,4,5,6,5,4,2],\ [3,4,5,6,5,4,3],\\
[1,3,4,5,6,5,4,3,1],\ [4,2,3,4,5,4,2,3,4],\ [2,3,4,5,6,5,4,2,3],\ [2,4,5,6,7,6,5,4,2],\\
[1,3,4,5,6,7,6,5,4,3,1],\ [4,2,3,4,5,6,5,4,2,3,4],\ [2,3,4,5,6,7,6,5,4,2,3],\\
[1,2,3,4,5,6,7,6,5,4,2,3,1],\ [1,3,4,5,6,7,8,7,6,5,4,3,1],\ [5,4,2,3,4,5,6,5,4,2,3,4,5],\\
[4,2,3,4,5,6,7,6,5,4,2,3,4],\ [2,3,4,5,6,7,8,7,6,5,4,2,3],\ [1,5,4,2,3,4,5,6,5,4,2,3,1,4,5],\\
[1,4,2,3,4,5,6,7,6,5,4,2,3,1,4],\ [1,2,3,4,5,6,7,8,7,6,5,4,2,3,1],\\
[4,2,3,4,5,6,7,8,7,6,5,4,2,3,4],\ [1,5,4,2,3,4,5,6,7,6,5,4,2,3,1,4,5],\\
[1,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4],\ [3,1,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3],\\
[1,6,5,4,2,3,4,5,6,7,6,5,4,2,3,1,4,5,6],\ [1,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,5],\\
[3,1,6,5,4,2,3,4,5,6,7,6,5,4,2,3,1,4,3,5,6],\\
[3,1,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5],\\
[4,3,1,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,4],\\
[3,1,6,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,6],\\
[3,1,7,6,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,6,7],\\
[2,4,3,1,7,6,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,4,2,6,7]\big\},
\end{gathered}$$

one obtains 150 solutions for w_i in (5.233) with $\ell_T(w_i)=2$ and w_i of type A_1^2:

$$\begin{gathered}
w_i \in \big\{[1,2,3,1,4,3,5,4,6,5,7,6,5,4,3,1,8,7,6,5,4,2,3,1],\\
[1,2,3,1,4,5,6,5,4,2,3,4,5,7,6,5,4,2,3,1,4,5,8,7,6,5,4,2,3,1],\ [1,2,3,1,4,5,6,7,8,7,6,5,4,2,3,1],\\
[1,2,3,4,2,3,1,7,6,5,4,2,3,1,4,3,5,4,2,6,7,8,7,6,5,4,2,3,1,4,3,5,4,2,6,7],\\
[1,2,3,4,2,3,5,4,6,5,7,6,5,4,2,3,8,7,6,5,4,2,3,1],\ [1,2,3,4,2,5,4,2,6,7,8,7,6,5,4,2,3,1],\\
[1,2,3,4,2,5,4,6,5,4,2,7,6,5,4,2,3,1],\ [1,2,3,4,2,5,4,6,5,4,2,7,8,7,6,5,4,2,3,1],\\
[1,2,3,4,3,5,4,6,5,4,3,7,6,5,4,2,3,1],\ [1,2,3,4,3,5,4,6,5,7,6,5,4,3,8,7,6,5,4,2,3,1],\\
[1,2,3,4,5,4,2,3,4,6,5,7,6,5,4,2,3,4,8,7,6,5,4,2,3,1],\ [1,2,3,4,5,4,6,5,4,7,8,7,6,5,4,2,3,1],\\
[1,2,3,4,5,4,6,7,6,5,4,2,3,1],\ [1,2,3,4,5,4,6,7,8,7,6,5,4,2,3,1],\\
[1,2,3,4,5,6,5,4,2,3,4,5,7,6,5,4,2,3,4,5,8,7,6,5,4,2,3,1],\ [1,2,3,4,5,6,5,7,8,7,6,5,4,2,3,1],\\
[1,2,4,2,3,1,7,6,5,4,2,3,1,4,3,5,4,2,6,7],\ [1,2,4,2,3,1,8,7,6,5,4,2,3,1,4,3,5,4,2,6,7,8],\\
[1,2,4,2,3,4,5,4,3,6,7,6,5,4,2,3,1,4],\ [1,2,4,2,3,4,5,4,3,6,7,8,7,6,5,4,2,3,1,4],\\
[1,2,4,2,3,4,5,4,6,5,4,3,7,6,5,4,2,3,1,4],\ [1,2,4,2,3,4,5,4,6,5,7,6,5,4,3,8,7,6,5,4,2,3,1,4],\\
[1,2,4,5,4,2,3,4,5,6,5,4,3,7,6,5,4,2,3,1,4,5],\ [1,2,4,5,4,2],\ [1,2,4,5,6,5,4,2],\\
[1,2,6,5,4,2,3,1,7,6,5,4,2,3,1,4,3,5,4,2,6,5,7,6],\ [1,3,1,4,2,3,4,5,4,6,5,7,6,5,4,2,8,7,6,5,4,2,3,1,4,3],\\
[1,3,1,4,2,5,4,2,3,1,8,7,6,5,4,2,3,1,4,3,5,4,2,6,5,4,3,1,7,8],\\
[1,3,1,4,2,7,6,5,4,2,3,1,8,7,6,5,4,2,3,1,4,3,5,4,2,6,5,4,3,1,7,6,8,7],\\
[1,3,1,4,3,1,5,6,7,8,7,6,5,4,3,1],\ [1,3,1,4,3,5,4,3,1,6,5,4,3,1],\ [1,3,1,4,3,5,4,3,1,6,7,8,7,6,5,4,3,1],\\
[1,3,1,4,5,4,2,3,4,5,6,5,4,2,7,8,7,6,5,4,2,3,1,4,3,5],\\
[1,3,1,4,5,4,2,3,4,5,6,5,7,6,5,4,2,8,7,6,5,4,2,3,1,4,3,5],\\
[1,3,1,4,5,6,5,4,2,3,4,5,6,7,6,5,4,2,8,7,6,5,4,2,3,1,4,3,5,6],\\
[1,3,1,4,5,6,5,4,3,1],\ [1,3,1,4,5,6,7,6,5,4,3,1],\ [1,3,4,2,3,4,5,4,2,6,7,8,7,6,5,4,2,3,1,4],\\
[1,3,4,2,3,4,5,4,6,5,7,6,5,4,2,8,7,6,5,4,2,3,1,4],\ [1,3,4,3,5,4,3,1],\\
[1,3,4,3,5,4,3,6,5,4,3,1],\ [1,3,4,5,4,2,3,4,5,6,5,4,2,7,6,5,4,2,3,1,4,5],
\end{gathered}$$

$$\begin{gathered}
[1,3,4,5,4,2,3,4,5,6,5,4,2,7,8,7,6,5,4,2,3,1,4,5],\\
[1,3,4,5,4,2,3,4,5,6,5,7,6,5,4,2,8,7,6,5,4,2,3,1,4,5],\ [1,3,4,5,4,3,1,7],\\
[1,3,4,5,4,6,5,4,7,6,5,4,3,1],\ [1,3,4,5,4,6,7,6,5,4,3,1],\ [1,3,4,5,4,6,7,8,7,6,5,4,3,1],\ [1,3,4,5,6,5,4,3,1,8],\\
[1,3,4,5,6,5,7,6,5,4,3,1],\ [1,3,4,5,6,7,6,8,7,6,5,4,3,1],\ [1,4,2,3,1,4,3,5,4,3,1,6,7,8,7,6,5,4,2,3,1,4],\\
[1,4,2,3,1,4,3,5,4,6,5,7,6,5,4,3,1,8,7,6,5,4,2,3,1,4],\ [1,4,2,3,1,4,5,6,7,8,7,6,5,4,2,3,1,4],\\
[1,4,2,3,4,5,4,2,3,4,6,5,7,6,5,4,2,3,4,8,7,6,5,4,2,3,1,4],\ [1,4,2,3,4,5,6,5,7,6,5,4,2,3,1,4],\\
[1,4,2,3,4,5,6,5,7,6,5,8,7,6,5,4,2,3,1,4],\ [1,4,5,4,2,3,4,5,6,5,4,2,3,1,4,5],\\
[1,4,5,4,2,3,4,5,6,7,6,5,4,2,3,1,4,5],\ [1,4,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,5],\\
[1,4],\ [1,5,4,2,3,1,4,5,6,5,4,2,3,1,4,5],\ [1,5,4,2,3,1,4,5,6,7,8,7,6,5,4,2,3,1,4,5],\\
[1,5,4,2,3,1,4,5],\ [1,5,4,2,3,4,5,6,5,4,2,3,1,4,5,8],\ [1,5,4,2,3,4,5,6,7,6,8,7,6,5,4,2,3,1,4,5],\\
[1,5,6,5,4,2,3,4,5,6,7,6,5,4,2,3,1,4,5,6],\ [1,6,5,4,2,3,1,4,5,6],\ [1,6],\ [1,7,6,5,4,2,3,1,4,5,6,7],\ [1,8],\\
[2,3,1,4,2,3,4,5,4,6,5,7,6,5,4,3,8,7,6,5,4,2,3,1,4,3],\\
[2,3,1,4,5,6,5,4,2,3,4,5,6,7,6,5,4,3,8,7,6,5,4,2,3,1,4,3,5,6],\ [2,3,1,6,5,4,2,3,1,4,3,5,6,7,6,5,4,2,3,1,4,3,5,6],\\
[2,3,4,2,3,1,4,7,6,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,4,2,6,7],\\
[2,3,4,2,3,5,4,2,3,6,7,8,7,6,5,4,2,3],\ [2,3,4,2,3,5,4,6,5,4,2,3,7,6,5,4,2,3],\\
[2,3,4,2,3,5,4,6,5,4,2,3,7,8,7,6,5,4,2,3],\ [2,3,4,2,3,5,4,6,5,7,6,5,4,2,3,8,7,6,5,4,2,3],\\
[2,3,4,2,5,4,2,6,5,4,2,3],\ [2,3,4,2,5,4,2,6,7,8,7,6,5,4,2,3],\ [2,3,4,2,5,4,6,5,4,2,7,6,5,4,2,3],\\
[2,3,4,2,5,4,6,5,4,2,7,8,7,6,5,4,2,3],\ [2,3,4,2,5,6,5,4,2,3],\ [2,3,4,2,5,6,7,6,5,4,2,3],\\
[2,3,4,3,5,4,6,5,4,3,7,6,5,4,2,3],\ [2,3,4,3,5,6,7,6,5,4,2,3],\ [2,3,4,3,5,6,7,8,7,6,5,4,2,3],\\
[2,3,4,5,4,6,5,4,2,3],\ [2,3,4,5,4,6,5,4,7,6,5,4,2,3],\ [2,3,4,5,6,5,4,2,3,8],\ [2,3,4,5,6,5,7,6,5,8,7,6,5,4,2,3],\\
[2,3,4,5,6,5,7,8,7,6,5,4,2,3],\ [2,3,4,5,6,7,6,8,7,6,5,4,2,3],\ [2,4,2,3,4,5,4,3,6,5,4,2,3,4],\\
[2,4,2,3,4,5,4,3,6,7,6,5,4,2,3,4],\ [2,4,2,3,4,5,4,3,6,7,8,7,6,5,4,2,3,4],\ [2,4,2,3,4,5,4,6,5,4,3,7,6,5,4,2,3,4],\\
[2,4,2,5,4,2,6,5,4,2],\ [2,4,2,5,6,5,4,2],\ [2,4,2,5,6,7,6,5,4,2],\ [2,4,2,6],\ [2,4,2,8],\\
[2,4,3,1,6,7,6,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,4,2,6,7],\ [2,4,5,4,2,7,8,7],\\
[2,4,5,4,2,7],\ [2,4,5,4,2,8],\ [2,4,5,4,6,5,4,2],\ [2,4,5,6,5,7,6,5,4,2],\ [3,1,4,2,3,4,5,6,5,7,8,7,6,5,4,2,3,1,4,3],\\
[3,1,4,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5],\ [3,1,4,5,4,6,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,6],\\
[3,1,4,6,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,6],\ [3,1,4,7,6,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,6,7],\\
[3,1,5,4,2,3,4,5,6,7,6,8,7,6,5,4,2,3,1,4,3,5],\ [3,1,5,6,5,4,2,3,4,5,6,7,6,5,4,2,3,1,4,3,5,6],\\
[3,1,5,6,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,6],\ [3,1,6,5,4,2,3,1,4,3,5,6],\\
[3,1,6,7,6,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,6,7],\ [3,1,7,6,5,4,2,3,1,4,3,5,6,7],\\
[3,1,8,7,6,5,4,2,3,1,4,3,5,6,7,8],\ [3,4,2,3,1,5,4,2,3,1,4,8,7,6,5,4,2,3,1,4,3,5,4,2,6,5,4,3,7,8],\\
[3,4,2,3,4,5,4,2,6,5,4,2,3,4],\ [3,4,2,3,4,5,4,2,6,7,8,7,6,5,4,2,3,4],\ [3,4,3,5,4,3],\ [3,4,3,6,7,6],\\
[3,4,3,6],\ [3,4,3,7],\ [3,4,5,4,3,7,8,7],\ [3,4,5,4,6,5,4,3],\ [3,4,5,6,5,4,3,8],\ [3,5],\ [3,7],\\
[4,2,3,1,5,4,2,3,1,4,3,5,6,7,8,7,6,5,4,2,3,1,4,3,5,4],\ [4,2,3,4,5,4,2,3,4,7],\ [4,2,3,4,5,6,5,4,2,3,4,8],\\
[4,2,3,4,5,6,5,7,6,5,4,2,3,4],\ [4,2,3,4,5,6,7,6,8,7,6,5,4,2,3,4],\ [4,2,3,4],\\
[4,3,1,5,4,2,3,4,5,6,7,6,8,7,6,5,4,2,3,1,4,3,5,4],\ [4,5,4,2,3,4,5,6,5,4,2,3,4,5],\ [4,5,4,8],\ [4,7,8,7],\\
[4,7],\ [4,8],\ [5,4,2,3,4,5,6,5,4,2,3,4,5,8],\ [5,4,2,3,4,5],\\
[5,4,3,1,6,5,4,2,3,1,4,3,5,4,6,5],\ [5,8],\ [6,5,4,2,3,4,5,6]\big\},
\end{gathered}$$

one obtains 100 solutions for w_i in (5.233) with $\ell_T(w_i) = 2$ and w_i of type A_2:

$$\begin{gathered}
w_i \in \big\{[1,2,3,1,4,5,4,6,7,8,7,6,5,4,2,3,1,4,3,5],\ [1,2,3,1,4,5,6,5,4,7,6,5,4,2,3,1,4,3,5,6],\\
[1,2,3,1,4,5,6,5,4,7,8,7,6,5,4,2,3,1,4,3,5,6],\ [1,2,3,1,4,5,6,7,8,7,6,5,4,2,3,1,4,3],\\
[1,2,3,4,3,1,5,6,7,6,5,4,3,8,7,6,5,4,2,3,1,4,3,5,4,2,6,7],\ [1,2,3,4,5,4,6,7,6,5,4,2,3,1,4,5],\\
[1,2,3,4,5,4,6,7,8,7,6,5,4,2,3,1,4,5],\ [1,2,3,4,5,6,7,6,5,4,2,3,1,4],\\
[1,2,3,4,5,6,7,6,5,4,2,3,1,8],\ [1,2,3,4,5,6,7,6,5,4,2,3],\ [1,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4],\\
[1,2,3,4,5,6,7,8,7,6,5,4,2,3],\ [1,2,4,2,3,1,4,5,6,7,6,5,4,2,3,1,4,3,5,6,7,8,7,6,5,4,2,3],\\
[1,2,4,2,3,4,5,6,7,6,5,4,3,1],\ [1,2,4,2,3,4,5,6,7,8,7,6,5,4,3,1],\ [1,2,4,5,4,2,3,4,5,6,5,4,3,1],
\end{gathered}$$

[1, 2, 4, 5, 4, 2, 3, 4, 5, 6, 7, 6, 5, 4, 3, 1], [1, 2, 4, 5, 4, 2, 3, 4, 5, 6, 7, 8, 7, 6, 5, 4, 3, 1], [1, 2, 4, 5, 6, 5, 4, 2, 3, 4, 5, 6, 7, 6, 5, 4, 3, 1],
[1, 3, 1, 4, 5, 4, 2, 3, 1, 4, 5, 6, 7, 8, 7, 6, 5, 4, 2, 3], [1, 3, 1, 4, 5, 6, 5, 4, 2, 3, 1, 4, 5, 6, 7, 6, 5, 4, 2, 3],
[1, 3, 1, 4, 5, 6, 5, 4, 2, 3, 1, 4, 5, 6, 7, 8, 7, 6, 5, 4, 2, 3], [1, 3, 1, 4, 5, 6, 7, 6, 5, 4, 2, 3, 1, 4, 5, 6, 7, 8, 7, 6, 5, 4, 2, 3],
[1, 3, 4, 3, 1, 5, 4, 2, 3, 1, 4, 5, 6, 7, 8, 7, 6, 5, 4, 2, 3, 4], [1, 3, 4, 5, 4, 2, 3, 1, 4, 5, 6, 5, 4, 2],
[1, 3, 4, 5, 4, 2, 3, 1, 4, 5, 6, 7, 6, 5, 4, 2], [1, 3, 4, 5, 4, 3, 1, 6, 7, 6], [1, 3, 4, 5, 4, 3, 1, 6], [1, 3, 4, 5, 4, 3],
[1, 3, 4, 5, 6, 5, 4, 2, 3, 1, 4, 5, 6, 7, 6, 5, 4, 2], [1, 3, 4, 5, 6, 5, 4, 3, 1, 7, 8, 7], [1, 3, 4, 5, 6, 5, 4, 3, 1, 7],
[1, 3, 4, 5, 6, 5, 4, 3], [1, 3, 4, 5, 6, 7, 6, 5, 4, 3, 1, 8], [1, 4, 2, 3, 1, 4, 5, 4, 6, 7, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 4],
[1, 4, 2, 3, 4, 5, 6, 5, 7, 6, 5, 4, 2, 3, 1, 4, 5, 6], [1, 4, 2, 3, 4, 5, 6, 7, 6, 5, 4, 2, 3, 1, 4, 5], [1, 4, 2, 3, 4, 5, 6, 7, 6, 5, 4, 2, 3, 1, 4, 8],
[1, 4, 2, 3, 4, 5, 6, 7, 6, 5, 4, 2, 3, 4], [1, 4, 2, 3, 4, 5, 6, 7, 8, 7, 6, 5, 4, 2, 3, 1, 4, 5], [1, 4, 2, 3, 4, 5, 6, 7, 8, 7, 6, 5, 4, 2, 3, 4],
[1, 5, 4, 2, 3, 4, 5, 6, 5, 4, 2, 3, 1, 4, 5, 7, 8, 7], [1, 5, 4, 2, 3, 4, 5, 6, 5, 4, 2, 3, 1, 4, 5, 7], [1, 5, 4, 2, 3, 4, 5, 6, 5, 4, 2, 3, 4, 5],
[1, 5, 4, 2, 3, 4, 5, 6, 7, 6, 5, 4, 2, 3, 1, 4, 5, 6], [1, 5, 4, 2, 3, 4, 5, 6, 7, 6, 5, 4, 2, 3, 1, 4, 5, 8],
[2, 3, 1, 4, 2, 3, 1, 4, 5, 6, 7, 6, 5, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 4, 2, 6, 7], [2, 3, 1, 4, 5, 6, 5, 4, 2, 3, 1, 4, 5, 6, 7, 6, 5, 4, 3, 1],
[2, 3, 1, 4, 5, 6, 5, 4, 2, 3, 1, 4, 5, 6, 7, 8, 7, 6, 5, 4, 3, 1], [2, 3, 1, 4, 5, 6, 7, 6, 5, 4, 2, 3, 1, 4, 5, 6, 7, 8, 7, 6, 5, 4, 3, 1],
[2, 3, 4, 3, 1, 5, 6, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 6, 7, 8, 7, 6, 5, 4, 2, 3, 1, 4], [2, 3, 4, 5, 4, 6, 5, 4, 2, 3, 4, 5],
[2, 3, 4, 5, 6, 5, 4, 2, 3, 4], [2, 3, 4, 5, 6, 5, 4, 2, 3, 7, 8, 7], [2, 3, 4, 5, 6, 5, 4, 2, 3, 7],
[2, 3, 4, 5, 6, 5, 4, 2], [2, 3, 4, 5, 6, 7, 6, 5, 4, 2, 3, 4], [2, 3, 4, 5, 6, 7, 6, 5, 4, 2, 3, 8], [2, 3, 4, 5, 6, 7, 6, 5, 4, 2],
[2, 3, 4, 5, 6, 7, 8, 7, 6, 5, 4, 2, 3, 4], [2, 4, 2, 3, 1, 7, 6, 5, 4, 2, 3, 4, 5, 6, 7, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 6, 7],
[2, 4, 2, 3, 4, 5, 4, 3], [2, 4, 2, 3, 4, 5, 6, 5, 4, 3], [2, 4, 2, 5, 6, 5], [2, 4, 2, 5], [2, 4, 5, 4, 2, 3, 4, 5, 6, 5, 4, 3],
[2, 4, 5, 4, 2, 6, 7, 6], [2, 4, 5, 4, 2, 6], [2, 4, 5, 6, 5, 4, 2, 7], [3, 1, 4, 2, 3, 4, 5, 6, 5, 7, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 6],
[3, 1, 4, 2, 3, 4, 5, 6, 7, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5], [3, 1, 4, 2, 3, 4, 5, 6, 7, 8, 7, 6, 5, 4, 2, 3, 1, 4],
[3, 1, 5, 4, 2, 3, 4, 5, 6, 7, 6, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 6, 7], [3, 1, 5, 4, 2, 3, 4, 5, 6, 7, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 4],
[3, 1, 5, 4, 2, 3, 4, 5, 6, 7, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 6], [3, 1, 5, 4, 2, 3, 4, 5, 6, 7, 8, 7, 6, 5, 4, 2, 3, 1, 4, 5],
[3, 1, 6, 5, 4, 2, 3, 4, 5, 6, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 6, 8], [3, 1, 6, 5, 4, 2, 3, 4, 5, 6, 7, 6, 5, 4, 2, 3, 1, 4, 5, 6],
[3, 1, 6, 5, 4, 2, 3, 4, 5, 6, 7, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 6, 7], [3, 4, 2, 3, 4, 5, 4, 2], [3, 4, 2, 3, 4, 5, 6, 5, 4, 2],
[3, 4, 2, 3, 4, 5, 6, 7, 6, 5, 4, 2], [3, 4, 3, 1, 5, 4, 2, 3, 4, 5, 6, 7, 8, 7, 6, 5, 4, 2, 3, 1, 4, 5], [3, 4, 3, 5, 6, 5],
[3, 4, 3, 5], [3, 4, 5, 4, 2, 3, 4, 5, 6, 5, 4, 2], [3, 4, 5, 4, 3, 6], [3, 4, 5, 4], [3, 4],
[4, 2, 3, 1, 4, 5, 4, 2, 3, 1, 4, 3, 5, 6, 7, 8, 7, 6, 5, 4, 3, 1], [4, 2, 3, 4, 5, 4, 2, 3, 4, 6, 7, 6], [4, 2, 3, 4, 5, 4, 2, 3, 4, 6],
[4, 2, 3, 4, 5, 6, 5, 4, 2, 3, 4, 5], [4, 2, 3, 4, 5, 6, 5, 4, 2, 3, 4, 7, 8, 7], [4, 2, 3, 4, 5, 6, 5, 4, 2, 3, 4, 7],
[4, 2, 3, 4, 5, 6, 7, 6, 5, 4, 2, 3, 4, 8], [4, 5], [5, 6], [6, 7], [7, 8]},

one obtains 75 solutions for w_i in (5.233) with $\ell_T(w_i) = 3$ and w_i of type A_1^3:

$w_i \in \{$[1, 2, 3, 1, 4, 2, 5, 4, 6, 5, 4, 2, 7, 8, 7, 6, 5, 4, 2, 3, 1], [1, 2, 3, 1, 4, 3, 5, 4, 3, 6, 5, 4, 3, 7, 6, 5, 4, 3, 1, 8, 7, 6, 5, 4, 2, 3, 1],
[1, 2, 3, 1, 4, 5, 4, 6, 5, 4, 2, 3, 4, 5, 7, 6, 5, 4, 2, 3, 1, 4, 5, 8, 7, 6, 5, 4, 2, 3, 1], [1, 2, 3, 1, 4, 5, 4, 6, 7, 8, 7, 6, 5, 4, 2, 3, 1],
[1, 2, 3, 4, 2, 3, 1, 6, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 4, 2, 6, 7, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 4, 2, 6, 7],
[1, 2, 3, 4, 2, 3, 5, 4, 6, 5, 4, 7, 6, 5, 4, 2, 3, 8, 7, 6, 5, 4, 2, 3, 1],
[1, 2, 3, 4, 2, 5, 4, 2, 3, 4, 6, 5, 4, 3, 7, 6, 5, 4, 2, 3, 4, 8, 7, 6, 5, 4, 2, 3, 1],
[1, 2, 3, 4, 3, 1, 5, 6, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 6, 8, 7, 6, 5, 4, 2, 3, 1],
[1, 2, 3, 4, 3, 5, 4, 3, 6, 5, 4, 3, 7, 6, 5, 4, 2, 3, 1], [1, 2, 3, 4, 3, 5, 4, 6, 5, 4, 7, 6, 5, 4, 3, 8, 7, 6, 5, 4, 2, 3, 1],
[1, 2, 3, 4, 5, 4, 6, 5, 4, 2, 3, 4, 5, 7, 6, 5, 4, 2, 3, 4, 5, 8, 7, 6, 5, 4, 2, 3, 1], [1, 2, 3, 4, 5, 6, 5, 4, 2, 3, 4, 5, 7, 6, 5, 4, 2, 3, 1],
[1, 2, 3, 4, 5, 6, 5, 4, 3, 1, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 4, 6, 5, 8, 7, 6, 5, 4, 2, 3, 1],
[1, 2, 4, 2, 3, 1, 4, 3, 5, 4, 3, 1, 6, 5, 7, 6, 5, 4, 3, 1, 8, 7, 6, 5, 4, 2, 3, 1, 4],
[1, 2, 4, 2, 3, 1, 5, 6, 5, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 4, 2, 6, 7], [1, 2, 4, 2, 3, 1, 6, 7, 6, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 4, 2, 6, 7, 8],
[1, 2, 4, 2, 5, 4, 2, 3, 4, 5, 6, 5, 4, 3, 7, 6, 5, 4, 2, 3, 1, 4, 5], [1, 2, 4, 2, 5, 6, 5, 4, 2], [1, 2, 4, 5, 4, 2, 8],
[1, 3, 1, 4, 2, 3, 4, 5, 4, 6, 5, 4, 7, 6, 5, 4, 2, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3],
[1, 3, 1, 4, 2, 3, 5, 4, 2, 3, 1, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 4, 2, 6, 5, 4, 3, 1, 7, 8],
[1, 3, 1, 4, 2, 3, 7, 6, 5, 4, 2, 3, 1, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 4, 2, 6, 5, 4, 3, 1, 7, 6, 8, 7],
[1, 3, 1, 4, 2, 5, 4, 2, 3, 1, 6, 7, 6, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 4, 2, 6, 5, 4, 3, 1, 7, 8],

$$\begin{gathered}
[1,3,1,4,3,1,5,6,7,6,8,7,6,5,4,3,1],\ [1,3,1,4,5,4,6,5,4,2,3,4,5,6,7,6,5,4,2,8,7,6,5,4,2,3,1,4,3,5,6],\\
[1,3,1,4,5,4,6,7,6,5,4,3,1],\ [1,3,4,3,5,4,3,1,7],\ [1,3,4,3,5,4,3,6,5,4,3,1,8],\\
[1,3,4,5,4,2,3,1,4,5,6,5,4,2,7,8,7,6,5,4,2,3,1,4,5],\ [1,4,2,3,1,4,5,6,5,7,6,5,8,7,6,5,4,2,3,1,4],\\
[1,4,2,3,4,5,4,2,3,4,6,7,8,7,6,5,4,2,3,1,4],\ [1,4,2,3,4,5,6,7,6,5,4,2,3,4,5,6,8,7,6,5,4,2,3,1,4],\\
[1,4,5,4,2,3,1,4,5,6,5,4,2,3,1,4,5],\ [1,4,5,4,2,3,1,4,5,6,7,8,7,6,5,4,2,3,1,4,5],\\
[1,4,5,4,2,3,4,5,6,5,4,2,3,1,4,5,8],\ [1,4,5,4,2,3,4,5,6,7,6,8,7,6,5,4,2,3,1,4,5],\\
[1,4,5,4,6,5,4,2,3,1,4,5,6],\ [1,4,8],\ [1,5,4,2,3,1,4,5,7],\\
[1,5,4,2,3,4,5,6,5,4,2,3,4,5,7,6,5,4,2,3,1,4,5],\ [1,6,7,6,5,4,2,3,1,4,5,6,7],\\
[2,3,1,4,2,5,6,5,4,2,3,4,5,6,7,6,5,4,3,8,7,6,5,4,2,3,1,4,3,5,6],\\
[2,3,1,5,6,5,4,2,3,1,4,3,5,6,7,6,5,4,2,3,1,4,3,5,6],\\
[2,3,4,2,3,1,4,6,7,6,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,4,2,6,7],\ [2,3,4,2,3,5,4,2,3,6,7,6,8,7,6,5,4,2,3],\\
[2,3,4,2,3,5,4,3,6,5,4,2,3,7,8,7,6,5,4,2,3],\ [2,3,4,2,3,5,4,6,5,4,7,6,5,4,2,3,8,7,6,5,4,2,3],\\
[2,3,4,2,3,5,6,7,6,5,4,2,3],\ [2,3,4,3,5,6,5,7,8,7,6,5,4,2,3],\ [2,3,4,5,4,6,5,4,2,3,8],\\
[2,3,4,5,6,5,4,2,3,4,5,7,6,5,4,2,3],\ [2,4,2,3,1,4,5,4,3,8,7,6,5,4,2,3,1,4,3,5,4,2,6,7,8],\\
[2,4,2,3,1,4,5,6,7,6,5,4,3,8,7,6,5,4,2,3,1,4,3,5,4,2,6,7,8],\ [2,4,2,3,4,5,4,3,6,5,4,2,3,4,8],\\
[2,4,2,3,4,5,4,3,6,7,6,8,7,6,5,4,2,3,4],\ [3,1,4,2,3,4,5,6,7,6,5,4,2,3,4,5,6,8,7,6,5,4,2,3,1,4,3],\\
[3,1,4,5,4,2,3,4,5,6,7,6,8,7,6,5,4,2,3,1,4,3,5],\ [3,1,4,5,4,6,5,4,2,3,1,4,3,5,6],\\
[3,1,4,5,4,8,7,6,5,4,2,3,1,4,3,5,6,7,8],\ [3,1,4,6,7,6,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,6,7],\\
[3,1,5,6,5,7,6,5,4,2,3,1,4,3,5,6,7],\ [3,1,6,5,4,2,3,1,4,3,5,6,8],\\
[3,1,6,5,4,2,3,4,5,6,7,6,5,4,2,3,4,5,6,8,7,6,5,4,2,3,1,4,3,5,6],\ [3,1,7,8,7,6,5,4,2,3,1,4,3,5,6,7,8],\\
[3,4,2,3,1,5,4,2,3,1,4,7,8,7,6,5,4,2,3,1,4,3,5,4,2,6,5,4,3,7,8],\ [3,4,3,5,4,3,7,8,7],\\
[3,4,5,4,6,5,4,3,8],\ [4,2,3,1,5,4,2,3,1,4,3,5,6,7,6,8,7,6,5,4,2,3,1,4,3,5,4],\\
[4,2,3,4,5,4,2,3,4,6,5,4,2,3,4],\ [4,2,3,4,5,4,2,3,4,6,7,8,7,6,5,4,2,3,4],\ [4,2,3,4,6],\\
[4,5,4,2,3,4,5,6,5,4,2,3,4,5,8],\ [5,4,2,3,4,5,7,8,7],\ [5,4,3,1,6,5,4,2,3,1,4,3,5,4,6,5,8],\ [5,6,5,4,2,3,4,5,6]\big\},
\end{gathered}$$

one obtains 165 solutions for w_i in (5.233) with $\ell_T(w_i) = 3$ and w_i of type $A_1 * A_2$:

$$\begin{gathered}
w_i \in \big\{[1,3,1,4,5,4,2,3,4,5,6,5,4,2,7,8,7,6,5,4,2,3,1,4,5],\\
[1,2,3,1,4,2,5,4,2,6,5,4,2,7,8,7,6,5,4,2,3,1,4,3,5],\\
[1,2,3,1,4,2,5,4,6,5,4,2,3,4,5,7,6,5,4,3,1,8,7,6,5,4,2,3,1],\\
[1,2,3,1,4,3,5,4,6,5,4,3,7,6,5,4,3,8,7,6,5,4,2,3,1,4,3,5,6],\\
[1,2,3,1,4,3,5,4,6,5,7,6,5,4,3,8,7,6,5,4,2,3,1,4,3],\ [1,2,3,1,4,3,5,4,6,5,7,6,5,4,3,8,7,6,5,4,2,3,1],\\
[1,2,3,1,4,5,4,6,5,4,7,6,5,4,2,3,1,4,3,5,6],\ [1,2,3,1,4,5,4,6,5,4,7,8,7,6,5,4,2,3,1,4,3,5,6],\\
[1,2,3,1,4,5,6,5,4,2,3,4,5,7,6,5,4,2,3,4,5,8,7,6,5,4,2,3,1],\ [1,2,3,1,4,5,6,5,7,8,7,6,5,4,2,3,1,4,3],\\
[1,2,3,4,2,3,5,4,6,5,4,7,6,5,4,2,3,4,5,8,7,6,5,4,2,3,1],\ [1,2,3,4,2,3,5,4,6,5,7,6,5,4,2,3,4,8,7,6,5,4,2,3,1],\\
[1,2,3,4,2,5,4,2,3,4,6,5,7,6,5,4,3,8,7,6,5,4,2,3,1],\ [1,2,3,4,2,5,4,2,6,5,4,2,7,6,5,4,2,3,1,4,5],\\
[1,2,3,4,2,5,4,2,6,5,4,2,7,8,7,6,5,4,2,3,1,4,5],\ [1,2,3,4,2,5,4,2,6,5,7,8,7,6,5,4,2,3,1],\\
[1,2,3,4,2,5,4,2,6,7,8,7,6,5,4,2,3,1,4],\ [1,2,3,4,2,5,4,6,5,4,2,3,4,5,7,6,5,4,3,8,7,6,5,4,2,3,1],\\
[1,2,3,4,2,5,4,6,5,4,2,7,6,5,4,2,3,1,8],\ [1,2,3,4,3,1,5,6,5,7,6,5,4,3,8,7,6,5,4,2,3,1,4,3,5,4,2,6,7],\\
[1,2,3,4,3,5,4,3,6,5,4,3,7,6,5,4,2,3,1,4,5],\ [1,2,3,4,3,5,4,6,5,4,3,7,6,5,4,2,3,1,4],\\
[1,2,3,4,3,5,4,6,5,4,3,7,6,5,4,2,3],\ [1,2,3,4,3,5,4,6,5,7,6,5,4,3,8,7,6,5,4,2,3,1,4],\\
[1,2,3,4,5,4,2,3,4,6,5,7,6,5,4,2,3,4,5,8,7,6,5,4,2,3,1],\\
[1,2,3,4,5,4,2,3,4,6,5,7,6,5,4,2,3,4,8,7,6,5,4,2,3,1,4],\\
[1,2,3,4,5,4,6,5,7,8,7,6,5,4,2,3,1],\ [1,2,3,4,5,4,6,7,6,5,4,2,3,1,8],\\
[1,2,3,4,5,6,5,4,3,1,7,6,5,4,2,3,1,4,3,5,6],\ [1,2,3,4,5,6,5,7,8,7,6,5,4,2,3],\\
[1,2,4,2,3,1,4,3,8,7,6,5,4,2,3,1,4,3,5,4,2,6,5,4,3,7,8],\\
[1,2,4,2,3,1,4,5,6,5,7,6,5,4,2,3,1,4,3,5,6,7,8,7,6,5,4,2,3],\\
[1,2,4,2,3,1,6,7,6,5,4,2,3,1,4,3,5,4,2,6,5,7,6],\ [1,2,4,2,3,4,5,4,2,3,4,6,5,7,6,5,4,2,8,7,6,5,4,2,3,1,4],
\end{gathered}$$

[1, 2, 4, 2, 3, 4, 5, 4, 3, 6, 5, 7, 6, 5, 4, 2, 3, 1, 4], [1, 2, 4, 2, 3, 4, 5, 4, 3, 6, 5, 7, 6, 5, 8, 7, 6, 5, 4, 2, 3, 1, 4],
[1, 2, 4, 2, 3, 4, 5, 4, 3, 6, 7, 6, 5, 4, 2, 3, 1, 4, 8], [1, 2, 4, 2, 3, 4, 5, 4, 3, 6, 7, 6, 5, 4, 2, 3, 4],
[1, 2, 4, 2, 3, 4, 5, 4, 3, 6, 7, 8, 7, 6, 5, 4, 2, 3, 4], [1, 2, 4, 2, 3, 4, 5, 4, 6, 5, 4, 3, 7, 6, 5, 4, 2, 3, 1, 4, 5],
[1, 2, 4, 2, 3, 4, 5, 4, 6, 5, 4, 3, 7, 6, 5, 4, 2, 3, 4], [1, 2, 4, 2, 3, 4, 5, 6, 5, 7, 6, 5, 4, 3, 1], [1, 2, 4, 5, 4, 2, 3, 4, 5, 6, 5, 4, 3, 1, 8],
[1, 2, 4, 5, 4, 2, 3, 4, 5, 6, 7, 6, 8, 7, 6, 5, 4, 3, 1], [1, 2, 4, 5, 4, 2, 6], [1, 2, 4, 5, 4, 6, 5, 4, 2, 3, 4, 5, 6, 7, 6, 5, 4, 3, 1],
[1, 3, 1, 4, 2, 3, 4, 5, 4, 6, 5, 4, 7, 6, 5, 4, 2, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 6],
[1, 3, 1, 4, 2, 3, 4, 5, 4, 6, 5, 7, 6, 5, 4, 2, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5],
[1, 3, 1, 4, 2, 3, 4, 5, 4, 6, 5, 7, 6, 5, 4, 2, 8, 7, 6, 5, 4, 2, 3, 1, 4],
[1, 3, 1, 4, 2, 3, 4, 5, 6, 7, 6, 5, 4, 2, 3, 1, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 4, 2, 6, 7],
[1, 3, 1, 4, 2, 5, 4, 2, 3, 1, 7, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 4, 2, 6, 5, 4, 3, 1, 7, 6, 8, 7], [1, 3, 1, 4, 3, 1, 5, 4, 6, 7, 8, 7, 6, 5, 4, 3, 1],
[1, 3, 1, 4, 3, 5, 4, 3, 1, 6, 5, 4, 3, 1, 7, 8, 7], [1, 3, 1, 4, 3, 5, 4, 3, 6, 5, 4, 2, 3, 1, 4, 5, 6, 7, 8, 7, 6, 5, 4, 2, 3],
[1, 3, 1, 4, 3, 5, 4, 3, 6, 5, 4, 3, 1], [1, 3, 1, 4, 5, 4, 2, 3, 1, 4, 5, 6, 7, 6, 8, 7, 6, 5, 4, 2, 3],
[1, 3, 1, 4, 5, 4, 2, 3, 4, 5, 6, 5, 4, 2, 7, 6, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5],
[1, 3, 1, 4, 5, 4, 2, 3, 4, 5, 6, 5, 7, 6, 5, 4, 2, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 6],
[1, 3, 1, 4, 5, 4, 2, 3, 4, 5, 6, 5, 7, 6, 5, 4, 2, 8, 7, 6, 5, 4, 2, 3, 1, 4, 5], [1, 3, 1, 4, 5, 6, 5, 4, 3, 1, 7],
[1, 3, 1, 6, 5, 4, 2, 3, 1, 4, 3, 5, 4, 6, 5, 7, 6, 5, 4, 2, 3], [1, 3, 4, 2, 3, 4, 5, 4, 2, 3, 4, 6, 5, 7, 6, 5, 4, 3, 8, 7, 6, 5, 4, 2, 3, 1, 4],
[1, 3, 4, 2, 3, 4, 5, 4, 2, 6, 5, 7, 6, 5, 8, 7, 6, 5, 4, 2, 3, 1, 4], [1, 3, 4, 2, 3, 4, 5, 4, 6, 5, 7, 6, 5, 4, 2, 8, 7, 6, 5, 4, 2, 3, 1, 4, 5],
[1, 3, 4, 2, 3, 5, 4, 2, 3, 1, 4, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 4, 2, 6, 7, 8],
[1, 3, 4, 2, 3, 5, 6, 7, 6, 5, 4, 2, 3, 1, 4, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 4, 2, 6, 7, 8],
[1, 3, 4, 3, 1, 5, 4, 2, 3, 1, 4, 5, 6, 7, 6, 8, 7, 6, 5, 4, 2, 3, 4], [1, 3, 4, 3, 5, 4, 2, 3, 1, 4, 5, 6, 5, 4, 2],
[1, 3, 4, 3, 5, 4, 2, 3, 1, 4, 5, 6, 7, 6, 5, 4, 2], [1, 3, 4, 3, 5, 4, 3], [1, 3, 4, 5, 4, 2, 3, 4, 5, 6, 5, 4, 2, 7, 6, 5, 4, 2, 3, 1, 4, 5, 8],
[1, 3, 4, 5, 4, 2, 3, 4, 5, 6, 5, 4, 2, 7, 6, 8, 7, 6, 5, 4, 2, 3, 1, 4, 5], [1, 3, 4, 5, 4, 6, 5, 7, 6, 5, 4, 3, 1],
[1, 3, 4, 5, 4, 6, 7, 6, 5, 4, 3, 1, 8], [1, 3, 4, 5, 6, 5, 4, 3, 8], [1, 4, 2, 3, 1, 4, 3, 5, 4, 3, 1, 6, 5, 7, 6, 5, 8, 7, 6, 5, 4, 2, 3, 1, 4],
[1, 4, 2, 3, 1, 4, 3, 5, 4, 3, 6, 7, 8, 7, 6, 5, 4, 2, 3, 1, 4], [1, 4, 2, 3, 1, 4, 3, 5, 4, 6, 5, 7, 6, 5, 4, 3, 8, 7, 6, 5, 4, 2, 3, 1, 4],
[1, 4, 2, 3, 1, 4, 3, 5, 6, 7, 8, 7, 6, 5, 4, 3, 1], [1, 4, 2, 3, 4, 5, 4, 3, 1, 6, 7, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 4],
[1, 4, 2, 3, 4, 5, 6, 5, 7, 6, 5, 4, 2, 3, 4], [1, 4, 3, 1, 5, 4, 2, 3, 1, 4, 3, 5, 4, 6, 7, 8, 7, 6, 5, 4, 2, 3, 4],
[1, 4, 5, 4, 2, 3, 4, 5, 6, 5, 4, 2, 3, 1, 4, 5, 7, 8, 7], [1, 4, 5, 4, 2, 3, 4, 5, 6, 5, 4, 2, 3, 1, 4, 5, 7],
[1, 4, 5, 4, 2, 3, 4, 5, 6, 5, 4, 2, 3, 4, 5], [1, 4, 5, 4, 2, 3, 4, 5, 6, 7, 6, 5, 4, 2, 3, 1, 4, 5, 8],
[1, 5, 4, 2, 3, 1, 4, 3, 5, 4, 6, 5, 4, 3, 1], [1, 5, 4, 2, 3, 1, 4, 3, 5, 4, 6, 7, 8, 7, 6, 5, 4, 3, 1], [1, 5, 4, 2, 3, 1, 4, 5, 6, 5, 4, 2, 3, 1, 4, 5, 7, 8, 7],
[1, 5, 4, 2, 3, 4, 5, 6, 5, 4, 2, 3, 4, 5, 8], [1, 5, 4, 2, 3, 4, 5], [1, 6, 5, 4, 2, 3, 4, 5, 6],
[2, 3, 1, 4, 2, 3, 1, 4, 5, 6, 5, 7, 6, 5, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 4, 2, 6, 7],
[2, 3, 1, 4, 2, 3, 4, 5, 4, 6, 5, 4, 7, 6, 5, 4, 3, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 6],
[2, 3, 1, 4, 2, 5, 4, 2, 6, 5, 4, 2, 3, 1, 4, 5, 6, 7, 8, 7, 6, 5, 4, 3, 1], [2, 3, 1, 4, 2, 5, 6, 5, 4, 2, 3, 1, 4, 5, 6, 7, 8, 7, 6, 5, 4, 3, 1],
[2, 3, 1, 4, 2, 5, 6, 7, 6, 5, 4, 2, 3, 1, 4, 5, 6, 7, 8, 7, 6, 5, 4, 3, 1],
[2, 3, 1, 4, 3, 1, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 4, 2, 6, 5, 7, 6, 8, 7, 6, 5, 4, 2, 3, 1, 4],
[2, 3, 1, 4, 5, 4, 6, 5, 4, 2, 3, 1, 4, 5, 6, 7, 6, 5, 4, 3, 1], [2, 3, 1, 4, 5, 4, 6, 5, 4, 2, 3, 1, 4, 5, 6, 7, 8, 7, 6, 5, 4, 3, 1],
[2, 3, 4, 2, 3, 1, 5, 6, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 6, 7, 8, 7, 6, 5, 4, 2, 3, 1, 4], [2, 3, 4, 2, 3, 5, 4, 2, 3, 6, 5, 7, 6, 5, 8, 7, 6, 5, 4, 2, 3],
[2, 3, 4, 2, 3, 5, 4, 2, 3, 6, 5, 7, 8, 7, 6, 5, 4, 2, 3], [2, 3, 4, 2, 3, 5, 4, 2, 6, 7, 8, 7, 6, 5, 4, 2, 3],
[2, 3, 4, 2, 3, 5, 4, 6, 5, 4, 2, 3, 7, 6, 5, 4, 2, 3, 8], [2, 3, 4, 2, 3, 5, 4, 6, 5, 4, 2, 3, 7, 6, 8, 7, 6, 5, 4, 2, 3],
[2, 3, 4, 2, 3, 5, 4, 6, 5, 4, 2, 7, 6, 5, 4, 2, 3], [2, 3, 4, 2, 3, 5, 4, 6, 5, 4, 2, 7, 8, 7, 6, 5, 4, 2, 3], [2, 3, 4, 2, 3, 5, 4, 6, 5, 4, 3, 7, 6, 5, 4, 2, 3],
[2, 3, 4, 2, 5, 4, 2, 6, 5, 4, 2, 3, 4], [2, 3, 4, 2, 5, 4, 2, 6, 5, 4, 2, 3, 7, 8, 7], [2, 3, 4, 2, 5, 4, 2, 6, 5, 7, 8, 7, 6, 5, 4, 2, 3],
[2, 3, 4, 2, 5, 4, 2, 6, 7, 8, 7, 6, 5, 4, 2, 3, 4], [2, 3, 4, 2, 5, 4, 6, 5, 4, 2, 3], [2, 3, 4, 2, 5, 4, 6, 5, 4, 2, 7, 6, 5, 4, 2, 3, 8],
[2, 3, 4, 2, 5, 4, 6, 5, 4, 7, 6, 5, 4, 2, 3], [2, 3, 4, 2, 5, 6, 5, 4, 2, 3, 7], [2, 3, 4, 3, 5, 4, 6, 5, 4, 3, 7, 6, 5, 4, 2, 3, 4],
[2, 3, 4, 3, 5, 4, 6, 5, 4, 7, 6, 5, 4, 2, 3], [2, 3, 4, 3, 5, 6, 7, 6, 5, 4, 2, 3, 8], [2, 3, 4, 5, 4, 6, 5, 4, 2, 3, 4, 5, 8],
[2, 3, 4, 5, 4, 6, 5, 4, 2], [2, 3, 4, 5, 6, 5, 4, 2, 3, 4, 8], [2, 3, 4, 5, 6, 5, 7, 6, 8, 7, 6, 5, 4, 2, 3],
[2, 3, 4, 5, 6, 7, 6, 8, 7, 6, 5, 4, 2, 3, 4], [2, 4, 2, 3, 1, 4, 7, 6, 5, 4, 2, 3, 4, 5, 6, 7, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 6, 7],
[2, 4, 2, 3, 1, 6, 7, 6, 5, 4, 2, 3, 4, 5, 6, 7, 8, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 6, 7], [2, 4, 2, 3, 1, 7, 6, 5, 4, 2, 3, 1, 4, 3, 5, 6, 7],

$$[2,4,2,3,1,8,7,6,5,4,2,3,1,4,3,5,6,7,8],\ [2,4,2,3,4,5,4,3,6,5,4,2,3,4,7,8,7],\ [2,4,2,3,4,5,4,3,6,5,4,2,3,4,7],$$
$$[2,4,2,3,4,5,4,3,6,5,7,6,5,4,2,3,4],\ [2,4,2,3,4,5,4,3,6,7,6,5,4,2,3,4,8],\ [2,4,2,3,4,5,6,5,4,3,8],$$
$$[2,4,2,5,4,6,5,4,2],\ [2,4,2,5,6,5,4,2,7],\ [2,4,2,5,8],\ [2,4,5,4,2,3,4,5,6,5,4,3,8],$$
$$[2,4,5,4,2,7,8],\ [2,4,5,6,5,4,2,3,1,7,6,5,4,2,3,1,4,3,5,6,7],\ [3,1,4,5,4,2,3,4,5,6,7,6,8,7,6,5,4,2,3,1,4,3,5,6,7],$$
$$[3,1,4,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,6],\ [3,1,4,5,6,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,6],$$
$$[3,1,4,6,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,6,7],\ [3,1,5,4,2,3,4,5,6,7,6,8,7,6,5,4,2,3,1,4,3,5,4],$$
$$[3,1,5,4,2,3,4,5,6,7,6,8,7,6,5,4,2,3,1,4,5],\ [3,1,5,6,5,4,2,3,1,4,3,5,4,6,5],$$
$$[3,1,5,6,5,4,2,3,4,5,6,7,6,5,4,2,3,1,4,3,5,6,8],\ [3,1,5,6,5,4,2,3,4,5,6,7,6,5,4,2,3,1,4,5,6],$$
$$[3,1,6,5,4,2,3,1,4,5,6],\ [3,1,7,6,5,4,2,3,1,4,5,6,7],$$
$$[3,4,2,3,4,5,4,2,3,1,4,8,7,6,5,4,2,3,1,4,3,5,6,7,8],\ [3,4,2,3,4,5,4,2,6,5,4,2,3,4,7,8,7],$$
$$[3,4,2,3,4,5,4,2,7],\ [3,4,2,3,4,5,6,5,7,6,5,4,2],\ [3,4,3,1,5,4,2,3,4,5,6,7,6,8,7,6,5,4,2,3,1,4,5],$$
$$[3,4,3,5,4,2,3,4,5,6,5,4,2],\ [3,4,3,6,7],\ [3,4,5,4,3,1,6,5,4,2,3,1,4,5,6],\ [3,4,7],$$
$$[4,2,3,1,4,5,4,2,3,1,4,3,5,6,7,6,8,7,6,5,4,3,1],\ [4,2,3,4,5,6,5,4,2,3,4,5,8],\ [4,5,8],\ [4,7,8],$$
$$[1,3,1,4,5,6,5,7,6,5,4,2,3,1,4,5,6,7,8,7,6,5,4,2,3]\big\},$$

one obtains 90 solutions for w_i in (5.233) with $\ell_T(w_i) = 3$ and w_i of type A_3:

$$w_i \in \big\{[1,2,3,1,4,3,1,5,6,7,6,5,8,7,6,5,4,2,3,1,4,3,5,4,2,6,7],$$
$$[1,2,3,1,4,3,5,4,6,5,7,6,5,4,3,1,8,7,6,5,4,2,3,1,4],\ [1,2,3,1,4,5,4,6,7,8,7,6,5,4,2,3,1,4,3,5,4],$$
$$[1,2,3,1,4,5,4,6,7,8,7,6,5,4,2,3,1,4,3,5,6],\ [1,2,3,1,4,5,4,6,7,8,7,6,5,4,2,3,1,4,5],$$
$$[1,2,3,1,4,5,6,5,4,7,6,5,4,2,3,1,4,3,5,6,8],\ [1,2,3,1,4,5,6,5,7,8,7,6,5,4,2,3,1,4,3,5,6],$$
$$[1,2,3,1,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5],\ [1,2,3,1,4,5,6,7,8,7,6,5,4,2,3,1,4],$$
$$[1,2,3,4,2,3,5,4,6,5,7,6,5,4,2,3,8,7,6,5,4,2,3],\ [1,2,3,4,2,5,4,2,6,7,8,7,6,5,4,2,3],$$
$$[1,2,3,4,2,5,4,6,5,4,2,7,6,5,4,2,3],\ [1,2,3,4,2,5,4,6,5,4,2,7,8,7,6,5,4,2,3],\ [1,2,3,4,5,4,6,7,6,5,4,2,3,1,4,5,8],$$
$$[1,2,3,4,5,6,7,6,5,4,2,3,1,4,5],\ [1,2,3,4,5,6,7,6,5,4,2,3,1,4,8],\ [1,2,3,4,5,6,7,6,5,4,2,3,4],$$
$$[1,2,3,4,5,6,7,6,5,4,2,3,8],\ [1,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,5],$$
$$[1,2,3,4,5,6,7,8,7,6,5,4,2,3,4],\ [1,2,4,2,3,1,4,3,5,4,3,1,6,7,8,7,6,5,4,3,1],$$
$$[1,2,4,2,3,1,4,5,6,7,6,5,4,2,3,1,4,5,6,7,8,7,6,5,4,2,3],\ [1,2,4,2,3,1,7,6,5,4,2,3,1,4,3,5,4,2,6,7,8],$$
$$[1,2,4,2,3,4,5,4,6,5,4,7,6,5,4,3,1],\ [1,2,4,2,3,4,5,4,6,7,6,5,4,3,1],\ [1,2,4,2,3,4,5,4,6,7,8,7,6,5,4,3,1],$$
$$[1,2,4,2,3,4,5,6,7,6,5,4,3,1,8],\ [1,2,4,5,4,2,3,4,5,6,5,4,3,1,7,8,7],\ [1,2,4,5,4,2,3,4,5,6,5,4,3,1,7],$$
$$[1,2,4,5,4,2,3,4,5,6,5,4,3],\ [1,2,4,5,4,2,3,4,5,6,5,7,6,5,4,3,1],\ [1,2,4,5,4,2,3,4,5,6,7,6,5,4,3,1,8],$$
$$[1,3,1,4,5,4,2,3,1,4,5,6,5,7,6,5,8,7,6,5,4,2,3],\ [1,3,1,4,5,4,2,3,1,4,5,6,5,7,8,7,6,5,4,2,3],$$
$$[1,3,1,4,5,4,2,3,1,4,5,6,7,8,7,6,5,4,2,3,4],\ [1,3,1,4,5,6,5,4,2,3,1,4,5,6,7,6,5,4,2,3,8],$$
$$[1,3,1,4,5,6,5,4,2,3,1,4,5,6,7,6,5,4,2],\ [1,3,1,4,5,6,5,4,2,3,1,4,5,6,7,6,8,7,6,5,4,2,3],$$
$$[1,3,4,2,3,4,5,4,2,6,7,8,7,6,5,4,2,3,4],\ [1,3,4,5,4,2,3,1,4,5,6,5,4,2,7],$$
$$[1,3,4,5,4,2,3,1,4,5,6,5,7,6,5,4,2],\ [1,3,4,5,4,2,3,4,5,6,5,4,2],$$
$$[1,3,4,5,4,3,1,6,7],\ [1,3,4,5,4,3,6],\ [1,3,4,5,6,5,4,3,1,7,8],\ [1,4,2,3,1,4,5,6,7,8,7,6,5,4,2,3,1,4,5],$$
$$[1,4,2,3,4,5,6,7,6,5,4,2,3,1,4,5,6],\ [1,4,2,3,4,5,6,7,6,5,4,2,3,1,4,5,8],\ [1,4,2,3,4,5,6,7,6,5,4,2,3,4,8],$$
$$[1,5,4,2,3,1,4,5,6,7,6],\ [1,5,4,2,3,1,4,5,6],\ [1,5,4,2,3,4,5,6,5,4,2,3,1,4,5,7,8],\ [1,6,5,4,2,3,1,4,5,6,7],$$
$$[2,3,1,4,2,3,4,5,4,6,5,7,6,5,4,3,8,7,6,5,4,2,3,1,4],\ [2,3,1,4,5,6,5,4,2,3,1,4,5,6,7,6,5,4,3,1,8],$$
$$[2,3,1,4,5,6,5,4,2,3,1,4,5,6,7,6,8,7,6,5,4,3,1],\ [2,3,1,4,5,6,5,4,2,3,4,5,6,7,6,5,4,3,1],\ [2,3,4,2,5,4,2,6,5,4,2],$$
$$[2,3,4,2,5,6,5,4,2],\ [2,3,4,2,5,6,7,6,5,4,2],\ [2,3,4,3,1,5,6,7,6,5,4,2,3,1,4,5,6,7,8,7,6,5,4,3,1],$$
$$[2,3,4,5,6,5,4,2,3,4,5],\ [2,3,4,5,6,5,4,2,3,4,7,8,7],\ [2,3,4,5,6,5,4,2,3,4,7],$$
$$[2,3,4,5,6,5,4,2,3,7,8],\ [2,3,4,5,6,5,4,2,7],\ [2,3,4,5,6,7,6,5,4,2,3,4,8],\ [2,4,2,3,4,5,4,3,6],$$
$$[2,4,2,3,4,5,4,6,5,4,3],\ [2,4,2,5,6],\ [2,4,5,4,2,6,7],\ [3,1,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,6],$$
$$[3,1,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,5],\ [3,1,4,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,5],$$
$$[3,1,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,6,7],\ [3,1,6,5,4,2,3,1,4,3,5,6,7,8,7],$$
$$[3,1,6,5,4,2,3,1,4,3,5,6,7],\ [3,1,7,6,5,4,2,3,1,4,3,5,6,7,8],$$

$$
\begin{gathered}
[3,4,2,3,1,4,5,4,2,3,4,5,6,7,8,7,6,5,4,3,1],\ [3,4,2,3,4,5,4,2,6,7,6],\ [3,4,2,3,4,5,4,2,6],\ [3,4,2,3,4,5,4,6,5,4,2],\\
[3,4,2,3,4,5,6,5,4,2,7],\ [3,4,3,5,6],\ [3,4,5],\ [4,2,3,4,5,4,2,3,4,6,7],\\
[4,2,3,4,5,6,5,4,2,3,4,7,8],\ [4,2,3,4,5,6,5],\ [4,2,3,4,5],\ [5,4,2,3,4,5,6]\big\},
\end{gathered}
$$

one obtains 15 solutions for w_i in (5.233) with $\ell_T(w_i)=4$ and w_i of type $A_1^2*A_2$:

$$
\begin{gathered}
w_i\in\big\{[1,2,3,1,4,5,4,6,5,4,2,3,4,5,7,6,5,4,2,3,4,5,8,7,6,5,4,2,3,1],\\
[1,2,3,4,3,1,5,6,5,7,6,5,4,2,3,1,4,3,5,4,6,5,8,7,6,5,4,2,3,1],\\
[1,2,3,4,5,4,6,5,4,2,3,4,5,7,6,5,4,2,3,1,4,5],\\
[1,2,3,4,5,4,6,5,4,3,1,7,6,5,4,2,3,1,4,3,5,6],\\
[1,3,1,4,2,3,4,5,6,5,7,6,5,4,2,3,1,8,7,6,5,4,2,3,1,4,3,5,4,2,6,7],\\
[1,3,1,4,2,3,5,4,2,3,1,7,8,7,6,5,4,2,3,1,4,3,5,4,2,6,5,4,3,1,7,6,8,7],\\
[1,4,3,1,5,4,2,3,1,4,3,5,4,6,7,6,8,7,6,5,4,2,3,4],\\
[1,4,5,4,2,3,1,4,5,6,5,4,2,3,1,4,5,7,8,7],\\
[1,4,5,4,2,3,4,5,6,5,4,2,3,4,5,8],\\
[2,4,2,3,1,4,5,4,3,6,7,6,8,7,6,5,4,2,3,1,4,3,5,4,2,6,7,8],\\
[2,4,2,3,1,4,6,7,6,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,6,7],\\
[3,1,4,2,3,4,5,6,5,7,6,5,4,2,3,4,5,6,8,7,6,5,4,2,3,1,4,3,5,6],\\
[3,1,5,6,5,4,2,3,1,4,3,5,4,6,5,8],\\
[3,4,2,3,4,5,4,2,3,1,4,7,8,7,6,5,4,2,3,1,4,3,5,6,7,8],\\
[4,2,3,4,5,4,2,3,4,6,5,4,2,3,4,7,8,7]\big\},
\end{gathered}
\tag{5.237}
$$

one obtains 45 solutions for w_i in (5.233) with $\ell_T(w_i)=4$ and w_i of type A_1*A_3:

$$
\begin{gathered}
w_i\in\big\{[1,2,3,1,4,2,5,4,2,6,5,4,2,7,8,7,6,5,4,2,3,1,4,5],\\
[1,2,3,1,4,2,5,4,6,5,4,2,3,4,5,7,6,5,4,3,8,7,6,5,4,2,3,1],\\
[1,2,3,1,4,3,1,5,6,5,7,6,5,8,7,6,5,4,2,3,1,4,3,5,4,2,6,7],\\
[1,2,3,1,4,3,5,4,6,5,4,7,6,5,4,3,8,7,6,5,4,2,3,1,4,3,5,6],\\
[1,2,3,1,4,5,4,6,5,4,7,6,5,4,2,3,1,4,3,5,6,8],\ [1,2,3,4,2,3,5,4,6,5,4,7,6,5,4,2,3,8,7,6,5,4,2,3],\\
[1,2,3,4,2,3,5,4,6,5,7,6,5,4,2,3,4,5,8,7,6,5,4,2,3,1],\ [1,2,3,4,2,5,4,2,3,4,6,5,4,7,6,5,4,3,8,7,6,5,4,2,3,1],\\
[1,2,3,4,2,5,4,2,6,5,4,2,7,6,5,4,2,3,1,4,5,8],\ [1,2,3,4,3,5,4,6,5,4,3,7,6,5,4,2,3,1,4,5],\\
[1,2,3,4,3,5,4,6,5,4,3,7,6,5,4,2,3,4],\ [1,2,3,4,5,6,5,4,2,3,4,5,7,6,5,4,2,3],\\
[1,2,4,2,3,1,4,5,6,5,7,6,5,4,2,3,1,4,5,6,7,8,7,6,5,4,2,3],\ [1,2,4,2,3,4,5,4,3,6,7,6,5,4,2,3,4,8],\\
[1,2,4,5,4,2,3,4,5,6,5,4,3,8],\ [1,3,1,4,2,3,4,5,4,6,5,7,6,5,4,2,8,7,6,5,4,2,3,1,4,3,5,6],\\
[1,3,1,4,2,3,4,5,4,6,5,7,6,5,4,2,8,7,6,5,4,2,3,1,4,5],\ [1,3,1,4,5,4,2,3,1,4,5,6,7,6,8,7,6,5,4,2,3,4],\\
[1,3,4,2,3,5,4,2,3,1,4,6,7,6,8,7,6,5,4,2,3,1,4,3,5,4,2,6,7,8],\ [1,3,4,3,5,4,2,3,1,4,5,6,5,4,2,7],\\
[1,3,4,3,5,4,2,3,4,5,6,5,4,2],\ [1,4,2,3,1,4,3,5,4,3,6,5,7,6,5,8,7,6,5,4,2,3,1,4],\\
[1,4,2,3,4,5,4,2,3,4,6,5,7,6,5,8,7,6,5,4,2,3,1,4],\ [1,4,2,3,4,5,4,2,3,4,6,7,8,7,6,5,4,2,3,4],\\
[1,4,5,4,2,3,4,5,6,5,4,2,3,1,4,5,7,8],\ [1,5,4,2,3,1,4,3,5,4,6,5,4,3,1,7,8,7],\\
[2,3,1,4,2,5,6,5,4,2,3,1,4,5,6,7,6,8,7,6,5,4,3,1],\ [2,3,1,4,5,4,6,5,4,2,3,1,4,5,6,7,6,5,4,3,1,8],\\
[2,3,1,4,5,4,6,5,4,2,3,4,5,6,7,6,5,4,3,1],\ [2,3,4,2,3,1,5,6,7,6,5,4,2,3,1,4,5,6,7,8,7,6,5,4,3,1],\\
[2,3,4,2,3,5,4,2,3,6,5,7,6,8,7,6,5,4,2,3],\ [2,3,4,2,3,5,4,2,6,5,7,8,7,6,5,4,2,3],\\
[2,3,4,2,3,5,4,6,5,4,7,6,5,4,2,3],\ [2,3,4,2,5,4,2,6,5,4,2,3,4,7,8,7],\\
[2,3,4,5,6,5,4,2,3,4,5,8],\ [2,4,2,3,1,4,5,4,8,7,6,5,4,2,3,1,4,3,5,6,7,8],\\
[2,4,2,3,1,5,6,5,7,6,5,4,2,3,1,4,3,5,6,7],\ [2,4,2,3,4,5,4,3,6,5,4,2,3,4,7,8],\ [2,4,2,3,4,5,4,6,5,4,3,8],\\
[3,1,4,2,3,4,5,6,7,6,5,4,2,3,4,5,6,8,7,6,5,4,2,3,1,4],\ [3,1,4,5,4,2,3,4,5,6,7,6,8,7,6,5,4,2,3,1,4,5],
\end{gathered}
$$

$$\begin{aligned}&[3,1,4,5,4,2,3,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,6,7],\ [3,1,4,5,4,6,5,4,2,3,1,4,3,5,6,7,8,7],\\ &\quad[3,1,4,5,4,6,5,4,2,3,1,4,5,6],\ [3,4,2,3,1,4,5,4,2,3,4,5,6,7,6,8,7,6,5,4,3,1]\big\},\end{aligned}\tag{5.238}$$

one obtains 5 solutions for w_i in (5.233) with $\ell_T(w_i)=4$ and w_i of type A_2^2:

$$\begin{aligned}w_i\in\big\{&[1,2,3,4,2,5,4,2,3,4,6,5,7,6,5,4,3,8,7,6,5,4,2,3,1,4],\\ &[1,2,4,2,3,4,5,4,3,6,5,7,6,5,4,2,3,4],\\ &[1,3,1,4,5,4,2,3,4,5,6,5,4,2,7,6,8,7,6,5,4,2,3,1,4,5],\\ &[2,3,1,4,2,5,4,6,5,4,2,3,1,4,5,6,7,8,7,6,5,4,3,1],\\ &[2,3,4,2,3,5,4,6,5,4,2,7,6,5,4,2,3,8]\big\},\end{aligned}\tag{5.239}$$

one obtains 18 solutions for w_i in (5.233) with $\ell_T(w_i)=4$ and w_i of type A_4:

$$\begin{aligned}w_i\in\big\{&[1,2,3,1,4,3,5,4,6,5,7,6,5,4,3,8,7,6,5,4,2,3,1,4],\ [1,2,3,1,4,5,6,7,8,7,6,5,4,2,3,1,4,3,5,6],\\ &[1,2,3,4,2,5,4,2,6,5,7,8,7,6,5,4,2,3],\ [1,2,3,4,2,5,4,2,6,7,8,7,6,5,4,2,3,4],\\ &[1,2,3,4,2,5,4,6,5,4,2,7,6,5,4,2,3,8],\ [1,2,3,4,5,6,7,6,5,4,2,3,1,4,5,8],\ [1,2,4,2,3,4,5,4,6,5,7,6,5,4,3,1],\\ &[1,2,4,2,3,4,5,4,6,7,6,5,4,3,1,8],\ [1,2,4,5,4,2,3,4,5,6,5,4,3,1,7,8],\\ &[1,3,1,4,5,4,2,3,1,4,5,6,5,7,6,8,7,6,5,4,2,3],\ [1,4,2,3,1,4,3,5,4,6,7,8,7,6,5,4,3,1],\\ &[1,5,4,2,3,4,5,6],\ [2,3,4,2,5,4,6,5,4,2],\ [2,3,4,2,5,6,5,4,2,7],\ [2,3,4,5,6,5,4,2,3,4,7,8],\\ &[2,4,2,3,1,7,6,5,4,2,3,1,4,3,5,6,7,8],\ [3,1,6,5,4,2,3,1,4,5,6,7],\ [3,4,2,3,4,5,4,2,6,7]\big\},\end{aligned}\tag{5.240}$$

and one obtains 5 solutions for w_i in (5.233) with $\ell_T(w_i)=4$ and w_i of type D_4:

$$\begin{aligned}w_i\in\big\{&[1,2,3,1,4,5,6,7,8,7,6,5,4,2,3,1,4,5],\ [1,2,3,4,5,6,7,6,5,4,2,3,4,8],\\ &[1,5,4,2,3,1,4,5,6,7],\ [3,1,6,5,4,2,3,1,4,3,5,6,7,8],\ [4,2,3,4,5,6]\big\},\end{aligned}\tag{5.241}$$

where $\{s_1,s_2,s_3,s_4,s_5,s_6,s_7,s_8\}$ is a simple system of generators of E_8, corresponding to the Dynkin diagram displayed in Figure 6, and each of them gives rise to $m/2$ elements of $\operatorname{Fix}_{NC^m(E_8)}(\phi^p)$ since i ranges from 1 to $m/2$. There are no solutions for w_i in (5.233) with w_i of type A_1^4.

Letting the computer find all solutions in cases (iii)–(v) would take years. However, the number of these solutions can be computed from our knowledge of the solutions in Case (ii) according to type, if this information is combined with the decomposition numbers in the sense of [14, 15, 17] (see the end of Section 2) and some elementary (multiset) permutation counting. The decomposition numbers for A_2, A_3, A_4, and D_4 of which we make use can be found in the appendix of [15].

To begin with, the number of solutions of (5.234) with $\ell_1=\ell_2=1$ is equal to

$$n_{1,1}:=150\cdot 2+100\cdot N_{A_2}(A_1,A_1)=600,$$

since an element of type A_1^2 can be decomposed in two ways into a product of two elements of absolute length 1, while for an element of type A_2 this can be done in $N_{A_2}(A_1,A_1)=3$ ways. Similarly, the number of solutions of (5.234) with $\ell_1=2$ and $\ell_2=1$ is equal to

$$n_{2,1}:=75\cdot 3+165\cdot(1+N_{A_2}(A_1,A_1))+90\cdot N_{A_3}(A_2,A_1)=1425,$$

the number of solutions of (5.234) with $\ell_1=3$ and $\ell_2=1$ is equal to

$$\begin{aligned}n_{3,1}:=15\cdot(2+N_{A_2}(A_1,A_1))+45\cdot(1+N_{A_3}(A_2,A_1))+5\cdot(2N_{A_2}(A_1,A_1))\\ +18\cdot(N_{A_4}(A_3,A_1)+N_{A_4}(A_1*A_2,A_1))+5\cdot(N_{D_4}(A_3,A_1)+N_{D_4}(A_1^3,A_1))=660,\end{aligned}$$

the number of solutions of (5.234) with $\ell_1 = \ell_2 = 2$ is equal to

$$n_{2,2} := 15 \cdot (2 + 2N_{A_2}(A_1, A_1)) + 45 \cdot (2N_{A_3}(A_2, A_1)) + 5 \cdot (2 + N_{A_2}(A_1, A_1)^2) \\ + 18 \cdot (N_{A_4}(A_2, A_2) + N_{A_4}(A_1^2, A_1^2) + 2N_{A_4}(A_2, A_1^2)) \\ + 5 \cdot (N_{D_4}(A_2, A_2) + 2N_{D_4}(A_2, A_1^2)) = 1195,$$

the number of solutions of (5.235) with $\ell_1 = \ell_2 = \ell_3 = 1$ is equal to

$$n_{1,1,1} := 75 \cdot 3! + 165 \cdot (3N_{A_2}(A_1, A_1)) + 90N_{A_3}(A_1, A_1, A_1) = 3375,$$

the number of solutions of (5.235) with $\ell_1 = 2$ and $\ell_2 = \ell_3 = 1$ is equal to

$$n_{2,1,1} := 15 \cdot (2 + N_{A_2}(A_1, A_1) + 2 \cdot 2 \cdot N_{A_2}(A_1, A_1)) + 45 \cdot (2N_{A_3}(A_2, A_1) + N_{A_3}(A_1, A_1, A_1)) \\ + 5 \cdot (2N_{A_2}(A_1, A_1) + 2N_{A_2}(A_1, A_1)^2) + 18 \cdot (N_{A_4}(A_2, A_1, A_1) + N_{A_4}(A_1^2, A_1, A_1)) \\ + 5 \cdot (N_{D_4}(A_2, A_1, A_1) + N_{D_4}(A_1^2, A_1, A_1)) = 2850,$$

and the number of solutions of (5.236) is equal to

$$n_{1,1,1,1} := 15 \cdot (12N_{A_2}(A_1, A_1)) + 45 \cdot (4N_{A_3}(A_1, A_1, A_1)) + 5 \cdot (6N_{A_2}(A_1, A_1)^2) \\ + 18 \cdot N_{A_4}(A_1, A_1, A_1, A_1) + 5 \cdot N_{D_4}(A_1, A_1, A_1, A_1) = 6750.$$

In total, we obtain

$$1 + (45 + 150 + 100 + 75 + 165 + 90 + 15 + 45 + 5 + 18 + 5)\frac{m}{2} + (n_{1,1} + 2n_{2,1} + 2n_{3,1} + n_{2,2})\binom{m/2}{2} \\ + (n_{1,1,1} + 3n_{2,1,1})\binom{m/2}{3} + n_{1,1,1,1}\binom{m/2}{4} = \frac{(5m+4)(3m+2)(5m+2)(15m+4)}{64}$$

elements in $\operatorname{Fix}_{NC^m(E_8)}(\phi^p)$, which agrees with the limit in (5.198h).

Finally, we turn to (5.198l). By Remark 3, the only choices for h_2 and m_2 to be considered are $h_2 = 1$ and $m_2 = 7$, $h_2 = 2$ and $m_2 = 8$, $h_2 = 2$ and $m_2 = 7$, $h_2 = 2$ and $m_2 = 4$, respectively $h_2 = m_2 = 2$. These correspond to the choices $p = 30m/7$, $p = 15m/8$, $p = 15m/7$, $p = 15m/4$, respectively $15m/2$, out of which only $p = 15m/8$ has not yet been discussed and belongs to the current case. The corresponding action of ϕ^p is given by

$$\phi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(*; c^2 w_{\frac{m}{8}+1} c^{-2}, c^2 w_{\frac{m}{8}+2} c^{-2}, \dots, c^2 w_m c^{-2}, c w_1 c^{-1}, \dots, c w_{\frac{m}{8}} c^{-1}\big),$$

so that we have to solve

$$t_1(c^{13}t_1c^{-13})(c^{11}t_1c^{-11})(c^9t_1c^{-9})(c^7t_1c^{-7})(c^5t_1c^{-5})(c^3t_1c^{-3})(ct_1c^{-1}) = c \qquad (5.242)$$

for t_1 with $\ell_T(t_1) = 1$. A computation with the help of Stembridge's *Maple* package `coxeter` [29] finds no solution. Hence, the left-hand side of (3.3) is equal to 1, as required.

6. Cyclic sieving II

In this section we present the second cyclic sieving conjecture due to Bessis and Reiner [7, Conj. 6.5].

Let $\psi : NC^m(W) \to NC^m(W)$ be the map defined by

$$(w_0; w_1, \dots, w_m) \mapsto \big(cw_m c^{-1}; w_0, w_1, \dots, w_{m-1}\big). \tag{6.1}$$

For $m = 1$, we have $w_0 = cw_1^{-1}$, so that this action reduces to the inverse of the *Kreweras complement* K_{id}^c as defined by Armstrong [1, Def. 2.5.3].

It is easy to see that $\psi^{(m+1)h}$ acts as the identity, where h is the Coxeter number of W (see (7.1) below). By slight abuse of notation, let C_2 be the cyclic group of order $(m+1)h$ generated by ψ. (The slight abuse consists in the fact that we insist on C_2 to be a cyclic group of order $(m+1)h$, while it may happen that the order of the action of ψ given in (6.1) is actually a proper divisor of $(m+1)h$.)

Given these definitions, we are now in the position to state the second cyclic sieving conjecture of Bessis and Reiner. By the results of [16] and of this paper, it becomes the following theorem.

Theorem 10. *For an irreducible well-generated complex reflection group W and any $m \ge 1$, the triple $(NC^m(W), \mathrm{Cat}^m(W;q), C_2)$, where $\mathrm{Cat}^m(W;q)$ is the q-analogue of the Fuß–Catalan number defined in* (3.2)*, exhibits the cyclic sieving phenomenon in the sense of Reiner, Stanton and White* [23]*. Here, n is the rank of W, $d_1, d_2, \dots, d_n$ are the degrees of W, h is the Coxeter number of W, and $[\alpha]_q := (1-q^\alpha)/(1-q)$.*

By definition of the cyclic sieving phenomenon, we have to prove that

$$|\operatorname{Fix}_{NC^m(W)}(\psi^p)| = \mathrm{Cat}^m(W;q)\big|_{q=e^{2\pi i p/(m+1)h}}, \tag{6.2}$$

for all p in the range $0 \le p < (m+1)h$.

7. Auxiliary results II

This section collects several auxiliary lemmas which allow us to reduce the problem of proving Theorem 10, respectively the equivalent statement (6.2), for the 26 exceptional groups listed in Section 2 to a finite problem. While Lemmas 12 and 13 cover special choices of the parameters, Lemmas 11 and 14 afford an inductive procedure. More precisely, if we assume that we have already verified Theorem 10 for all groups of smaller rank, then Lemmas 11 and 14, together with Lemmas 12 and 15, reduce the verification of Theorem 10 for the group that we are currently considering to a finite problem; see Remark 4. The final lemma of this section, Lemma 16, disposes of complex reflection groups with a special property satisfied by their degrees.

Let $p = a(m+1) + b$, $0 \le b < m+1$. We have

$$\begin{aligned}\psi^p\big((w_0; w_1, \dots, w_m)\big) \\ = (c^{a+1}w_{m-b+1}c^{-a-1}; c^{a+1}w_{m-b+2}c^{-a-1}, \dots, c^{a+1}w_m c^{-a-1}, \\ c^a w_0 c^{-a}, \dots, c^a w_{m-b} c^{-a}\big).\end{aligned} \tag{7.1}$$

Lemma 11. *It suffices to check* (6.2) *for p a divisor of $(m+1)h$. More precisely, let p be a divisor of $(m+1)h$, and let k be another positive integer with $\gcd(k, (m+1)h/p) = 1$,* *then we have*

$$\operatorname{Cat}^m(W;q)\big|_{q=e^{2\pi i p/(m+1)h}} = \operatorname{Cat}^m(W;q)\big|_{q=e^{2\pi i kp/(m+1)h}} \tag{7.2}$$

and

$$|\operatorname{Fix}_{NC^m(W)}(\psi^p)| = |\operatorname{Fix}_{NC^m(W)}(\psi^{kp})|. \tag{7.3}$$

Proof. For (7.3), this follows in the same way as (4.3) in Lemma 3.

For (7.2), we must argue differently than in Lemma 3. Let us write $\zeta = e^{2\pi i p/(m+1)h}$. For a given group W, we write $S_1(W)$ for the set of all indices i such that $\zeta^{d_i - h} = 1$, and we write $S_2(W)$ for the set of all indices i such that $\zeta^{d_i} = 1$. By the rule of de l'Hospital, we have

$$\operatorname{Cat}^m(W;q)\big|_{q=e^{2\pi i p/(m+1)h}} = \begin{cases} 0 & \text{if } |S_1(W)| > |S_2(W)|,\\ \frac{\prod_{i\in S_1(W)}(mh+d_i)}{\prod_{i\in S_2(W)} d_i}\frac{\prod_{i\notin S_1(W)}(1-\zeta^{d_i-h})}{\prod_{i\notin S_2(W)}(1-\zeta^{d_i})}, & \text{if } |S_1(W)| = |S_2(W)|. \end{cases}$$

It follows from [21, Theorem 10.33, with the automorphism σ specialised to complex conjugation], that the case $|S_1(W)| < |S_2(W)|$ cannot occur.

We claim that, for the case where $|S_1(W)| = |S_2(W)|$, the factors in the quotient of products

$$\frac{\prod_{i\notin S_1(W)}(1-\zeta^{d_i-h})}{\prod_{i\notin S_2(W)}(1-\zeta^{d_i})}$$

cancel pairwise. If we assume the correctness of the claim, it is obvious that we get the same result if we replace ζ by ζ^k, where $\gcd(k,(m+1)h/p) = 1$, hence establishing (7.2).

In order to see that our claim is indeed valid, we proceed in a case-by-case fashion, making appeal to the classification of irreducible well-generated complex reflection groups, which we recalled in Section 2. First of all, since $d_n = h$, the set $S_1(W)$ is always non-empty as it contains the element n. Hence, if we want to have $|S_1(W)| = |S_2(W)|$, the set $S_2(W)$ must be non-empty as well. In other words, the integer $(m+1)h/p$ must divide at least one of the degrees $d_1, d_2, \dots, d_n$. Writing M for $(m+1)h/p$, what needs to be checked is whether the *multi*sets (that is, multiplicities of elements must be taken into account)

$$\{(d_i - h) \bmod M : i \notin S_1(W)\} \quad \text{and} \quad \{d_i \bmod M : i \notin S_2(W)\}$$

are the same. Since, for a fixed irreducible well-generated complex reflection group, there is only a finite number of possibilities for M, this amounts to a routine verification. □

Lemma 12. *Let p be a divisor of $(m+1)h$. If p is divisible by $m+1$, then* (6.2) *is true.*

Proof. According to (7.1), the action of ψ^p on $NC^m(W)$ is described by

$$\psi^p\big((w_0;w_1,\dots,w_m)\big) = \big(c^{p/(m+1)}w_0c^{-p/(m+1)};c^{p/(m+1)}w_1c^{-p/(m+1)},\dots,c^{p/(m+1)}w_mc^{-p/(m+1)}\big).$$

Hence, if $(w_0;w_1,\dots,w_m)$ is fixed by ψ^p, then each individual w_i must be fixed under conjugation by $c^{p/(m+1)}$.

Using the notation $W' = \mathrm{Cent}_W(c^{p/(m+1)})$, the previous observation means that $w_i \in W'$, $i = 1, 2, \dots, m$. By the theorem of Springer cited in the proof of Lemma 4 and by (4.5), the tuples $(w_0; w_1, \dots, w_m)$ fixed by ψ^p are in fact identical with the elements of $NC^m(W')$, which implies that

$$|\operatorname{Fix}_{NC^m(W)}(\psi^p)| = |NC^m(W')|. \tag{7.4}$$

Application of Theorem 1 with W replaced by W' and of the "limit rule" (4.4) then yields that

$$|NC^m(W')| = \prod_{\substack{1\le i\le n\\ \frac{(m+1)h}{p} \mid d_i}} \frac{mh + d_i}{d_i} = \mathrm{Cat}^m(W;q)\big|_{q=e^{2\pi i p/(m+1)h}}. \tag{7.5}$$

Combining (7.4) and (7.5), we obtain (6.2). This finishes the proof of the lemma. ☐

Lemma 13. *Equation* (6.2) *holds for all divisors p of $m+1$.*

Proof. We have

$$\mathrm{Cat}^m(W;q)\big|_{q=e^{2\pi i p/(m+1)h}} = \begin{cases} 0 & \text{if } p < m+1,\\ m+1 & \text{if } p = m+1. \end{cases}$$

On the other hand, if $(w_0; w_1, \dots, w_m)$ is fixed by ψ^p, then, because of the action (7.1), we must have $w_0 = w_p = \cdots = w_{m-p+1}$ and $w_0 = c w_{m-p+1} c^{-1}$. In particular, $w_0 \in \mathrm{Cent}_W(c)$. By the theorem of Springer cited in the proof of Lemma 4, the subgroup $\mathrm{Cent}_W(c)$ is itself a complex reflection group whose degrees are those degrees of W that are divisible by h. The only such degree is h itself, hence $\mathrm{Cent}_W(c)$ is the cyclic group generated by c. Moreover, by (4.5), we obtain that $w_0 = \varepsilon$ or $w_0 = c$. If $p = m+1$, the set $\operatorname{Fix}_{NC^m(W)}(\psi^p)$ consists of the $m+1$ elements $(w_0; w_1, \dots, w_m)$ obtained by choosing $w_i = c$ for a particular i between 0 and m, all other w_j's being equal to ε. If $p < m+1$, then there is no element in $\operatorname{Fix}_{NC^m(W)}(\psi^p)$. ☐

Lemma 14. *Let W be an irreducible well-generated complex reflection group of rank n, and let $p = m_1 h_1$ be a divisor of $(m+1)h$, where $m+1 = m_1 m_2$ and $h = h_1 h_2$. We assume that* $\gcd(h_1, m_2) = 1$. *Suppose that Theorem 10 has already been verified for all irreducible well-generated complex reflection groups with rank $< n$. If h_2 does not divide all degrees d_i, then Equation* (6.2) *is satisfied.*

Proof. Let us write $h_1 = a m_2 + b$, with $0 \le b < m_2$. The condition $\gcd(h_1, m_2) = 1$ translates into $\gcd(b, m_2) = 1$. From (7.1), we infer that

$$\begin{multline} \psi^p\big((w_0; w_1, \dots, w_m)\big)\\ = \big(c^{a+1} w_{m-m_1 b+1} c^{-a-1}; c^{a+1} w_{m-m_1 b+2} c^{-a-1}, \dots, c^{a+1} w_m c^{-a-1},\\ c^a w_0 c^{-a}, \dots, c^a w_{m-m_1 b} c^{-a}\big). \end{multline} \tag{7.6}$$

Supposing that $(w_0; w_1, \dots, w_m)$ is fixed by ψ^p, we obtain the system of equations

$$\begin{aligned} w_i &= c^{a+1} w_{i+m-m_1 b+1} c^{-a-1}, \quad i = 0, 1, \dots, m_1 b - 1,\\ w_i &= c^a w_{i-m_1 b} c^{-a}, \quad i = m_1 b, m_1 b + 1, \dots, m, \end{aligned}$$

which, after iteration, implies in particular that

$$w_i = c^{b(a+1)+(m_2-b)a} w_i c^{-b(a+1)-(m_2-b)a} = c^{h_1} w_i c^{-h_1}, \quad i = 0, 1, \dots, m.$$

It is at this point where we need $\gcd(b, m_2) = 1$. The last equation shows that each w_i, $i = 0, 1, \ldots, m$, lies in $\mathrm{Cent}_W(c^{h_1})$. By the theorem of Springer cited in the proof of Lemma 4, this centraliser subgroup is itself a complex reflection group, W' say, whose degrees are those degrees of W that are divisible by $h/h_1 = h_2$. Since, by assumption, h_2 does not divide *all* degrees, W' has rank strictly less than n. Again by assumption, we know that Theorem 10 is true for W', so that in particular,

$$|\operatorname{Fix}_{NC^m(W')}(\psi^p)| = \operatorname{Cat}^m(W'; q)\big|_{q=e^{2\pi i p/(m+1)h}}.$$

The arguments above together with (4.5) show that $\operatorname{Fix}_{NC^m(W)}(\psi^p) = \operatorname{Fix}_{NC^m(W')}(\psi^p)$. On the other hand, it is straightforward to see that

$$\operatorname{Cat}^m(W; q)\big|_{q=e^{2\pi i p/(m+1)h}} = \operatorname{Cat}^m(W'; q)\big|_{q=e^{2\pi i p/(m+1)h}}.$$

This proves (6.2) for our particular p, as required. □

Lemma 15. *Let W be an irreducible well-generated complex reflection group of rank n, and let $p = m_1 h_1$ be a divisor of $(m+1)h$, where $m+1 = m_1 m_2$ and $h = h_1 h_2$. We assume that $\gcd(h_1, m_2) = 1$. If $m_2 > n$ then*

$$\operatorname{Fix}_{NC^m(W)}(\psi^p) = \emptyset.$$

Proof. Let us suppose that $(w_0; w_1, \ldots, w_m) \in \operatorname{Fix}_{NC^m(W)}(\psi^p)$ and that there exists a $j \geq 1$ such that $w_j \neq \varepsilon$. By (7.6), it then follows for such a j that also $w_k \neq \varepsilon$ for all $k \equiv j - l m_1 b \pmod{m+1}$, where, as before, b is defined as the unique integer with $h_1 = a m_2 + b$ and $0 \leq b < m_2$. Since, by assumption, $\gcd(b, m_2) = 1$, there are exactly m_2 such k's which are distinct mod $m+1$. However, this implies that the sum of the absolute lengths of the w_i's, $0 \leq i \leq m$, is at least $m_2 > n$, a contradiction. □

Remark 4. (1) If we put ourselves in the situation of the assumptions of Lemma 14, then we may conclude that Equation (6.2) only needs to be checked for pairs (m_2, h_2) subject to the following restrictions:

$$m_2 \geq 2, \quad \gcd(h_1, m_2) = 1, \quad \text{and } h_2 \text{ divides all degrees of } W. \tag{7.7}$$

Indeed, Lemmas 12 and 14 together imply that Equation (6.2) is always satisfied except if $m_2 \geq 2$, h_2 divides all degrees of W, and $\gcd(h_1, m_2) = 1$.

(2) Still putting ourselves in the situation of Lemma 14, if $m_2 > n$ and $m_2 h_2$ does not divide any of the degrees of W, then Equation (6.2) is satisfied. Indeed, Lemma 15 says that in this case the left-hand side of (6.2) equals 0, while it is obvious that in this case the right-hand side of (6.2) equals 0 as well.

(3) It should be observed that this leaves a finite number of choices for m_2 to consider, whence a finite number of choices for (m_1, m_2, h_1, h_2). Altogether, there remains a finite number of choices for $p = h_1 m_1$ to be checked.

Lemma 16. *Let W be an irreducible well-generated complex reflection group of rank n with the property that $d_i \mid h$ for $i = 1, 2, \ldots, n$. Then Theorem 10 is true for this group W.*

Proof. By Lemma 11, we may restrict ourselves to divisors p of $(m+1)h$.

Suppose that $e^{2\pi i p/(m+1)h}$ is a d_i-th root of unity for some i. In other words, $(m+1)h/p$ divides d_i. Since d_i is a divisor of h by assumption, the integer $(m+1)h/p$ also divides

h. But this is equivalent to saying that $m+1$ divides p, and Equation (6.2) holds by Lemma 12.

Now assume that $(m+1)h/p$ does not divide any of the d_i's. Then, clearly, the right-hand side of (6.2) equals 0. Inspection of the classification of all irreducible well-generated complex reflection groups, which we recalled in Section 2, reveals that all groups satisfying the hypotheses of the lemma have rank $n \le 2$. Except for the groups contained in the infinite series $G(d,1,n)$ and $G(e,e,n)$ for which Theorem 2 has been established in [16], these are the groups $G_5, G_6, G_9, G_{10}, G_{14}, G_{17}, G_{18}, G_{21}$. We now discuss these groups case by case, keeping the notation of Lemma 14. In order to simplify the argument, we note that Lemma 15 implies that Equation (6.2) holds if $m_2 > 2$, so that in the following arguments we always may assume that $m_2 = 2$.

Case G_5. The degrees are $6, 12$, and therefore Remark 4.(1) implies that Equation (6.2) is always satisfied.

Case G_6. The degrees are $4, 12$, and therefore, according to Remark 4.(1), we need only consider the case where $h_2 = 4$ and $m_2 = 2$, that is, $p = 3(m+1)/2$. Then (7.6) becomes

$$\psi^p\big((w_0; w_1, \dots, w_m)\big) = \big(c^2 w_{\frac{m+1}{2}} c^{-2}; c^2 w_{\frac{m+3}{2}} c^{-2}, \dots, c^2 w_m c^{-2}, c w_0 c^{-1}, \dots, c w_{\frac{m-1}{2}} c^{-1}\big). \tag{7.8}$$

If $(w_0; w_1, \dots, w_m)$ is fixed by ψ^p, there must exist an i with $0 \le i \le \frac{m-1}{2}$ such that $\ell_T(w_i) = 1$, $w_i c w_i c^{-1} = c$, and all w_j, $j \ne i, \frac{m+1}{2} + i$, equal ε. However, with the help of the *GAP* package `CHEVIE` [12], one verifies that there is no such solution to this equation. Hence, the left-hand side of (6.2) is equal to 0, as required.

Case G_9. The degrees are $8, 24$, and therefore, according to Remark 4.(1), we need only consider the case where $h_2 = 8$ and $m_2 = 2$, that is, $p = 3(m+1)/2$. This is the same p as for G_6. Again, the *GAP* package `CHEVIE` [12] finds no solution. Hence, the left-hand side of (6.2) is equal to 0, as required.

Case G_{10}. The degrees are $12, 24$, and therefore, according to Remark 4.(1), we need only consider the case where $h_2 = 12$ and $m_2 = 2$, that is, $p = 3(m+1)/2$. This is the same p as for G_6. Again, the *GAP* package `CHEVIE` [12] finds no solution. Hence, the left-hand side of (6.2) is equal to 0, as required.

Case G_{14}. The degrees are $6, 24$, and therefore Remark 4.(1) implies that Equation (6.2) is always satisfied.

Case G_{17}. The degrees are $20, 60$, and therefore, according to Remark 4.(1), we need only consider the cases where $h_2 = 20$ and $m_2 = 2$, respectively that $h_2 = 4$ and $m_2 = 2$. In the first case, $p = 3(m+1)/2$, which is the same p as for G_6. Again, the *GAP* package `CHEVIE` [12] finds no solution. In the second case, $p = 15(m+1)/2$. Then (7.6) becomes

$$\psi^p\big((w_0; w_1, \dots, w_m)\big) \\ = \big(c^8 w_{\frac{m+1}{2}} c^{-8}; c^8 w_{\frac{m+3}{2}} c^{-8}, \dots, c^8 w_m c^{-8}, c^7 w_0 c^{-7}, \dots, c^7 w_{\frac{m-1}{2}} c^{-7}\big). \tag{7.9}$$

By Lemma 6, every element of $NC(W)$ is fixed under conjugation by c^3, and, thus, on elements fixed by ψ^p, the above action of ψ^p reduces to the one in (7.8). This action was already discussed in the first case. Hence, in both cases, the left-hand side of (6.2) is equal to 0, as required.

Case G_{18}. The degrees are $30, 60$, and therefore Remark 4.(1) implies that Equation (6.2) is always satisfied.

Case G_{21}. The degrees are $12, 60$, and therefore, according to Remark 4.(1), we need only consider the cases where $h_2 = 5$ and $m_2 = 2$, respectively that $h_2 = 15$ and $m_2 = 2$. In the first case, $p = 5(m + 1)/2$, so that (7.6) becomes

$$\psi^p\big((w_0; w_1, \dots, w_m)\big) \\ = (c^3 w_{\frac{m+1}{2}} c^{-3}; c^3 w_{\frac{m+3}{2}} c^{-3}, \dots, c^3 w_m c^{-3}, c^2 w_0 c^{-2}, \dots, c^2 w_{\frac{m-1}{2}} c^{-2}\big). \quad (7.10)$$

If $(w_0; w_1, \dots, w_m)$ is fixed by ψ^p, there must exist an i with $0 \le i \le \frac{m-1}{2}$ such that $\ell_T(w_i) = 1$ and $w_i c^2 w_i c^{-2} = c$. However, with the help of the *GAP* package `CHEVIE` [12], one verifies that there is no such solution to this equation. In the second case, $p = 15(m + 1)/2$. Then (7.6) becomes the action in (7.9). By Lemma 6, every element of $NC(W)$ is fixed under conjugation by c^5, and, thus, on elements fixed by ψ^p, the action of ψ^p in (7.9) reduces to the one in the first case. Hence, in both cases, the left-hand side of (6.2) is equal to 0, as required.

This completes the proof of the lemma. □

8. Case-by-case verification of Theorem 10

We now perform a case-by-case verification of Theorem 10. It should be observed that the action of ψ (given in (6.1)) is exactly the same as the action of ϕ (given in (3.1)) with m replaced by $m + 1$ *on the components* $w_1, w_2, \dots, w_{m+1}$, that is, if we disregard the 0-th component of the elements of the generalised non-crossing partitions involved. The only difference which arises is that, while the $(m + 1)$-tuples $(w_0; w_1, \dots, w_m)$ in (6.1) must satisfy $w_0 w_1 \cdots w_m = c$, for $w_1, w_2, \dots, w_{m+1}$ in (3.1) we only must have $w_1 w_2 \cdots w_{m+1} \le_T c$. The condition for $(w_0; w_1, \dots, w_m)$ of being in $\mathrm{Fix}_{NC^m(W)}(\psi^p)$ is therefore exactly the same as the condition on $w_1, w_2, \dots, w_{m+1}$ for the element $(\varepsilon; w_1, \dots, w_m, w_{m+1})$ being in $\mathrm{Fix}_{NC^{m+1}(W)}(\phi^p)$. Consequently, we may use the counting results from Section 5, except that we have to restrict our attention to those elements $(w_0; w_1, \dots, w_m, w_{m+1}) \in NC^{m+1}(W)$ for which $w_1 w_2 \cdots w_{m+1} = c$, or, equivalently, $w_0 = \varepsilon$.

In the sequel we write ζ_d for a primitive d-th root of unity.

Case G_4. The degrees are $4, 6$, and hence we have

$$\mathrm{Cat}^m(G_4; q) = \frac{[6m + 6]_q \, [6m + 4]_q}{[6]_q \, [4]_q}.$$

Let ζ be a $6(m + 1)$-th root of unity. In what follows, we abbreviate the assertion that "ζ is a primitive d-th root of unity" as "$\zeta = \zeta_d$." The following cases on the right-hand side of (6.2) do occur:

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_4; q) = m + 1, \quad \text{if } \zeta = \zeta_6, \zeta_3, \tag{8.1a}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_4; q) = \tfrac{3m+3}{2}, \quad \text{if } \zeta = \zeta_4,\ 2 \mid (m + 1), \tag{8.1b}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_4; q) = \mathrm{Cat}^m(G_4), \quad \text{if } \zeta = -1 \text{ or } \zeta = 1, \tag{8.1c}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_4; q) = 0, \quad \text{otherwise.} \tag{8.1d}$$

We must now prove that the left-hand side of (6.2) in each case agrees with the values exhibited in (8.1). The only cases not covered by Lemmas 4 and 5 are the ones in (8.1b) and (8.1d). On the other hand, the only case left to consider according to Remark 4 is the case where $h_2 = m_2 = 2$, that is the case (8.1b) where $p = 3(m+1)/2$. In particular, $m+1$ must be divisible by 2. The action of ψ^p is the same as the one in (7.8). Hence, the counting problem is the same as there, except that the underlying group now is G_4. With the help of the *GAP* package `CHEVIE` [12], one finds that each of the 3 (complex) reflections in G_4 which are less than the (chosen) Coxeter element is a valid choice for w_i, and each of these choices gives rise to $(m+1)/2$ elements in $\mathrm{Fix}_{NC^m(G_4)}(\psi^p)$ since the index i ranges from 0 to $(m-1)/2$.

Hence, in total, we obtain $3\frac{m+1}{2} = \frac{3m+3}{2}$ elements in $\mathrm{Fix}_{NC^m(G_4)}(\psi^p)$, which agrees with the limit in (8.1b).

Case G_8. The degrees are $8, 12$, and hence we have

$$\mathrm{Cat}^m(G_8; q) = \frac{[12m+12]_q\,[12m+8]_q}{[12]_q\,[8]_q}.$$

Let ζ be a $12(m+1)$-th root of unity. The following cases on the right-hand side of (6.2) do occur:

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_8; q) = m+1, \quad \text{if } \zeta = \zeta_{12}, \zeta_6, \zeta_3, \tag{8.2a}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_8; q) = \tfrac{3m+3}{2}, \quad \text{if } \zeta = \zeta_8,\ 2 \mid (m+1), \tag{8.2b}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_8; q) = \mathrm{Cat}^m(G_8), \quad \text{if } \zeta = \zeta_4, -1, 1, \tag{8.2c}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_8; q) = 0, \quad \text{otherwise.} \tag{8.2d}$$

We must now prove that the left-hand side of (6.2) in each case agrees with the values exhibited in (8.2). The only cases not covered by Lemmas 4 and 5 are the ones in (8.2b) and (8.2d). On the other hand, the only case left to consider according to Remark 4 is the case where $h_2 = 4$ and $m_2 = 2$, that is the case (8.2b) where $p = 3(m+1)/2$. In particular, $m+1$ must be divisible by 2. The action of ψ^p is the same as the one in (7.8). Hence, the counting problem is the same as there, except that the underlying group now is G_8. With the help of the *GAP* package `CHEVIE` [12], one finds that each of the 3 (complex) reflections in G_8 which are less than the (chosen) Coxeter element is a valid choice for w_i, and each of these choices gives rise to $(m+1)/2$ elements in $\mathrm{Fix}_{NC^m(G_8)}(\psi^p)$ since the index i ranges from 0 to $(m-1)/2$.

Hence, in total, we obtain $3\frac{m+1}{2} = \frac{3m+3}{2}$ elements in $\mathrm{Fix}_{NC^m(G_8)}(\psi^p)$, which agrees with the limit in (8.2b).

Case G_{16}. The degrees are $20, 30$, and hence we have

$$\mathrm{Cat}^m(G_{16}; q) = \frac{[30m+30]_q\,[30m+20]_q}{[30]_q\,[20]_q}.$$

Let ζ be a $30(m+1)$-th root of unity. The following cases on the right-hand side of (6.2) do occur:

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{16};q)=m+1,\quad \text{if } \zeta=\zeta_{30},\zeta_{15},\zeta_6,\zeta_3, \tag{8.3a}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{16};q)=\tfrac{3m+3}{2},\quad \text{if } \zeta=\zeta_{20},\zeta_4,\ 2\mid(m+1), \tag{8.3b}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{16};q)=\operatorname{Cat}^m(G_{16}),\quad \text{if } \zeta=\zeta_{10},\zeta_5,-1,1, \tag{8.3c}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{16};q)=0,\quad \text{otherwise.} \tag{8.3d}$$

We must now prove that the left-hand side of (6.2) in each case agrees with the values exhibited in (8.3). The only cases not covered by Lemmas 4 and 5 are the ones in (8.3b) and (8.3d). On the other hand, the only cases left to consider according to Remark 4 are the cases where $h_2=10$ and $m_2=2$, respectively $h_2=m_2=2$. Both cases belong to (8.3b). In the first case, we have $p=3(m+1)/2$, while in the second case we have $p=15(m+1)/2$. In particular, $m+1$ must be divisible by 2. In the first case, the action of ψ^p is the same as the one in (7.8). Hence, the counting problem is the same as there, except that the underlying group now is G_{16}. With the help of the *GAP* package `CHEVIE` [12], one finds that each of the 3 (complex) reflections in G_{16} which are less than the (chosen) Coxeter element is a valid choice for w_i, and each of these choices gives rise to $(m+1)/2$ elements in $\operatorname{Fix}_{NC^m(G_{16})}(\psi^p)$ since the index i ranges from 0 to $(m-1)/2$. On the other hand, if $p=15(m+1)/2$, then the action of ψ^p is the same as the one in (7.9). By Lemma 6, every element of $NC(G_{16})$ is fixed under conjugation by c^3, and, thus, on elements fixed by ψ^p, the action of ψ^p reduces to the one in the first case.

Hence, in total, we obtain $3\frac{m+1}{2}=\frac{3m+3}{2}$ elements in $\operatorname{Fix}_{NC^m(G_{16})}(\psi^p)$, which agrees with the limit in (8.3b).

Case G_{20}. The degrees are $12, 30$, and hence we have

$$\operatorname{Cat}^m(G_{20};q)=\frac{[30m+30]_q\,[30m+12]_q}{[30]_q\,[12]_q}.$$

Let ζ be a $30(m+1)$-th root of unity. The following cases on the right-hand side of (6.2) do occur:

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{20};q)=m+1,\quad \text{if } \zeta=\zeta_{30},\zeta_{15},\zeta_{10},\zeta_5, \tag{8.4a}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{20};q)=\tfrac{5m+5}{2},\quad \text{if } \zeta=\zeta_{12},\zeta_4,\ 2\mid(m+1), \tag{8.4b}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{20};q)=\operatorname{Cat}^m(G_{20}),\quad \text{if } \zeta=\zeta_6,\zeta_3,-1,1, \tag{8.4c}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{20};q)=0,\quad \text{otherwise.} \tag{8.4d}$$

We must now prove that the left-hand side of (6.2) in each case agrees with the values exhibited in (8.4). The only cases not covered by Lemmas 4 and 5 are the ones in (8.4b) and (8.4d). On the other hand, the only cases left to consider according to Remark 4 are the cases where $h_2=6$ and $m_2=2$, respectively $h_2=m_2=2$. Both cases belong to (8.4b). In the first case, we have $p=5(m+1)/2$, while in the second case we have $p=15(m+1)/2$. In particular, $m+1$ must be divisible by 2. In the first case, the action of ψ^p is the same as the one in (7.10). Hence, the counting problem is the same as there, except that the underlying group now is G_{20}. With the help of the *GAP* package `CHEVIE` [12], one finds that each of the 5 (complex) reflections in G_{20} which are less than the (chosen) Coxeter element is a valid choice for w_i, and each of these choices gives rise to $(m+1)/2$ elements in $\mathrm{Fix}_{NC^m(G_{20})}(\psi^p)$ since the index i ranges from 0 to $(m-1)/2$. On the other hand, if $p = 15(m+1)/2$, then the action of ψ^p is the same as the one in (7.9). By Lemma 6, every element of $NC(G_{20})$ is fixed under conjugation by c^5, and, thus, on elements fixed by ψ^p, the action of ψ^p reduces to the one in the first case.

Hence, in total, we obtain $5\frac{m+1}{2} = \frac{5m+5}{2}$ elements in $\mathrm{Fix}_{NC^m(G_{20})}(\psi^p)$, which agrees with the limit in (8.4b).

Case $G_{23} = H_3$. The degrees are $2, 6, 10$, and hence we have

$$\mathrm{Cat}^m(H_3;q) = \frac{[10m+10]_q\,[10m+6]_q\,[10m+2]_q}{[10]_q\,[6]_q\,[2]_q}.$$

Let ζ be a $10(m+1)$-th root of unity. The following cases on the right-hand side of (6.2) do occur:

$$\lim_{q\to\zeta}\mathrm{Cat}^m(H_3;q) = m+1, \quad \text{if } \zeta=\zeta_{10},\zeta_5, \tag{8.5a}$$

$$\lim_{q\to\zeta}\mathrm{Cat}^m(H_3;q) = \tfrac{5m+5}{3}, \quad \text{if } \zeta=\zeta_6,\zeta_3,\ 3\mid(m+1), \tag{8.5b}$$

$$\lim_{q\to\zeta}\mathrm{Cat}^m(H_3;q) = \mathrm{Cat}^m(H_3), \quad \text{if } \zeta=-1 \text{ or } \zeta=1, \tag{8.5c}$$

$$\lim_{q\to\zeta}\mathrm{Cat}^m(H_3;q) = 0, \quad \text{otherwise.} \tag{8.5d}$$

We must now prove that the left-hand side of (6.2) in each case agrees with the values exhibited in (8.5). The only cases not covered by Lemmas 4 and 5 are the ones in (8.5b) and (8.5d). On the other hand, the only cases left to consider according to Remark 4 are the cases where $h_2 = 1$ and $m_2 = 3$, $h_2 = 2$ and $m_2 = 3$, respectively $h_2 = m_2 = 2$. respectively . These correspond to the choices $p = 10(m+1)/3$, $p = 5(m+1)/3$, respectively $p = 5(m+1)/2$. The first two cases belong to (8.5b), while $p = 5(m+1)/2$ belongs to (8.5d).

In the case that $p = 5(m+1)/3$, the action of ψ^p is given by

$$\psi^p\big((w_0;w_1,\dots,w_m)\big) \\ = \big(c^2w_{\frac{m+1}{3}}c^{-2};c^2w_{\frac{m+4}{3}}c^{-2},\dots,c^2w_mc^{-2},cw_0c^{-1},\dots,cw_{\frac{m-2}{3}}c^{-1}\big).$$

Hence, for an i with $0 \le i \le \frac{m-2}{3}$, we must find an element $w_i = t_1$, where t_1 satisfies (5.9), and all other w_j, $j \notin \{i, i+\frac{m+1}{3}, i+\frac{2(m+1)}{3}\}$, are set equal to ε. We have found five solutions to the counting problem (5.9) in (5.10). Each of them gives rise to $(m+1)/3$ elements in $\mathrm{Fix}_{NC^m(H_3)}(\psi^p)$ since the index i ranges from 0 to $(m-2)/3$. On the other hand, if $p = 10(m+1)/3$, then the action of ψ^p is given by

$$\psi^p\big((w_0;w_1,\dots,w_m)\big) \\ = \big(c^4w_{\frac{2m+2}{3}}c^{-4};c^4w_{\frac{2m+5}{3}}c^{-4},\dots,c^4w_mc^{-4},c^3w_0c^{-3},\dots,c^3w_{\frac{2m-1}{3}}c^{-3}\big).$$

By Lemma 6, every element of $NC(H_3)$ is fixed under conjugation by c^5, and, thus, on elements fixed by ψ^p, the action of ψ^p reduces to the one in the first case.

Hence, in total, we obtain $5\frac{m+1}{3} = \frac{5m+5}{3}$ elements in $\mathrm{Fix}_{NC^m(H_3)}(\psi^p)$, which agrees with the limit in (8.5b).

If $p = 5(m+1)/2$, then the action of ψ^p is the same as the one in (7.10). The computation at the end of Case H_3 in Section 5 did not find any solutions, which is in agreement with (8.5d).

Case G_{24}. The degrees are $4, 6, 14$, and hence we have

$$\mathrm{Cat}^m(G_{24}; q) = \frac{[14m+14]_q\,[14m+6]_q\,[14m+4]_q}{[14]_q\,[6]_q\,[4]_q}.$$

Let ζ be a $14(m+1)$-th root of unity. The following cases on the right-hand side of (6.2) do occur:

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_{24}; q) = m+1, \quad \text{if } \zeta = \zeta_{14}, \zeta_7, \tag{8.6a}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_{24}; q) = \tfrac{7m+7}{3}, \quad \text{if } \zeta = \zeta_6, \zeta_3,\ 3 \mid (m+1), \tag{8.6b}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_{24}; q) = \mathrm{Cat}^m(G_{24}), \quad \text{if } \zeta = -1 \text{ or } \zeta = 1, \tag{8.6c}$$

$$\lim_{q\to\zeta} \mathrm{Cat}^m(G_{24}; q) = 0, \quad \text{otherwise.} \tag{8.6d}$$

We must now prove that the left-hand side of (6.2) in each case agrees with the values exhibited in (8.6). The only cases not covered by Lemmas 4 and 5 are the ones in (8.6b) and (8.6d). On the other hand, the only cases left to consider according to Remark 4 are the cases where $h_2 = 1$ and $m_2 = 3$, $h_2 = 2$ and $m_2 = 3$, respectively $h_2 = m_2 = 2$. These correspond to the choices $p = 14(m+1)/3$, $p = 7(m+1)/3$, respectively $p = 7(m+1)/2$. The first two cases belong to (8.6b), while $p = 7(m+1)/2$ belongs to (8.6d).

In the case that $p = 14(m+1)/3$ or $p = 7(m+1)/3$, we have found seven solutions to the counting problem (5.18) in (5.19), and each of them gives rise to $(m+1)/3$ elements in $\mathrm{Fix}_{NC^m(G_{24})}(\psi^p)$ (in the style as discussed in Case H_3). Hence, in total, we obtain $7\frac{m+1}{3} = \frac{7m+7}{3}$ elements in $\mathrm{Fix}_{NC^m(G_{24})}(\psi^p)$, which agrees with the limit in (8.6b).

If $p = 7(m+1)/2$, the relevant counting problem is (5.25). However, no element $(w_0; w_1, \dots, w_m) \in \mathrm{Fix}_{NC^m(G_{24})}(\psi^p)$ can be produced in this way since the counting problem imposes the restriction that $\ell_T(w_0) + \ell_T(w_1) + \dots + \ell_T(w_m)$ be even, which is absurd. This is in agreement with the limit in (8.6d).

Case G_{25}. The degrees are $6, 9, 12$, and hence we have

$$\mathrm{Cat}^m(G_{25}; q) = \frac{[12m+12]_q\,[12m+9]_q\,[12m+6]_q}{[12]_q\,[9]_q\,[6]_q}.$$

Let ζ be a $12(m+1)$-th root of unity. The following cases on the right-hand side of (6.2) do occur:

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{25};q) = m+1, \quad \text{if } \zeta=\zeta_{12},\zeta_4, \tag{8.7a}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{25};q) = \tfrac{4m+4}{3}, \quad \text{if } \zeta=\zeta_9,\ 3\mid(m+1), \tag{8.7b}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{25};q) = (m+1)(2m+1), \quad \text{if } \zeta=\zeta_6,-1 \tag{8.7c}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{25};q) = \operatorname{Cat}^m(G_{25}), \quad \text{if } \zeta=\zeta_3,1, \tag{8.7d}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{25};q) = 0, \quad \text{otherwise.} \tag{8.7e}$$

We must now prove that the left-hand side of (6.2) in each case agrees with the values exhibited in (8.7). The only cases not covered by Lemmas 4 and 5 are the ones in (8.7b) and (8.7e). On the other hand, the only case left to consider according to Remark 4 is the case where $h_2=m_2=3$. This corresponds to the choice $p=4(m+1)/3$, which belongs to (8.7b). We have found four solutions to the counting problem (5.30) in (5.31), and each of them gives rise to $(m+1)/3$ elements in $\operatorname{Fix}_{NC^m(G_{25})}(\psi^p)$ (in the style as discussed in Case H_3). Hence, in total, we obtain $4\frac{m+1}{3}=\frac{4m+4}{3}$ elements in $\operatorname{Fix}_{NC^m(G_{25})}(\psi^p)$, which agrees with the limit in (8.7b).

Case G_{26}. The degrees are $6,12,18$, and hence we have

$$\operatorname{Cat}^m(G_{26};q)=\frac{[18m+18]_q\,[18m+12]_q\,[18m+6]_q}{[18]_q\,[12]_q\,[6]_q}.$$

Let ζ be a $14(m+1)$-th root of unity. The following cases on the right-hand side of (6.2) do occur:

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{26};q) = m+1, \quad \text{if } \zeta=\zeta_{18},\zeta_9, \tag{8.8a}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{26};q) = \operatorname{Cat}^m(G_{26}), \quad \text{if } \zeta=\zeta_6,\zeta_3,-1,1, \tag{8.8b}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{26};q) = 0, \quad \text{otherwise.} \tag{8.8c}$$

We must now prove that the left-hand side of (6.2) in each case agrees with the values exhibited in (8.8). The only case not covered by Lemmas 4 and 5 is the one in (8.8c). On the other hand, the only cases left to consider according to Remark 4 are the cases where $h_2=6$ and $m_2=2$, respectively $h_2=m_2=2$. These correspond to the choices $p=3(m+1)/2$, respectively $p=9(m+1)/2$, both of which belong to (8.8c). The relevant counting problem is (5.35). However, no element $(w_0;w_1,\dots,w_m)\in\operatorname{Fix}_{NC^m(G_{26})}(\psi^p)$ can be produced in this way since the counting problem imposes the restriction that $\ell_T(w_0)+\ell_T(w_1)+\dots+\ell_T(w_m)$ be even, which is absurd. This is in agreement with the limit in (8.8c).

Case G_{27}. The degrees are $6,12,30$, and hence we have

$$\operatorname{Cat}^m(G_{27};q)=\frac{[30m+30]_q\,[30m+12]_q\,[30m+6]_q}{[30]_q\,[12]_q\,[6]_q}.$$

Let ζ be a $14(m+1)$-th root of unity. The following cases on the right-hand side of (6.2) do occur:

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{27};q)=m+1,\quad \text{if } \zeta=\zeta_{30},\zeta_{15},\zeta_{10},\zeta_5, \tag{8.9a}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{27};q)=\operatorname{Cat}^m(G_{27}),\quad \text{if } \zeta=\zeta_6,\zeta_3,-1,1, \tag{8.9b}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{27};q)=0,\quad \text{otherwise.} \tag{8.9c}$$

We must now prove that the left-hand side of (6.2) in each case agrees with the values exhibited in (8.9). The only case not covered by Lemmas 4 and 5 is the one in (8.9c). On the other hand, the only cases left to consider according to Remark 4 are the cases where $h_2=6$ and $m_2=3$, $h_2=m_2=3$, $h_2=6$ and $m_2=2$, respectively $h_2=m_2=2$. These correspond to the choices $p=5(m+1)/3$, $10(m+1)/3$, $5(m+1)/2$, respectively $15(m+1)/2$, all of which belong to (8.9c).

If $p=5(m+1)/3$ or $p=10(m+1)/3$, the computation with the help of the *GAP* package `CHEVIE` [12] at the end of Case G_{27} in Section 5 did not find any solutions for the corresponding counting problem. This is in agreement with the limit in (8.9c).

In the case that $5(m+1)/2$ or $15(m+1)/2$, the relevant counting problem is (5.42). However, no element $(w_0;w_1,\dots,w_m)\in\operatorname{Fix}_{NC^m(G_{27})}(\psi^p)$ can be produced in this way since the counting problem imposes the restriction that $\ell_T(w_0)+\ell_T(w_1)+\dots+\ell_T(w_m)$ be even, which is absurd. This is again in agreement with the limit in (8.9c).

CASE $G_{28}=F_4$. The degrees are $2,6,8,12$, and hence we have

$$\operatorname{Cat}^m(F_4;q)=\frac{[12m+12]_q\,[12m+8]_q\,[12m+6]_q\,[12m+2]_q}{[12]_q\,[8]_q\,[6]_q\,[2]_q}.$$

Let ζ be a $12(m+1)$-th root of unity. The following cases on the right-hand side of (6.2) do occur:

$$\lim_{q\to\zeta}\operatorname{Cat}^m(F_4;q)=m+1,\quad \text{if } \zeta=\zeta_{12}, \tag{8.10a}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(F_4;q)=\tfrac{3m+3}{2},\quad \text{if } \zeta=\zeta_8,\ 2\mid(m+1), \tag{8.10b}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(F_4;q)=(m+1)(2m+1),\quad \text{if } \zeta=\zeta_6,\zeta_3, \tag{8.10c}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(F_4;q)=\tfrac{(m+1)(3m+2)}{2},\quad \text{if } \zeta=\zeta_4, \tag{8.10d}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(F_4;q)=\operatorname{Cat}^m(F_4),\quad \text{if } \zeta=-1 \text{ or } \zeta=1, \tag{8.10e}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(F_4;q)=0,\quad \text{otherwise.} \tag{8.10f}$$

We must now prove that the left-hand side of (6.2) in each case agrees with the values exhibited in (8.10). The only cases not covered by Lemmas 4 and 5 are the ones in (8.10b) and (8.10f). On the other hand, according to Remark 4, the are no choices for h_2 and m_2 left to be considered.

CASE G_{29}. The degrees are $4,8,12,20$, and hence we have

$$\operatorname{Cat}^m(G_{29};q)=\frac{[20m+20]_q\,[20m+12]_q\,[20m+8]_q\,[20m+4]_q}{[20]_q\,[12]_q\,[8]_q\,[4]_q}.$$

Let ζ be a $20(m+1)$-th root of unity. The following cases on the right-hand side of (6.2) do occur:

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{29};q) = m+1, \quad \text{if } \zeta = \zeta_{20}, \zeta_{10}, \zeta_5, \tag{8.11a}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{29};q) = \operatorname{Cat}^m(G_{29}), \quad \text{if } \zeta = \zeta_4, -1, 1, \tag{8.11b}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{29};q) = 0, \quad \text{otherwise.} \tag{8.11c}$$

We must now prove that the left-hand side of (6.2) in each case agrees with the values exhibited in (8.11). The only case not covered by Lemmas 4 and 5 is the one in (8.11c). On the other hand, the only cases left to consider according to Remark 4, the only choices for h_2 and m_2 to be considered are $h_2 = 1$ and $m_2 = 3$, $h_2 = 2$ and $m_2 = 3$, $h_2 = 4$ and $m_2 = 3$, $h_2 = 4$ and $m_2 = 2$, respectively $h_2 = m_2 = 4$. These correspond to the choices $p = 20(m+1)/3$, $p = 10(m+1)/3$, $p = 5(m+1)/3$, $p = 5(m+1)/2$, respectively $p = 5(m+1)/4$, all of which belong to (8.11c).

In the case that $p = 20(m+1)/3$, $p = 10(m+1)/3$, or $p = 5(m+1)/3$, the relevant counting problem is (5.55). However, no element $(w_0; w_1, \dots, w_m) \in \operatorname{Fix}_{NC^m(G_{27})}(\psi^p)$ can be produced in this way since the counting problem imposes the restriction that $\ell_T(w_0) + \ell_T(w_1) + \cdots + \ell_T(w_m)$ be divisible by 3, which is absurd. This is in agreement with the limit in (8.11c).

In the case that $p = 5(m+1)/2$, the relevant counting problem is (5.64), for which we did not find any solutions. This is again in agreement with the limit in (8.11c).

In the case that $p = 5(m+1)/4$, the computation at the end of Case G_{29} in Section 5 did not find any solutions, which is as well in agreement with the limit in (8.11c).

Case $G_{30} = H_4$. The degrees are $2, 12, 20, 30$, and hence we have

$$\operatorname{Cat}^m(H_4;q) = \frac{[30m+30]_q\,[30m+20]_q\,[30m+12]_q\,[30m+2]_q}{[30]_q\,[20]_q\,[12]_q\,[2]_q}.$$

Let ζ be a $30(m+1)$-th root of unity. The following cases on the right-hand side of (6.2) do occur:

$$\lim_{q\to\zeta} \operatorname{Cat}^m(H_4;q) = m+1, \quad \text{if } \zeta = \zeta_{30}, \zeta_{15}, \tag{8.12a}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(H_4;q) = \tfrac{3m+3}{2}, \quad \text{if } \zeta = \zeta_{20},\ 2 \mid (m+1), \tag{8.12b}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(H_4;q) = \tfrac{5m+5}{2}, \quad \text{if } \zeta = \zeta_{12},\ 2 \mid (m+1), \tag{8.12c}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(H_4;q) = \tfrac{(m+1)(3m+2)}{2}, \quad \text{if } \zeta = \zeta_{10}, \zeta_5, \tag{8.12d}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(H_4;q) = \tfrac{(m+1)(5m+2)}{2}, \quad \text{if } \zeta = \zeta_6, \zeta_3, \tag{8.12e}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(H_4;q) = \tfrac{(m+1)(15m+1)}{4}, \quad \text{if } \zeta = \zeta_4,\ 2 \mid (m+1), \tag{8.12f}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(H_4;q) = \operatorname{Cat}^m(H_4), \quad \text{if } \zeta = -1 \text{ or } \zeta = 1, \tag{8.12g}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(H_4;q) = 0, \quad \text{otherwise.} \tag{8.12h}$$

We must now prove that the left-hand side of (6.2) in each case agrees with the values exhibited in (8.12). The only cases not covered by Lemmas 4 and 5 are the ones in

(8.12b), (8.12c), (8.12f), and (8.12h). On the other hand, the only cases left to consider according to Remark 4 are the cases where $h_2 = 2$ and $m_2 = 4$, respectively $h_2 = m_2 = 2$. These correspond to the choices $p = 15(m+1)/2$, respectively $p = 15(m+1)/4$, out of which the first belongs to (8.12f), while the second belongs to (8.12h).

In the case that $p = 15(m+1)/2$, the action of ψ^p is the same as the one in (7.9). We have found eight solutions to the counting problem (5.80) in (5.83), each of them giving rise to $(m+1)/2$ elements in $\operatorname{Fix}_{NC^m(H_4)}(\psi^p)$ since the index i (in (5.80)) ranges from 0 to $(m-1)/2$, and we have found 30 solutions to the counting problem (5.82) in (5.84), each of them giving rise to $\binom{(m+1)/2}{2}$ elements in $\operatorname{Fix}_{NC^m(H_4)}(\psi^p)$ since $0 \le i_1 < i_2 \le (m-1)/2$ (in (5.82)). Hence, we obtain $8\frac{m+1}{2} + 30\binom{(m+1)/2}{2} = \frac{(m+1)(15m+1)}{4}$ elements in $\operatorname{Fix}_{NC^m(H_4)}(\psi^p)$, which agrees with the limit in (8.12f).

If $p = 15(m+1)/4$, the computation at the end of Case H_4 in Section 5 did not find any solutions, which is in agreement with the limit in (8.12h).

CASE G_{32}. The degrees are $12, 18, 24, 30$, and hence we have

$$\operatorname{Cat}^m(G_{32}; q) = \frac{[30m+30]_q\,[30m+24]_q\,[30m+18]_q\,[30m+12]_q}{[30]_q\,[24]_q\,[18]_q\,[12]_q}.$$

Let ζ be a $30(m+1)$-th root of unity. The following cases on the right-hand side of (6.2) do occur:

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{32}; q) = m+1, \quad \text{if } \zeta = \zeta_{30}, \zeta_{15}, \zeta_{10}, \zeta_5, \tag{8.13a}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{32}; q) = \tfrac{5m+5}{4}, \quad \text{if } \zeta = \zeta_{24}, \zeta_8,\ 4 \mid (m+1), \tag{8.13b}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{32}; q) = \tfrac{(5m+5)(5m+3)}{8}, \quad \text{if } \zeta = \zeta_{12}, \zeta_4,\ 2 \mid (m+1), \tag{8.13c}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{32}; q) = \operatorname{Cat}^m(G_{32}), \quad \text{if } \zeta = \zeta_6, \zeta_3, -1, 1, \tag{8.13d}$$

$$\lim_{q\to\zeta} \operatorname{Cat}^m(G_{32}; q) = 0, \quad \text{otherwise.} \tag{8.13e}$$

We must now prove that the left-hand side of (6.2) in each case agrees with the values exhibited in (8.13). The only cases not covered by Lemmas 4 and 5 are the ones in (8.13b), (8.13c), and (8.13e). On the other hand, the only cases left to consider according to Remark 4 are the cases where $h_2 = 2$ and $m_2 = 4$, $h_2 = 6$ and $m_2 = 4$, $h_2 = m_2 = 3$, $h_2 = 6$ and $m_2 = 3$, $h_2 = m_2 = 2$, respectively $h_2 = 6$ and $m_2 = 2$. These correspond to the choices $p = 15(m+1)/4$, $p = 5(m+1)/4$, $p = 10(m+1)/3$, $p = 5(m+1)/3$, $p = 15(m+1)/2$, respectively $p = 5(m+1)/2$, out of which the first two belong to (8.13b), the next two belong to (8.13e), and the last two belong to (8.13c).

In the case that $p = 15(m+1)/4$ or $p = 5(m+1)/4$, we have found five solutions to the counting problem (5.88) in (5.89), each of them giving rise to $(m+1)/4$ elements in $\operatorname{Fix}_{NC^m(G_{32})}(\psi^p)$. Hence, we obtain $5\frac{m+1}{4} = \frac{5m+5}{4}$ elements in $\operatorname{Fix}_{NC^m(G_{32})}(\psi^p)$, which agrees with the limit in (8.13b).

In the case that $p = 10(m+1)/3$ or $p = 5(m+1)/3$, the relevant counting problem is (5.95). However, no element $(w_0; w_1, \dots, w_m) \in \operatorname{Fix}_{NC^m(G_{32})}(\psi^p)$ can be produced in this way since the counting problem imposes the restriction that $\ell_T(w_0) + \ell_T(w_1) + \dots + \ell_T(w_m)$ be divisible by 3, which is absurd. This is in agreement with the limit in (8.13e).

In the case that $p = 15(m+1)/2$ or $p = 5(m+1)/2$, we have found ten solutions to the counting problem (5.102) in (5.105), each of them giving rise to $(m+1)/2$ elements in $\operatorname{Fix}_{NC^m(G_{32})}(\psi^p)$, and we have found 25 solutions to the counting problem (5.104) in (5.106), each of them giving rise to $\binom{(m+1)/2}{2}$ elements in $\operatorname{Fix}_{NC^m(G_{32})}(\psi^p)$. Hence, we obtain $10\frac{m+1}{2} + 25\binom{(m+1)/2}{2} = \frac{(5m+5)(5m+3)}{8}$ elements in $\operatorname{Fix}_{NC^m(G_{32})}(\psi^p)$, which agrees with the limit in (8.13c).

Case G_{33}. The degrees are $4, 6, 10, 12, 18$, and hence we have

$$\operatorname{Cat}^m(G_{33};q) = \frac{[18m+18]_q\,[18m+12]_q\,[18m+10]_q\,[18m+6]_q\,[18m+4]_q}{[18]_q\,[12]_q\,[10]_q\,[6]_q\,[4]_q}.$$

Let ζ be a $18(m+1)$-th root of unity. The following cases on the right-hand side of (6.2) do occur:

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{33};q) = m+1, \quad \text{if } \zeta=\zeta_{18},\zeta_9, \tag{8.14a}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{33};q) = \tfrac{9m+9}{5}, \quad \text{if } \zeta=\zeta_{10},\zeta_5,\ 5\mid(m+1), \tag{8.14b}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{33};q) = \tfrac{(m+1)(3m+2)(3m+1)}{2}, \quad \text{if } \zeta=\zeta_6,\zeta_3, \tag{8.14c}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{33};q) = \operatorname{Cat}^m(G_{33}), \quad \text{if } \zeta=-1 \text{ or } \zeta=1, \tag{8.14d}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{33};q) = 0, \quad \text{otherwise.} \tag{8.14e}$$

We must now prove that the left-hand side of (6.2) in each case agrees with the values exhibited in (8.14). The only cases not covered by Lemmas 4 and 5 are the ones in (8.14b) and (8.14e). On the other hand, the only cases left to consider according to Remark 4 are the cases where $h_2 = 1$ and $m_2 = 5$, $h_2 = 2$ and $m_2 = 5$, $h_2 = 2$ and $m_2 = 4$, respectively $h_2 = m_2 = 2$. These correspond to the choices $p = 18(m+1)/5$, $p = 9(m+1)/5$, $p = 9(m+1)/4$, respectively $p = 9(m+1)/2$, out of which the first two belong to (8.14b), while the others belong to (8.14e).

In the case that $p = 18(m+1)/5$ or $p = 9(m+1)/5$, we have found nine solutions to the counting problem (5.115) in (5.116). Hence, we obtain $9\frac{m+1}{5} = \frac{9m+9}{5}$ elements in $\operatorname{Fix}_{NC^m(G_{33})}(\psi^p)$, which agrees with the limit in (8.14b).

If $p = 9(m+1)/4$, the computation at the end of Case G_{33} in Section 5 did not find any solutions, which is again in agreement with the limit in (8.13e).

In the case that $p = 9(m+1)/2$, the relevant counting problems are (5.122) and (5.124). However, no element $(w_0; w_1, \dots, w_m) \in \operatorname{Fix}_{NC^m(G_{33})}(\psi^p)$ can be produced in this way since the counting problem imposes the restriction that $\ell_T(w_0)+\ell_T(w_1)+\dots+\ell_T(w_m)$ be even, which is absurd. This is in agreement with the limit in (8.14e).

Case G_{34}. The degrees are $6, 12, 18, 24, 30, 42$, and hence we have

$$\operatorname{Cat}^m(G_{34};q) = \frac{[42m+42]_q\,[42m+30]_q\,[42m+24]_q}{[42]_q\,[30]_q\,[24]_q} \times \frac{[42m+18]_q\,[42m+12]_q\,[42m+6]_q}{[18]_q\,[12]_q\,[6]_q}.$$

Let ζ be a $42(m+1)$-th root of unity. The following cases on the right-hand side of (6.2) do occur:

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{34};q)=m+1,\quad \text{if } \zeta=\zeta_{42},\zeta_{21},\zeta_{14},\zeta_7, \tag{8.15a}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{34};q)=\operatorname{Cat}^m(G_{34}),\quad \text{if } \zeta=\zeta_6,\zeta_3,-1,1, \tag{8.15b}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(G_{34};q)=0,\quad \text{otherwise.} \tag{8.15c}$$

We must now prove that the left-hand side of (6.2) in each case agrees with the values exhibited in (8.15). The only case not covered by Lemmas 4 and 5 is the one in (8.15c). On the other hand, the only cases left to consider according to Remark 4 are the cases where $h_2=1$ and $m_2=5$, $h_2=2$ and $m_2=5$, $h_2=3$ and $m_2=5$, $h_2=6$ and $m_2=5$, $h_2=2$ and $m_2=4$, $h_2=6$ and $m_2=4$, $h_2=m_2=3$, $h_2=6$ and $m_2=3$, $h_2=m_2=2$, $h_2=6$ and $m_2=2$, respectively $h_2=6$ and $m_2=6$. These correspond to the choices $p=42(m+1)/5$, $p=21(m+1)/5$, $p=14(m+1)/5$, $p=7(m+1)/5$, $p=21(m+1)/4$, $p=7(m+1)/4$, $p=14(m+1)/3$, $p=7(m+1)/3$, $p=21(m+1)/2$, $p=7(m+1)/2$, respectively $p=7(m+1)/6$, all of which belong to (8.15c).

In the case that $p=42(m+1)/5$, $p=21(m+1)/5$, $p=14(m+1)/5$, or $p=7(m+1)/5$, the relevant counting problem is (5.128). However, no element $(w_0;w_1,\dots,w_m)\in \operatorname{Fix}_{NC^m(G_{34})}(\psi^p)$ can be produced in this way since the counting problem imposes the restriction that $\ell_T(w_0)+\ell_T(w_1)+\dots+\ell_T(w_m)$ be divisible by 5, which is absurd. This is in agreement with the limit in (8.15c).

In the case that $p=21(m+1)/4$ or $p=7(m+1)/4$, the relevant counting problem is (5.140). However, no element $(w_0;w_1,\dots,w_m)\in \operatorname{Fix}_{NC^m(G_{34})}(\psi^p)$ can be produced in this way since the counting problem imposes the restriction that $\ell_T(w_0)+\ell_T(w_1)+\dots+\ell_T(w_m)$ be divisible by 4, which is absurd. This is in agreement with the limit in (8.15c).

In the case that $p=14(m+1)/3$ or $p=7(m+1)/3$, the relevant counting problems are (5.146) and (5.148). However, the computations with the help of the *GAP* package `CHEVIE` [12] performed in Case G_{34} in Section 5 did not find any solutions for (5.146) or (5.148). This is in agreement with the limit in (8.15c).

In the case that $p=21(m+1)/2$, the relevant counting problems are (5.157), (5.158), and (5.159). However, the computations with the help of the *GAP* package `CHEVIE` [12] performed in Case G_{34} in Section 5 found no w_i with $\ell_T(w_i)=3$ in (5.157), and hence no solutions for (w_{i_1},w_{i_2}) with $\ell_T(w_{i_1})+\ell_T(w_{i_2})=3$ in (5.158), and no solutions for $(w_{i_1},w_{i_2},w_{i_3})$ in (5.159). This is in agreement with the limit in (8.15c).

If $p=7(m+1)/6$, the computation at the end of Case G_{34} in Section 5 did not find any solutions, which is also in agreement with the limit in (8.15c).

CASE $G_{35}=E_6$. The degrees are $2,5,6,8,9,12$, and hence we have

$$\operatorname{Cat}^m(E_6;q)=\frac{[12m+12]_q\,[12m+9]_q\,[12m+8]_q\,[12m+6]_q\,[12m+5]_q\,[12m+2]_q}{[12]_q\,[9]_q\,[8]_q\,[6]_q\,[5]_q\,[2]_q}.$$

Let ζ be a $12(m+1)$-th root of unity. The following cases on the right-hand side of (6.2) do occur:

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_6;q)=m+1,\quad \text{if } \zeta=\zeta_{12}, \tag{8.16a}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_6;q)=\tfrac{4m+4}{3},\quad \text{if } \zeta=\zeta_9,\ 3\mid(m+1), \tag{8.16b}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_6;q)=\tfrac{3m+3}{2},\quad \text{if } \zeta=\zeta_8,\ 2\mid(m+1), \tag{8.16c}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_6;q)=(m+1)(2m+1),\quad \text{if } \zeta=\zeta_6, \tag{8.16d}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_6;q)=\tfrac{(m+1)(3m+2)}{2},\quad \text{if } \zeta=\zeta_4, \tag{8.16e}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_6;q)=\tfrac{(m+1)(4m+3)(2m+1)}{3},\quad \text{if } \zeta=\zeta_3, \tag{8.16f}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_6;q)=\tfrac{(m+1)(3m+2)(2m+1)(6m+1)}{2},\quad \text{if } \zeta=-1, \tag{8.16g}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_6;q)=\operatorname{Cat}^m(E_6),\quad \text{if } \zeta=1, \tag{8.16h}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_6;q)=0,\quad \text{otherwise.} \tag{8.16i}$$

We must now prove that the left-hand side of (6.2) in each case agrees with the values exhibited in (8.16). The only cases not covered by Lemmas 4 and 5 are the ones in (8.16b), (8.16c), and (8.16i). On the other hand, the only cases left to consider according to Remark 4 are the cases where $h_2=1$ and $m_2=5$, respectively $h_2=2$ and $m_2=5$. These correspond to the choices $p=12(m+1)/5$, respectively $p=6(m+1)/5$, both of which belong to (8.16i).

In the case that $p=12(m+1)/5$, the relevant counting problem is (5.172). However, no element $(w_0;w_1,\dots,w_m)\in\operatorname{Fix}_{NC^m(E_6)}(\psi^p)$ can be produced in this way since the counting problem imposes the restriction that $\ell_T(w_0)+\ell_T(w_1)+\cdots+\ell_T(w_m)$ be divisible by 5, which is absurd. This is in agreement with the limit in (8.16i).

If $p=6(m+1)/5$, the computation at the end of Case E_6 in Section 5 did not find any solutions, which is also in agreement with the limit in (8.16i).

Case $G_{36}=E_7$. The degrees are $2,6,8,10,12,14,18$, and hence we have

$$\operatorname{Cat}^m(E_7;q)=\frac{[18m+18]_q\,[18m+14]_q\,[18m+12]_q}{[18]_q\,[14]_q\,[12]_q}\times\frac{[18m+10]_q\,[18m+8]_q\,[18m+6]_q\,[18m+2]_q}{[10]_q\,[8]_q\,[6]_q\,[2]_q}.$$

Let ζ be a $18(m+1)$-th root of unity. The following cases on the right-hand side of (6.2) do occur:

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_7;q)=m+1,\quad \text{if } \zeta=\zeta_{18},\zeta_9, \tag{8.17a}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_7;q)=\tfrac{9m+9}{7},\quad \text{if } \zeta=\zeta_{14},\zeta_7,\ 7\mid(m+1), \tag{8.17b}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_7;q)=\frac{(m+1)(3m+2)(3m+1)}{2},\quad \text{if } \zeta=\zeta_6,\zeta_3, \tag{8.17c}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_7;q)=\operatorname{Cat}^m(E_7),\quad \text{if } \zeta=-1 \text{ or } \zeta=1, \tag{8.17d}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_7;q)=0,\quad \text{otherwise.} \tag{8.17e}$$

We must now prove that the left-hand side of (6.2) in each case agrees with the values exhibited in (8.17). The only cases not covered by Lemmas 4 and 5 are the ones in (8.17b) and (8.17e). On the other hand, the only cases left to consider according to Remark 4 are the cases where $h_2=1$ and $m_2=7$, $h_2=2$ and $m_2=7$, $h_2=1$ and $m_2=5$, $h_2=2$ and $m_2=5$, $h_2=2$ and $m_2=4$, respectively $h_2=m_2=2$. These correspond to the choices $p=18(m+1)/7$, $p=9(m+1)/7$, $p=18(m+1)/5$, $p=9(m+1)/5$, $p=9(m+1)/4$, respectively $p=9(m+1)/2$, out of which the first two belong to (8.16b), and all others belong to (8.16i).

In the case that $p=18(m+1)/7$ or $p=9(m+1)/7$, we have found nine solutions to the counting problem (5.176) in (5.177). Hence, we obtain $9\frac{m+1}{7}=\frac{9m+9}{7}$ elements in $\operatorname{Fix}_{NC^m(E_7)}(\psi^p)$, which agrees with the limit in (8.17b).

In the case that $p=18(m+1)/5$ or $p=9(m+1)/5$, the relevant counting problem is (5.186). However, no element $(w_0;w_1,\dots,w_m)\in\operatorname{Fix}_{NC^m(E_7)}(\psi^p)$ can be produced in this way since the counting problem imposes the restriction that $\ell_T(w_0)+\ell_T(w_1)+\dots+\ell_T(w_m)$ be divisible by 5, which is absurd. This is in agreement with the limit in (8.17e).

In the case that $p=9(m+1)/4$, the relevant counting problem is (5.192). However, no element $(w_0;w_1,\dots,w_m)\in\operatorname{Fix}_{NC^m(E_7)}(\psi^p)$ can be produced in this way since the counting problem imposes the restriction that $\ell_T(w_0)+\ell_T(w_1)+\dots+\ell_T(w_m)$ be divisible by 4, which is absurd. This is again in agreement with the limit in (8.17e).

In the case that $p=9(m+1)/2$, the relevant counting problems are (5.195), (5.196), and (5.197). However, no element $(w_0;w_1,\dots,w_m)\in\operatorname{Fix}_{NC^m(E_7)}(\psi^p)$ can be produced in this way since the counting problem imposes the restriction that $\ell_T(w_0)+\ell_T(w_1)+\dots+\ell_T(w_m)$ be even, which is absurd. This is also in agreement with the limit in (8.17e).

CASE $G_{37}=E_8$. The degrees are $2,8,12,14,18,20,24,30$, and hence we have

$$\operatorname{Cat}^m(E_8;q)=\frac{[30m+30]_q\,[30m+24]_q\,[30m+20]_q\,[30m+18]_q}{[30]_q\,[24]_q\,[20]_q\,[18]_q}\\ \times\frac{[30m+14]_q\,[30m+12]_q\,[30m+8]_q\,[30m+2]_q}{[14]_q\,[12]_q\,[8]_q\,[2]_q}.$$

Let ζ be a $30(m+1)$-th root of unity. The following cases on the right-hand side of (6.2) do occur:

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q) = m+1, \quad \text{if } \zeta=\zeta_{30},\zeta_{15}, \tag{8.18a}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q) = \tfrac{5m+5}{4}, \quad \text{if } \zeta=\zeta_{24},\ 4\mid(m+1), \tag{8.18b}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q) = \tfrac{3m+3}{2}, \quad \text{if } \zeta=\zeta_{20},\ 2\mid(m+1), \tag{8.18c}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q) = \tfrac{(5m+5)(5m+3)}{8}, \quad \text{if } \zeta=\zeta_{12},\ 2\mid(m+1), \tag{8.18d}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q) = \tfrac{(m+1)(3m+2)}{2}, \quad \text{if } \zeta=\zeta_{10},\zeta_5, \tag{8.18e}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q) = \frac{(5m+5)(15m+7)}{16}, \quad \text{if } \zeta=\zeta_8,\ 4\mid(m+1), \tag{8.18f}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q) = \frac{(m+1)(5m+4)(5m+3)(5m+2)}{24}, \quad \text{if } \zeta=\zeta_6,\zeta_3, \tag{8.18g}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q) = \frac{(m+1)(5m+3)(15m+7)(15m+1)}{64}, \quad \text{if } \zeta=\zeta_4,\ 2\mid(m+1), \tag{8.18h}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q) = \operatorname{Cat}^m(E_8), \quad \text{if } \zeta=-1 \text{ or } \zeta=1, \tag{8.18i}$$

$$\lim_{q\to\zeta}\operatorname{Cat}^m(E_8;q) = 0, \quad \text{otherwise.} \tag{8.18j}$$

We must now prove that the left-hand side of (6.2) in each case agrees with the values exhibited in (8.18). The only cases not covered by Lemmas 4 and 5 are the ones in (8.18b), (8.18c), (8.18d), (8.18f), (8.18h), and (8.18j). On the other hand, the only cases left to consider according to Remark 4 are the cases where $h_2=2$ and $m_2=8$, $h_2=1$ and $m_2=7$, $h_2=2$ and $m_2=7$, $h_2=2$ and $m_2=4$, respectively $h_2=m_2=2$. These correspond to the choices $p=15(m+1)/8$, $p=30(m+1)/7$, $p=15(m+1)/7$, $p=15(m+1)/4$, respectively $15(m+1)/2$, out of which the first three belong to (8.18j), the fourth belongs to (8.18f), and the last belongs to (8.18h).

If $p=15(m+1)/8$, the relevant counting problem is (5.242). However, the computation at the end of Case E_8 in Section 5 did not find any solutions, which is in agreement with the limit in (8.18j). Hence, the left-hand side of (6.2) is equal to 0, as required.

In the case that $p=30(m+1)/7$ or $p=15(m+1)/7$, the relevant counting problem is (5.213). However, no element $(w_0;w_1,\dots,w_m)\in\operatorname{Fix}_{NC^m(E_8)}(\psi^p)$ can be produced in this way since the counting problem imposes the restriction that $\ell_T(w_0)+\ell_T(w_1)+\dots+\ell_T(w_m)$ be divisible by 7, which is absurd. This is also in agreement with the limit in (8.18j).

In the case that $p=15(m+1)/4$, the relevant counting problems are (5.227) and (5.228). We have found 45 solutions w_i to (5.227) of type A_1^2 in (5.229), and we have found 20 solutions w_i to (5.227) of type A_2 in (5.230), which implied 150 solutions for (w_{i_1},w_{i_2}) to (5.228). The first two give rise to to $(45+20)\frac{m+1}{4}=65\frac{m+1}{4}$ elements in $\operatorname{Fix}_{NC^m(E_8)}(\psi^p)$, while the third give rise to $150\binom{(m+1)/4}{2}$ elements in $\operatorname{Fix}_{NC^m(E_8)}(\psi^p)$. Hence, we obtain $65\frac{m+1}{4}+150\binom{(m+1)/4}{2}=\frac{(5m+5)(15m+7)}{16}$ elements in $\operatorname{Fix}_{NC^m(E_8)}(\psi^p)$, which agrees with the limit in (8.18f).

In the case that $p = 15(m+1)/2$, the relevant counting problems are (5.233), (5.234), (5.235), and (5.236). We have found 15 solutions w_i to (5.233) of type $A_1^2 * A_2$ in (5.237), we have found 45 solutions w_i to (5.233) of type $A_1 * A_3$ in (5.238), we have found 5 solutions w_i to (5.233) of type A_2^2 in (5.239), we have found 18 solutions w_i to (5.233) of type A_4 in (5.240), we have found 5 solutions w_i to (5.233) of type D_4 in (5.241), each giving rise to $(m+1)/2$ elements in $\operatorname{Fix}_{NC^m(E_8)}(\psi^p)$. Using the notation from there, these imply $2n_{3,1} + n_{2,2} = 2 \cdot 660 + 1195 = 2515$ solutions for (w_{i_1}, w_{i_2}) to (5.234) with $\ell_T(w_{i_1}) + \ell_T(w_{i_2}) = 4$, each giving rise to $\binom{(m+1)/2}{2}$ elements in $\operatorname{Fix}_{NC^m(E_8)}(\psi^p)$. They also imply $3n_{2,1,1} = 3 \cdot 2850 = 8550$ solutions for $(w_{i_1}, w_{i_2}, w_{i_3})$ to (5.235) with $\ell_T(w_{i_1}) + \ell_T(w_{i_2}) + \ell_T(w_{i_3}) = 4$, each giving rise to $\binom{(m+1)/2}{3}$ elements in $\operatorname{Fix}_{NC^m(E_8)}(\psi^p)$. Finally, they imply as well $n_{1,1,1,1} = 6750$ solutions for $(w_{i_1}, w_{i_2}, w_{i_3}, w_{i_4})$ to (5.236), each giving rise to $\binom{(m+1)/2}{4}$ elements in $\operatorname{Fix}_{NC^m(E_8)}(\psi^p)$.

In total, we obtain

$$(15+45+5+18+5)\frac{m+1}{2} + 2515\binom{(m+1)/2}{2} + 8550\binom{(m+1)/2}{3} + 6750\binom{(m+1)/2}{4} \\ = \frac{(m+1)(5m+3)(15m+7)(15m+1)}{64}$$

elements in $\operatorname{Fix}_{NC^m(E_8)}(\psi^p)$, which agrees with the limit in (8.18h).

REFERENCES

[1] D. Armstrong, *Generalized noncrossing partitions and combinatorics of Coxeter groups*, Mem. Amer. Math. Soc., vol. 202, no. 949, Amer. Math. Soc., Providence, R.I., 2009.

[2] C. A. Athanasiadis, *Generalized Catalan numbers, Weyl groups and arrangements of hyperplanes*, Bull. London Math. Soc. **36** (2004), 294–302.

[3] C. A. Athanasiadis, *On a refinement of the generalized Catalan numbers for Weyl groups*, Trans. Amer. Math. Soc. **357** (2005), 179–196.

[4] D. Bessis, *The dual braid monoid*, Ann. Sci. École Norm. Sup. (4) **36** (2003), 647–683.

[5] D. Bessis, *Finite complex reflection groups are $K(\pi, 1)$*, preprint, `ar`χ`iv:math/0610777`.

[6] D. Bessis and R. Corran, *Non-crossing partitions of type (e, e, r)*, Adv. Math. **202** (2006), 1–49.

[7] D. Bessis and V. Reiner, *Cyclic sieving and noncrossing partitions for complex reflection groups*, preprint, `ar`χ`iv:0701792`.

[8] T. Brady and C. Watt, *$K(\pi, 1)$'s for Artin groups of finite type*, Geom. Dedicata **94** (2002), 225–250.

[9] F. Chapoton, *Enumerative properties of generalized associahedra*, Séminaire Lotharingien Combin. **51** (2004), Article B51b, 16 pp.

[10] P. Edelman, *Chain enumeration and noncrossing partitions*, Discrete Math. **31** (1981), 171–180.

[11] S. Fomin and N. Reading, *Generalized cluster complexes and Coxeter combinatorics*, Int. Math. Res. Notices **44** (2005), 2709–2757.

[12] M. Geck, G. Hiss, F. Lübeck, G. Malle and G. Pfeiffer, *CHEVIE — a system for computing and processing generic character tables for finite groups of Lie type*, Appl. Algebra Engrg. Comm. Comput. **7** (1996), 175–210.

[13] J. E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge University Press, Cambridge, 1990.

[14] C. Krattenthaler, *The F-triangle of the generalised cluster complex*, in: Topics in Discrete Mathematics, dedicated to Jarik Nešetřil on the occasion of his 60th birthday, M. Klazar, J. Kratochvil, M. Loebl, J. Matoušek, R. Thomas and P. Valtr (eds.), Springer–Verlag, Berlin, New York, 2006, pp. 93–126.

[15] C. Krattenthaler, *The M-triangle of generalised non-crossing partitions for the types E_7 and E_8*, Séminaire Lotharingien Combin. **54** (2006), Article B54l, 34 pages.

[16] C. Krattenthaler, Non-crossing partitions on an annulus, in preparation.
[17] C. Krattenthaler and T. W. Müller, *Decomposition numbers for finite Coxeter groups and generalised non-crossing partitions*, Trans. Amer. Math. Soc. (to appear); `ar`χ`iv:0704.0199`.
[18] C. Krattenthaler and T. W. Müller, *Cyclic sieving for generalised non-crossing partitions associated to complex reflection groups of exceptional type*, preprint; `ar`χ`iv:1001.0028`.
[19] G. Kreweras, *Sur les partitions non croisées d'un cycle*, Discrete Math. **1** (1972), 333–350.
[20] G. I. Lehrer and J. Michel, *Invariant theory and eigenspaces for unitary reflection groups*, C. R. Math. Acad. Sci. Paris **336** (2003), 795–800.
[21] G. I. Lehrer and D. E. Taylor, *Unitary reflection groups*, Cambridge University Press, Cambridge, 2009.
[22] P. Orlik and L. Solomon, *Unitary reflection groups and cohomology*, Invent. Math. **59** (1980), 77–94.
[23] V. Reiner, D. Stanton and D. White, *The cyclic sieving phenomenon*, J. Combin. Theory Ser. A **108** (2004), 17–50.
[24] V. Ripoll, *Orbites d'Hurwitz des factorisations primitives d'un élément de Coxeter*, J. Algebra (to appear); `ar`χ`iv:0903.3604`.
[25] G. C. Shephard and J. A. Todd, *Finite unitary reflection groups*, Canad. J. Math. **6** (1954), 274–304.
[26] T. A. Springer, *Regular elements of finite reflection groups*, Invent. Math. **25** (1974), 159–198.
[27] J. R. Stembridge, Some hidden relations involving the ten symmetry classes of plane partitions, *J. Combin. Theory Ser. A* **68** (1994), 372–409.
[28] J. R. Stembridge, Canonical bases and self-evacuating tableaux, *Duke Math. J.* **82** (1996). 585–606,
[29] J. R. Stembridge, *coxeter*, *Maple* package for working with root systems and finite Coxeter groups; available at `http://www.math.lsa.umich.edu/~jrs`.

School of Mathematical Sciences, Queen Mary & Westfield College, University of London, Mile End Road, London E1 4NS, United Kingdom.
`http://www.maths.qmw.ac.uk/~twm/`.

Fakultät für Mathematik, Universität Wien, Nordbergstrasse 15, A-1090 Vienna, Austria. WWW: `http://www.mat.univie.ac.at/~kratt`.